

ZooKeys 602: 1–358 (2016)
doi: 10.3897/zookeys.602.8136
<http://zookeys.pensoft.net>

CATALOGUE

 A peer-reviewed open-access journal
ZooKeys
Launched to accelerate biodiversity research

A nomenclator of extant and fossil taxa of the Melanopsidae (Gastropoda, Cerithioidea)

Thomas A. Neubauer¹

¹ Geological-Paleontological Department, Natural History Museum Vienna, 1010 Vienna, Austria

Corresponding author: Thomas A. Neubauer (thomas.neubauer@nhm-wien.ac.at)

Academic editor: T. Backeljau | Received 15 February 2016 | Accepted 17 June 2016 | Published 5 July 2016

<http://zoobank.org/65EFA276-7345-4AC6-9B78-DBE7E98D6103>

Citation: Neubauer TA (2016) A nomenclator of extant and fossil taxa of the Melanopsidae (Gastropoda, Cerithioidea). ZooKeys 602: 1–358. doi: 10.3897/zookeys.602.8136

Abstract

This nomenclator provides details on all published names in the family-, genus-, and species-group, as well as for a few infrasubspecific names introduced for, or attributed to, the family Melanopsidae. It includes nomenclaturally valid names, as well as junior homonyms, junior objective synonyms, nomina nuda, common incorrect subsequent spellings, and as far as possible discussion on the current status in taxonomy. The catalogue encompasses three family-group names, 79 genus-group names, and 1381 species-group names. All of them are given in their original combination and spelling (except mandatory corrections requested by the Code), along with their original source. For each family- and genus-group name, the original classification and the type genus and type species, respectively, are given. Data provided for species-group taxa are type locality, type horizon (for fossil taxa), and type specimens, as far as available.

Keywords

Non-marine snails, catalogue, nomenclature, taxonomy, www-references

Table of contents

Introduction.....	2
Part I: Supraspecific taxa in Melanopsidae	3
Part IA. Family-group names.....	3
Part IB. Genus-group names	4
Part II: Species-group and infrasubspecific names.....	18
<i>Melanopsis</i> species introduced by André Etienne Justin Pascal Joseph Francois d'Audebard de Férussac.....	19
Validity of names mentioned in Férussac's and Deshayes' works	20
Names introduced by Jean-Pierre Sylvestre de Grateloup in 1828 and 1838	23
The taxonomic concepts applied by Jules René Bourguignat, Paul Pallary, Jean-Marie Pérès and Ferdinand Starmühlner	23
Note on the information provided	24
Nomenclatural notes	25
Acknowledgements.....	316
References	317

Introduction

The family Melanopsidae (Caenogastropoda: Cerithioidea) is one of the most diverse groups of non-marine gastropods in Earth history (Strong et al. 2008). Today, the family occurs in southern to eastern Europe, northern Africa, and the Middle East, as well as New Zealand and some south Pacific islands (Bănărescu 1990, Glaubrecht 1993, 1996, Altaba 1998, Strong et al. 2008). Its record dates back at least into the late Cretaceous of Europe (Bandel 2000, Neubauer et al. 2016a). While the fossil record of New Zealand species is fairly restricted, the record for Europe, northern Africa and the Middle East yielded many hundreds of species-group taxa. Early species were pure brackish-water dwellers, which they remained until the early Miocene, when they started to conquer freshwater (Glaubrecht 1996, Neubauer et al. 2016a). This change in life style coincided with a series of adaptive radiations in the middle and late Miocene, producing hundreds of species, and forming the basis for the modern clades and present diversity and distribution (Neubauer et al. 2016a).

The first detailed listing of names of living “Melanidae” – at that time the Melanopsidae were considered a subfamily of that group – was published by Brot (1874–1879). Soon later, Bourguignat (1884) followed with his “Histoire des mélaniens du système européen”, in which he described numerous new species, but few of them are currently still accepted. Pallary (1926a, b) provided a list of all published names of fossil and recent Melanopsidae, yet without any comments. Wenz (1929) summarized all Cenozoic (Paleocene–Pleistocene) names and gave extensive synonymy lists. Besides, there are papers on the fossil and recent Melanopsidae of Italy (Pantaneli 1886b) and Spain (Pallary 1924, Azpeitia Moros 1929). In addi-

tion, Pallary (1916a, b, 1920, 1925) provided a detailed account of nomenclatural mistakes concerning this family and introduced a great many of replacement names for existing homonyms.

A comprehensive annotated list of melanopsid names is, however, entirely missing. This catalogue presents information for all published names in the family-, genus- and species-group, as well as for a few infrasubspecific names. Discussed are nomenclaturally valid, invalid (e.g., junior objective synonyms, junior homonyms) and unavailable names (e.g., nomina nuda, misspellings), following the rules of the fourth edition of the International Code of Zoological Nomenclature (ICZN 1999, henceforth referred to as “the Code”), incorporating later added amendments (see <http://www.nhm.ac.uk/hosted-sites/iczn/code/index.jsp>). To facilitate comparison, the general outline of the present work follows the excellent nomenclator of Valvatidae by Haszprunar (2014).

Part I: Supraspecific taxa in Melanopsidae

The supraspecific phylogenetic relationships within the Melanopsidae are poorly resolved (Strong et al. 2011). Most of the subfamilies, genera and subgenera were established for particular morphological characteristics, most of which have been later shown to be highly variable and, moreover, polyphyletic (e.g., Neubauer et al. 2014a). Few of them are currently used in taxonomy.

The fossil Stomatopsinae Stache, 1889 (type genus: *Stomatopsis* Stache in Sandberger, 1871), originally introduced as subfamily of the “Melaniidae” (although Stache 1889 used the suffix “-idae”) but considered as a junior synonym of the Melanopsidae by Bouchet and Rocroi (2005: 248), are not included here. Their systematic position is highly uncertain and requires a more detailed re-examination.

Part IA. Family-group names

AMPHIMELANIINAE P. Fischer & Crosse, 1891

Original source. Fischer and Crosse 1880–1902: 312.

Original classification. Subfamily of Melaniidae.

Type genus. *Amphimelania* P. Fischer, 1885 (junior objective synonym of *Holandriana* Bourguignat, 1884).

Remarks. Considered a junior synonym of the Melanopsidae by Bouchet and Rocroi (2005: 248).

FAGOTIINAE Starobogatov in Starobogatov et al., 1992

Original source. Starobogatov et al. 1992: 58.

Original classification. Subfamily of Melanopsidae.

Type genus. *Fagotia* Bourguignat, 1884.

Remarks. Considered a junior synonym of the Melanopsidae by Bouchet and Rocroi (2005: 248).

MELANOPSIDAE H. Adams & A. Adams, 1854

Original source. Adams and Adams 1853–1858: 309.

Original classification. Subfamily of Melaniidae.

Type genus. *Melanopsis* Férussac in Férussac & Férussac, 1807.

Part IB. Genus-group names

In the following list, not all genus-group names are accompanied by a type species. Especially Bourguignat, who introduced the greatest number of melanopsid (sub) genera, rarely designated type species. Before 1931, a type species fixation was not a requirement for being available (ICZN 1999, Art. 12, 13.3). Original classifications of genus-level taxa are omitted for genera that were introduced without clear family classification. Purely fossil genera are marked by a dagger.

Aciculariana Bourguignat, 1884

Original source. Bourguignat 1884: 51.

Original classification. Subgenus of *Microcolpia*.

Remarks. Bourguignat (1884) apparently intended to base this subgenus on *M. acicularis*. If originally or subsequently designated as type species, *Aciculariana* would be an objective synonym of *Microcolpia*.

Acroxiana Bourguignat, 1884

Original source. Bourguignat 1884: 32.

Original classification. Subgenus of *Fagotia*.

Amphimelania P. Fischer, 1885 [invalid]

Original source. Fischer and Crosse 1880–1902: 701.

Original classification. Subgenus of *Melania*.

Type species. *Melania holandrii* Pfeiffer, 1828, by original designation.

Remarks. Replacement name for *Melanella* Swainson, 1840, non Bowdich, 1822. Junior objective synonym of *Holandriana* Bourguignat, 1884, with the same type species.

***Asmena* Gistel, 1848 [invalid]**

Original source. Gistel 1848: xi.

Remarks. Unnecessary substitute name for *Melanopsis* Férussac in Férussac & Férussac, 1807.

† *Battistiana* Pallary, 1920 [invalid]

Original source. Pallary 1920b: 108.

Original classification. Genus of Melanopsidae.

Type species. † *Buccinum fosile* Gmelin, 1791, by typification of replaced name.

Remarks. Established as a replacement name for *Pannonia* Pallary, 1916, wrongly assumed by Pallary (1920b) to be preoccupied by *Pannona* Lörenthey, 1902, and in fact a junior homonym of *Pannonia* Dollfus, 1912.

***Belusiana* Bourguignat, 1884**

Original source. Bourguignat 1884: 75.

Original classification. Subgenus of *Melanopsis*.

† *Boistelia* Cossmann, 1909

Original source. Cossmann 1909: 183.

Original classification. Section of *Melanopsis*.

Type species. † *Melanoptychia paradoxa* Brusina, 1892, by original designation.

***Buccinoidiana* Bourguignat, 1884**

Original source. Bourguignat 1884: 73.

Original classification. Subgenus of *Melanopsis*.

***Caledomelanopsis* Germain, 1934 [unavailable]**

Original source. Germain 1934: 146.

Original classification. Genus of Melanopsidae.

Remarks. Germain (1934) neither provided a diagnosis, nor did he indicate a type species.

† *Calodiona* Stefanescu, 1896

Original source. Stefanescu 1896: 126, 131.

Original classification. Subgenus of *Melanopsis*.

Type species. † *Melanopsis bergeroni* Stefanescu, 1896, by monotypy.

† *Campylostylus* Sandberger, 1870

Original source. Sandberger 1870–1875: 90.

Original classification. Subgenus of *Melanopsis*.

Type species. † *Melanopsis galloprovincialis* Mathéron, 1843, by monotypy.

† *Canthidomus* Swainson, 1840

Original source. Swainson 1840: 202, 342.

Original classification. Subgenus of *Melanopsis*.

Type species. † *Melanopsis bouei* Férussac, 1823, by subsequent designation by Herrmannsen (1846: 167).

Cariosiana Bourguignat, 1877

Original source. Bourguignat 1877: 88.

Original classification. Subgenus of *Melanopsis*.

Type species. *Murex cariosus* Linnaeus, 1767, by original designation.

Ceneona Gistel, 1848 [invalid]

Original source. Gistel 1848: 169.

Remarks. Unnecessary substitute name for *Melanopsis* Férussac in Férussac & Férussac, 1807.

Charpentieriana Bourguignat, 1884

Original source. Bourguignat 1884: 75.

Original classification. Subgenus of *Melanopsis*.

***Costatiana* Bourguignat, 1884**

Original source. Bourguignat 1884: 76.

Original classification. Subgenus of *Melanopsis*.

***Couphaniana* Bourguignat, 1884**

Original source. Bourguignat 1884: 74.

Original classification. Subgenus of *Melanopsis*.

Type species. *Melanopsis coupha* Bourguignat, 1884, by monotypy.

***Crassiana* Bourguignat, 1884 [invalid]**

Original source. Bourguignat 1884: 8.

Original classification. Subgenus of *Melanella*.

Remarks. Junior homonym of *Crassiana* Servain, 1882 (Bivalvia, Unionidae).

† *Duabiana* Starobogatov & Anistratenko in Anistratenko, 1993

Original source. Anistratenko 1993: 69, textfig. 1.

Original classification. Subgenus of *Melanopsis*.

Type species. † *Melanopsis cylindrica* Anistratenko, 1993, by original designation.

***Dufouriana* Bourguignat, 1884**

Original source. Bourguignat 1884: 75.

Original classification. Subgenus of *Melanopsis*.

***Esperiana* Bourguignat, 1877**

Original source. Bourguignat 1877: 88.

Original classification. Subgenus of *Melanopsis*.

Type species. *Melanopsis esperi* Férussac, 1823, by original designation.

***Eumorphiana* Bourguignat, 1884**

Original source. Bourguignat 1884: 76.

Original classification. Subgenus of *Melanopsis*.

***Fagotia* Bourguignat, 1884 [invalid]**

Original source. Bourguignat 1884: 30.

Original classification. Subgenus of *Fagotia*.

Type species. *Melanopsis esperi* Férussac, 1823, by subsequent designation by Wenz (1939: 690).

Remarks. Wenz (1939) declared that *Melanopsis esperi* was the type species by monotypy, but 22 nominal species were originally included in *Fagotia*. Junior objective synonym of *Esperiana*.

***Fagotiana* Bourguignat, 1884**

Original source. Bourguignat 1884: 7.

Original classification. Subgenus of *Melanella*.

Type species. *Melanella fagotiana* Bourguignat, 1884, by tautonymy.

***Graellsiana* Bourguignat, 1884**

Original source. Bourguignat 1884: 75.

Original classification. Subgenus of *Melanopsis*.

† *Handmannia* Cossmann, 1889 [invalid]

Original source. Cossmann 1889: 1101.

Type species. † *Melanopsis pygmaea* Hörnes, 1856, by original designation.

Remarks. Established as a substitute name for *Homalia* Handmann, 1887, which Cossmann (1899), despite being aware of the difference, treated as a junior homonym of “*Homala* Schumacher” (actually *Omala* Schumacher, 1817, unjustified emendation to “*Homala*” by Agassiz 1847).

***Holandriana* Bourguignat, 1884**

Original source. Bourguignat 1884: 7.

Original classification. Subgenus of *Melanella*.

Type species. *Melania holandrii* Pfeiffer, 1828, by subsequent designation by Welter-Schultes (2012: 36).

† *Homalia* Handmann, 1887

Original source. Handmann 1887: 12.

Original classification. Subgenus of *Melanopsis*.

† *Hyphantria* Handmann, 1887 [invalid]

Original source. Handmann 1887: 37.

Original classification. Subgenus of *Melanopsis*.

Remarks. Junior homonym of *Hyphantria* Harris, 1841 (Lepidoptera).

Laevigatiana Bourguignat, 1884

Original source. Bourguignat 1884: 8.

Original classification. Subgenus of *Melanella*.

Letourneuxiana Bourguignat, 1884 [invalid]

Original source. Bourguignat 1884: 32.

Original classification. Subgenus of *Fagotia*.

Remarks. Bourguignat (1884) introduced three different divisions called *Letourneuxiana*, one for *Fagotia*, one for *Melanella* and one for *Melanopsis*. All are junior homonyms of *Letourneuxiana* Silva e Castro, 1883 (Unionidae).

Letourneuxiana Bourguignat, 1884 [invalid]

Original source. Bourguignat 1884: 8.

Original classification. Subgenus of *Melanella*.

Remarks. Junior homonym of *Letourneuxiana* Silva e Castro, 1883 (Unionidae).

Letourneuxiana Bourguignat, 1884 [invalid]

Original source. Bourguignat 1884: 75.

Original classification. Subgenus of *Melanopsis*.

Type species. *Melanopsis letourneuxi* Bourguignat, 1884, by monotypy.

Remarks. Junior homonym of *Letourneuxiana* Silva e Castro, 1883 (Unionidae).

***Locardiana* Bourguignat, 1884**

Original source. Bourguignat 1884: 32.

Original classification. Subgenus of *Fagotia*.

Type species. *Fagotia locardiana* Bourguignat, 1884, by tautonymy.

***Lorcaniana* Bourguignat, 1884**

Original source. Bourguignat 1884: 74.

Original classification. Subgenus of *Melanopsis*.

***Lortetiana* Bourguignat, 1884**

Original source. Bourguignat 1884: 76.

Original classification. Subgenus of *Melanopsis*.

Type species. *Melanopsis lortetiana* Locard, 1883, by tautonymy.

***Lyrcea* H. Adams & A. Adams, 1854**

Original source. Adams and Adams 1853–1858: 310.

Original classification. Subgenus of *Melanopsis*.

Type species. *Melanopsis dufourii* Férussac, 1822, by subsequent designation by Cossmann (1909: 174).

Remarks. “*Lyrcea*” as mentioned in Wenz (1929: 2647) and many other authors is an incorrect subsequent spelling.

† *Macrospira* Sandberger, 1872 [invalid]

Original source. Sandberger 1870–1875: 222, 248.

Original classification. Subgenus of *Melanopsis*.

Remarks. Junior homonym of *Macrospira* Guilding in Swainson, 1840. Rovereto (1899: 109) introduced *Stilospirula* as replacement name.

***Maresiana* Bourguignat, 1884**

Original source. Bourguignat 1884: 77.

Original classification. Subgenus of *Melanopsis*.

† *Martinia* Handmann, 1887 [invalid]

Original source. Handmann 1887: 19.

Original classification. Subgenus of *Melanopsis*.

Type species. † *Melanopsis martiniana* Férussac, 1823 [objective synonym of *M. fossilis* (Gmelin, 1791)], by subsequent designation by Cossmann (1909: 174).

Remarks. Junior homonym of *Martinia* M'Coy in M'Coy & Griffith, 1844 (Brachiopoda).

† *Megalonoda* Kollmann, 1984

Original source. Kollmann 1984: 56.

Original classification. Genus of Melanopsidae.

Type species. † *Purpuroidea reussi* Hörnes, 1856, by original designation.

Melafusus Swainson, 1840

Original source. Swainson 1840: 201, 341.

Original classification. Subgenus of *Melanopsis*.

Remarks. Briefly described but no species included.

Melanella Swainson, 1840 [invalid]

Original source. Swainson 1840: 199, 341.

Original classification. Subgenus of *Melania*.

Remarks. No originally included species. Junior homonym of *Melanella* Bowdich, 1822. Fischer (1885: 701) introduced *Amphimelania* as replacement name, which is invalid, too, because it is a junior objective synonym of *Holandriana*, with the same type species.

Melanithes Swainson, 1840

Original source. Swainson 1840: 202, 341.

Original classification. Subgenus of *Melanopsis*.

Remarks. Based on some of Férussac's (1823) fossil (Eocene to Pliocene) *Melanopsis* species but no type species was designated.

***Melanopsis* J.B.L.d'A. de Férussac in J.B.L. d'A. de Férussac & A.E.J.P.J.F. d'A. de Férussac, 1807**

Original source. Férussac and Férussac 1807: 70.

Type species. *Melania costata* Olivier, 1804, by subsequent designation by Gray (1847: 153).

Remarks. Correct authority is denoted on p. xii of Férussac and Férussac (1807).

† *Melanoptychia* Neumayr, 1880

Original source. Neumayr 1880c: 480.

Original classification. Genus of Melanopsidae.

Type species. † *Melanoptychia bittneri* Neumayr, 1880, by subsequent designation by Cossmann (1909: 182).

† *Melanosteira* Andreae, 1893

Original source. Andreae 1893: 172.

Original classification. Genus of Melanopsidae.

Type species. † *Melanopsis aetolica* Neumayr, 1876, by subsequent designation by Cossmann (1909: 179).

Remarks. The name is not available from Oppenheim (1891), who used the vernacular “Melanosteiren”. In a book review of that paper, Andreae (1893) Latinized the name and made it thereby available.

† *Melanostira* Cossmann, 1909 [invalid]

Original source. Cossmann 1909: 179.

Remarks. Unjustified emendation and therefore junior objective synonym of *Melanosteira* Andreae, 1893.

***Mesopotamia* Pallary, 1939 [unavailable]**

Original source. Pallary 1939: 98.

Original classification. Subgenus of *Melanopsis*.

Remarks. Unavailable because no type species was designated (Art. 13.3).

***Microcolpia* Bourguignat, 1884**

Original source. Bourguignat 1884: 49.

Original classification. Genus of Melaniidae (note that Bourguignat actually only gave the vernacular “Mélaniens”).

Type species. *Melanopsis acicularis* Férussac, 1823, by subsequent designation by Cossmann (1909: 159).

Remarks. “*Microcalpia*” as mentioned in Fischer (1886: 705) is an incorrect subsequent spelling.

***Mingrelicana* Bourguignat, 1884**

Original source. Bourguignat 1884: 74.

Original classification. Subgenus of *Melanopsis*.

Type species. *Melanopsis mingrelica* Mousson, 1863, by subsequent designation by Izzatullaev and Starobogatov (1984: 1476).

***Myosotidiana* Bourguignat, 1884**

Original source. Bourguignat 1884: 74.

Original classification. Subgenus of *Melanopsis*.

Type species. *Melanopsis myosotidaea* Bourguignat, 1884, by monotypy.

***Nodosiana* Bourguignat, 1884**

Original source. Bourguignat 1884: 76.

Original classification. Subgenus of *Melanopsis*.

Type species. *Melanopsis nodosa* Férussac, 1822, by monotypy.

***Olivieriana* Bourguignat, 1884**

Original source. Bourguignat 1884: 74.

Original classification. Subgenus of *Melanopsis*.

***Pakaurangia* Finlay, 1926**

Original source. Finlay 1926: 380, 476.

Original classification. Subgenus of *Zemelanopsis*.

Type species. *Melanopsis waitaraensis* Marwick, 1926, by original designation.

Remarks. Finlay (1926) included also another species, the fossil *Coptochetus zelandicus* Marshall, 1918 (p. 265, pl. 22, fig. 13), in this subgenus. That species is, however, not a Melanopsidae but belongs to the marine genus *Exilia* Conrad, 1860 after Beu and Maxwell (1990: 414) and is consequently not treated in this catalogue. Finally, Bouchet and Poppe (1995: 503) synonymized *Pakaurangia* with the marine deep-water genus *Calliotectum* (Volutidae).

† *Pannonia* Pallary, 1916 [invalid]

Original source. Pallary 1916: 76.

Original classification. Subgenus of *Melanopsis*.

Type species. † *Buccinum fossile* Gmelin, 1791, by original designation.

Remarks. Replacement name for *Martinia* Handmann, 1887, non M'Coy in M'Coy & Griffith, 1844. Junior homonym of *Pannonia* Dollfus, 1912 (see *Battistiana*).

Parreyssiana Bourguignat, 1884

Original source. Bourguignat 1884: 76.

Original classification. Subgenus of *Melanopsis*.

Type species. *Melanopsis parreyssi* Philippi, 1847, by monotypy.

† *Pauluccia* Brusina, 1902

Original source. Brusina 1902: viii.

Original classification. Subgenus of *Melanopsis*.

Pechaudiana Bourguignat, 1884

Original source. Bourguignat 1884: 77.

Original classification. Subgenus of *Melanopsis*.

Potamactebiana Bourguignat, 1884

Original source. Bourguignat 1884: 52.

Original classification. Subgenus of *Microcolpia*.

Type species. *Melanopsis potamactebia* Bourguignat, 1870, by subsequent designation by Anistratenko and Anistratenko (2001: 150).

***Praeclariana* Bourguignat, 1884**

Original source. Bourguignat 1884: 51.

Original classification. Subgenus of *Microcolpia*.

Type species. *Microcolpia praeclara* Bourguignat, 1884, by monotypy.

***Praemorsiana* Bourguignat, 1877**

Original source. Bourguignat 1877: 88.

Original classification. Subgenus of *Melanopsis*.

Type species. *Buccinum praemorsum* Linnaeus, 1758, by original designation.

***Pseudhemisus* Nevill, 1884 [invalid]**

Original source. Nevill 1884: 214.

Original classification. Subgenus of *Melanopsis*.

Type species. *Melanopsis esperi* Férussac, 1823, by original designation.

Remarks. Junior objective synonym of *Esperiana* Bourguignat, 1876, with the same type species.

† *Pseudofagotia* Anistratenko, 1993

Original source. Anistratenko 1993: 72.

Original classification. Genus of Fagotiinae.

Type species. † *Pseudofagotia lineata* Anistratenko, 1993, by original designation.

***Pyramidaliana* Bourguignat, 1884**

Original source. Bourguignat 1884: 51.

Original classification. Subgenus of *Microcolpia*.

† *Sasykiana* Gozhik in Gozhik & Datsenko, 2007

Original source. Gozhik and Datsenko 2007: 98.

Original classification. Subgenus of *Fagotia*.

Type species. † *Fagotia (Sasykiana) plena* Gozhik in Gozhik & Datsenko, 2007, by original designation.

***Saulcyana* Bourguignat, 1884**

Original source. Bourguignat 1884: 75.

Original classification. Subgenus of *Melanopsis*.

***Scalariana* Bourguignat, 1884**

Original source. Bourguignat 1884: 74.

Original classification. Subgenus of *Melanopsis*.

***Seignettiana* Bourguignat, 1884**

Original source. Bourguignat 1884: 74.

Original classification. Subgenus of *Melanopsis*.

***Servainiana* Bourguignat, 1884**

Original source. Bourguignat 1884: 51.

Original classification. Subgenus of *Microcolpia*.

***Sesteriana* Bourguignat, 1884**

Original source. Bourguignat 1884: 75.

Original classification. Subgenus of *Melanopsis*.

***Sistaniana* Izzatullaev & Starobogatov, 1984**

Original source. Izzatullaev and Starobogatov 1984: 1476.

Original classification. Subgenus of *Melanopsis*.

Type species. *Melanopsis sistanica* Izzatullaev & Starobogatov, 1984, by original designation.

***Smendovia* Tournouër, 1882**

Original source. Tournouër 1882: 59.

Original classification. Genus of Melanopsinae.

Type species. † *Melanopsis thomasi* Tournouër, 1877, by original designation.

***Speciosiana* Bourguignat, 1884**

Original source. Bourguignat 1884: 8.

Original classification. Subgenus of *Melanella*.

Type species. *Melanella speciosa* Bourguignat, 1884, by monotypy.

† *Spiridionia* Cossmann, 1909

Original source. Cossmann 1909: 178.

Original classification. Subgenus of *Melanopsis*.

Type species. † *Melanopsis austriaca* Handmann, 1882, by original designation.

Remarks. Replacement name for *Hyphantria* Handmann, 1887, non Harris, 1841 (Lepidoptera).

† *Stilospirula* Rovereto, 1899

Original source. Rovereto 1899: 109.

Original classification. Genus of “Melaniidae”.

Type species. † *Melanopsis proboscidea* Deshayes, 1862, by subsequent designation by Cossmann (1909: 172; note that Cossmann made an unjustified emendation by changing the name to “*Stylospirula*”).

Remarks. Replacement name for *Macrospira* Sandberger, 1872, non Guilding in Swainson, 1840.

† *Stylospirula* Cossmann, 1899 [invalid]

Original source. Cossmann 1899: 147.

Remarks. Unjustified emendation and therefore junior objective synonym of *Stylospirula* Rovereto, 1899.

† *Turripontica* Anistratenko, 1993

Original source. Anistratenko 1993: 75.

Original classification. Genus of Fagotiinae.

Type species. † *Turripontica aciculina* Anistratenko, 1993, by original designation.

***Villeserriana* Bourguignat, 1884**

Original source. Bourguignat 1884: 51.

Original classification. Subgenus of *Microcolpia*.

Type species. *Microcolpia villeserriana* Bourguignat, 1884, by tautonymy.

***Zemelanopsis* Finlay, 1926**

Original source. Finlay 1926: 380, 476.

Type species. *Melanopsis trifasciata* Gray, 1843, by original designation.

Part II: Species-group and infrasubspecific names

The number of living melanopsid species-group taxa accepted in taxonomy today ranges between 25 and 50 (Strong et al. 2008); the IUCN Red List includes 36 species. Fossil species are far more numerous, with over 450 accepted names (Neubauer et al. 2016a). Beyond that, a plethora of homonyms and synonyms have been described in the Melanopsidae. The catalogue includes taxa initially described in this family and subsequently classified in other families and vice versa. It comprises a total of 1381 names, of which 71 are unavailable (mostly nomina nuda), 252 are available but invalid (junior homonyms and junior objective synonyms) and 15 are unresolved (requiring the action of a First Reviser).

Not treated by the present work are taxa not described for, and currently not considered to belong to, the Melanopsidae but classified therein previously. In the 19th century, many species introduced in the group that nowadays is understood as the family Melanopsidae, especially regarding varieties of *Holandriana holandrii* (Pfeiffer, 1828), have been classified within the genus *Melania* Lamarck, 1799 (e.g., Rossmässler 1838–1844, Brusina 1866, Kobelt 1881, Westerlund 1886, 1898). In order to avoid mixing with “real” *Melania* (which is a junior objective synonym of *Thiara* Röding, 1798, the type species of the family Thiaridae), only such *Melania* species-group taxa attributed to the *Holandriana holandrii* species-complex, or those that have been claimed to be related to that group, are included herein. Further taxa that are not included are: species introduced within the genera *Faunus* Montfort, 1810, *Hemisinus* Swainson, 1840, *Dicolpus* Philippi, 1887 [invalid: junior homonym of *Dicolpus* Gerstaecker, 1884 (Insecta)], *Loxotrema* Gabb, 1868, and *Boggsia* Olsson, 1929 that have been occasionally classified in the Melanopsidae (e.g., Brot 1874–1879, Cossmann 1909, Squires 1998, 1999, Squires and Saul 1997) but are not considered to belong there today; the *Bullia* (*Bullioopsis*) species described by Conrad (1830, 1862) from middle Miocene deposits of Maryland, USA (*B. integra* Conrad, 1862, *B. marylandica* Conrad, 1862, and *B. quadrata* Conrad, 1830) that he later considered melanopsids (Conrad 1866) but which belong to the marine Nassariidae after Allmon (1990); Pale-

ocene species of the Stomatopsinae Stache, 1889 (genera *Stomatopsis* Stache in Sandberger, 1871, *Stomatopsella* Stache, 1889, and *Megastomopsis* Stache, 1889) which was considered a subfamily of Melanopsidae by Bouchet & Rocroi (2005) but probably does not belong there; species introduced in the genus *Coptostylus* Sandberger, 1872, which has occasionally been classified within the Melanopsidae, but is currently assigned to the Thiaridae (Pacaud and Le Renard 1995: 156).

Basic sources used for the present assembly were the catalogues of Brot (1874–1879), Bourguignat (1884), Pallary (1926a, b) and Wenz (1929), as well as internet resources such as the Biodiversity Heritage Library (<http://biodiversitylibrary.org/>), the Global Names Index (<http://gni.globalnames.org/>) and the Worldwide Mollusc Species Data Base (<http://www.bagniliggia.it/WMSD/WMSDhome.htm>). However, many names listed in those online repositories are obvious misspellings deriving from automatic digitization procedures that have not been critically reviewed (compare discussion in Haszprunar 2014). Such errors are not included here because they do not come from the published literature. Some of the names currently listed in those databases could neither be traced back to any publication, nor could they be linked to an evident misspelling. These names are on purpose excluded from the present paper to avoid introducing potential misspellings or nomina nuda into the literature. Nevertheless, I will be grateful to receive information on names presently not included here, as well as corrections, explanations and additions.

Some of the publications consulted for this work display problematic cases, regarding their actual publication dates or the nomenclatural or taxonomic concepts applied therein. These issues require careful examinations, which are provided below.

***Melanopsis* species introduced by André Etienne Justin Pascal Joseph Francois d'Audebard de Férussac**

Férussac described several new species of *Melanopsis* in his “Histoire naturelle générale et particulière des mollusques terrestres et fluviatiles”, as well as in the “Monographie des espèces vivantes et fossiles du genre mélanopside”. The problem is to determine the exact publication dates of those works and, therefore, when and in which work which species was described.

The “Histoire naturelle” is an extremely comprehensive work, containing two volumes of text and 210 plates, which were published between March 1819 and August 1851 in 35 “livraisons”. It has been subject of intensive debates regarding the exact publication dates – for details see Kennard (1942a–c). The work was commenced by Férussac, who issued the first 28 livraisons between 1819 and 1832. After his death in 1836, Gérard Paul Deshayes finished the work by adding another seven livraisons between 1839 and 1851, including new explanations for the plates. The work includes two plates of “Mélanopsides fossiles”. The first plate was published in livraison 15 on 13 April 1822, but its captions came along with livraison 17, which was published on 2 November 1822 (Kennard 1842b: 106). The second plate appeared in livraison 21 on 27 September 1823 including its captions (Kennard 1842a: 16).

To increase confusion, there is a disagreement between the original captions provided by Férussac and the ones supplied by Deshayes with livraison 29 in 1839 (Table 1; see also Kennard 1842b: 106). Deshayes made several corrections and synonymizations and even introduced two new names in the altered captions for plate 2, i.e., *Melanopsis subcarinata* and *Melanopsis nodosa* var. *longa*. While the original captions for plate 2 are available to me, I could not find the captions for plate 1. Both plates were also published par for par in Férussac's (1823) monograph on the Melanopsidae. The captions for plate 2 are the same in both works (except minor differences regarding the details provided), so we can assume the same for plate 1. However, the exact wording of the original captions of 1822 needs to be seen in the future, in order to finally ascertain the availability of the associated names. A different problem appears for plate 2. While the year of publication of the species illustrated there is undoubtedly 1823, the original publication itself is uncertain – either the livraison 21 of the “Histoire naturelle” [27 September 1823] or the “Monographie” [exact date unknown]. This matter could unfortunately not be solved.

Validity of names mentioned in Férussac's and Deshayes' works

Apart from the uncertainties detailed above, the inconsistent formatting of Férussac's (1823) monograph led later authors to adopt names as valid that were certainly not intended as such by Férussac himself. He introduced several new *Melanopsis* species and varieties, latter of which he denoted by Greek letters followed by Latin descriptions. Quite often, he used the same Latin words to characterize different varieties and apparently did not mean to introduce available names (e.g., Férussac 1823: 154). New names (or new combinations) were always denoted by him with “nobis”. Table 2 summarizes the available species-group names and unnamed varieties introduced and described by Férussac (1823).

The variety *M. buccinoidea* var. “γ) ANTIQUUA” [sic] (Férussac 1823: 149) poses a special case. It appeared first in the legend of plate 1 of the “Mélanoïdes fossiles” in the “Histoire naturelle” but – if the format was the same as in the monograph – not in italic font as are the other names. In the monograph, it is the only term on the subspecific level Férussac gave in small caps, a format he otherwise used only for vernacular names of species. Because of this and the fact that he did not add “nobis” it is unlikely that Férussac wanted to introduce a new name; rather, he may have intended to indicate its status as a fossil (Latin *antiqua* = old). If it had been introduced as a distinct taxon, the name would have been invalid as an objective junior synonym of *Melanopsis fusiformis* Sowerby, 1822, which Férussac listed in synonymy.

Férussac (1822) also introduced two additional names (*inflata*, *elongata*) in the captions of plate 1 of the “Mélanoïdes fossiles”, ranked below var. γ [“ANTIQUUA”]. In the monograph (1823) they are clearly marked with “nobis”. If “*antiquua*” [= *fusiformis*] was considered as a distinct taxon, these names would be of infrasubspecific rank, which is not governed by the Code. Both names would have nonetheless become

Table 1. Férussac's original figure legends of the two plates of "Mélanoïpsides fossiles" (1822/1823) in the "Histoire naturelle", compared with the altered legends supplied by Deshayes in 1839 when finalizing the work. Note that Férussac's original figure legends for plate 1 are unavailable to me and thus based on the legends for plate 7 in Férussac (1823), which are supposed to be identical (see text for details).

Plate	Férussac's original legend (1822/1823)	Deshayes' altered legend (1839)
1	Fig. 1, 2. <i>Melanopsis Buccinoidea</i> , var. γ) antiqua: <i>inflata</i> . Des environs d'Epernay. <i>Melanopsis fusiformis</i> , Sowerby, <i>Min. conch.</i> , tab. 232, fig. 1, 5, 7	Fig. 1–7. Melanopsis fusiformis , SOWERBY. Diverses variétés d'Epernay, de Cuiseaux et d'Angleterre
	Fig. 3. La même plus jeune, du lieu dit les Rozières, près d'Épemay	
	Fig. 4. <i>Melanops. Buccinoidea?</i> var. ε) <i>Minuta</i> ; <i>fossilis</i> . De Cuiseaux, près St.-Amour, dans le bassin de la Saône. <i>Nota.</i> Les tours de spire sont un peu trop détachés dans la figure	
	Fig. 5. <i>Melanops. Buccinoidea</i> , var. γ) antiqua: <i>elongata</i> . Des environs d'Epernay. <i>Melanops. fusiformis</i> , Sowerby, loc. cit., fig. 2, 3, 6	
	Fig. 6. La même, de l'île de Wight. Sowerby, <i>id.</i>	
	Fig. 7. La même plus âgée, des environs d'Epernay	
	Fig. 8. <i>Melanops. Buccinoidea</i> , var. α). <i>Fossilis</i> . de Sestos	8–11. — buccinoïdes , FÉRUSAC. Des îles de la Grèce et d'Italie
	Fig. 9. <i>Melanops. Buccinoidea</i> , var. γ) antiqua: <i>inflata</i> . Du dépôt situé entre St.-Germinini et Carsoli; Italie	
	Fig. 10. <i>Melanops. Buccinoidea</i> , var. α). <i>fossilis</i> . De l'île de Rhodes	
	Fig. 11. <i>Melanops. Buccinoidea</i> , var. δ). <i>fossilis</i> . Des dépôts situés entre St.-Germini et Carsoli, et entre Otricoli et le Vigne	
	Fig. 12. <i>Melanops. incerta?</i> var. de Sestos	12. — incerta , FÉR. De Sestos
	Fig. 13. <i>Melanops. nodosa</i> . Du dépôt situé entre Otricoli et le Vigne, route de Rome à Foligno	13. — nodosa , FÉRUSAC. Otricoli
	Fig. 14. <i>Melanops. costata</i> , de Sestos	14–15. — costata , FÉR. De Sestos
	Fig. 15. La même plus petite, de Sestos	
	Fig. 16. <i>Melanops. Dufourii</i> , var. ε). <i>Fossilis, maxima</i> . Des environs de Dax	16. — Dufourii , FÉR. Dax
2	Fig. 1. <i>Melanopsis buccinoidea</i> , var. a) <i>Bulimus antediluvianus</i> , POIRET.	Fig. 1, 2, 4, 5. Melanopsis buccinoidea , FÉRUSAC. Variétés de Carsoli, Italie
	Fig. 2. Id. var. γ) <i>Antiquua</i> . Du dépôt situé entre Saint-Germini et Carsoli	
	Fig. 3. Id. même var. Plus âgée. Du même lieu	3. — subcarinata , DESH. d'Italie
	Fig. 4. Id	
	Fig. 5. <i>Melanops. Dufourii</i> , var. a. De Dax	
	Fig. 6. <i>Melanops. incerta</i> , var. a). Jeune de Sestos	6. — Audebardi , DESH. De Sestos. (Confondue par Férus. avec l' <i>incerta</i>)
	Fig. 7. <i>Melanops. atra</i> , fossile? De l'île de Luçon	7. — atra , FÉRUS. De Luçon
	Fig. 8. <i>Melanops. nodosa</i> , var. a) <i>cylindracea</i> . Dans une roche calcaire d'Athènes	8. — nodosa , FÉRUS. Var. <i>longa</i> d'Athènes
	Fig. 9. <i>Melanops. Bouei</i> . De la Moravie	9–10. — Bouei , FÉRUSAC. De la Moravie
	Fig. 10. Id. var.	
	Fig. 11, 13. <i>Melanops. Martiniana</i> . Grand exemplaires	11–13. — martiniana , FÉRUS. De la Moravie
	Fig. 12. La même coquille; individus moins âgés	

Table 2. Available species-group names (in italics) and unnamed varieties in Férussac (1823). Only the names in bold were first introduced in that work. The assigned status as Recent (R) or fossil (F) is based on Férussac's concept of the taxon in question, whether or not this reflects the latest opinion. Abbreviations: Lam. – Lamarck; O. – Olivier.

Species	Variety	Infrasub-specific name	Remarks	Fossil/ recent
<i>buccinoidea</i> O.				R+F
	α			R+F
	β			R
	γ ("ANTIQUUA")		If considered available, it was a junior objective synonym of <i>M. fusiformis</i> Sowerby, 1822	F
		<i>inflata</i>	Already available from Férussac's "Histoire naturelle"	F
		<i>elongata</i>	Already available from Férussac's "Histoire naturelle"	F
	δ			F
	ϵ			F
<i>dufourii</i>			Already available from Férussac's "Histoire naturelle"	R+F
	α			F
	β			R
	γ			R
	δ			R
	ϵ			F
	ζ			R
	η			R
	θ			R
<i>martiniana</i>			Junior objective synonym of <i>M. fossilis</i> (Gmelin, 1791)	F
<i>incerta</i>				F
<i>costata</i> O.				R+F
	α ("Fasciata")		Nomen nudum	?
<i>costellata</i>			Junior objective synonym of <i>M. cariosa</i> (Linnaeus, 1767)	R
	α			R
<i>nodosa</i>			Already available from Férussac's "Histoire naturelle"	R+F
	α			F
<i>bouei</i>				F
<i>decussata</i>				R
<i>esperi</i>				R
<i>acicularis</i>				R+F
	α			R
	β			R
<i>atra</i> Lam.				R
<i>spinosus</i> Lam.				R

available as species-group names after Art. 45.6.4.1 at least from Pallary (1916: 77), who used them as valid species names. At that time, however, both of them were junior homonyms and thus invalid.

Pallary (1916: 77–78) considered several of the descriptive Latin terms as available names, i.e., “*M. antiqua* Férussac”, “*M. affinis* Férussac”, “*M. magna* Férussac”, “*M. minuta* Férussac”, “*M. parva* Férussac” and “*M. subtuberculata* Férussac”. Since Pallary (1916) clearly associated those names with illustrations in Férussac (1823), they became available thereby, but with the authority “Pallary, 1916”. This actually also includes *Melanopsis antiqua*, which had been an unavailable name until then.

Names introduced by Jean-Pierre Sylvestre de Grateloup in 1828 and 1838

A similar problem as for Férussac’s monograph appears in works by Grateloup (1828, 1838). He labelled varieties with Latin letters, occasionally followed by Latin terms (e.g., “*olivula*”) and Latin descriptions and sometimes marked with “Nob.”. In his 1828-work he attributed the Latin terms *minor* and *major* to varieties of two different species. Apparently, most of these terms were meant as keywords rather than real names. This becomes obvious also from Grateloup (1838), where he introduced the new name *Melanopsis gibbosula* for specimens treated as “var. b. *minor*” in Grateloup (1828).

In order to bring stability to the problem, I propose to use only the Latin terms marked by Grateloup with “Nob.” as available names.

The taxonomic concepts applied by Jules René Bourguignat, Paul Pallary, Jean-Marie Pérès and Ferdinand Starmühlner

The works by Bourguignat and Pallary between 1853 and 1939 extended the list of melanopsid names enormously (Figure 1). Both authors applied very detailed taxonomic concepts and introduced dozens of species and varieties for even minor morphological deviations or differences in shell color. This procedure artificially inflated present-day melanopsid diversity, and few of those names are actually used today (Glaubrecht 2011). Nonetheless, most of the names are nomenclaturally available.

Unfortunately, Bourguignat and Pallary did not foresee that subspecies (as well as forma and variety names published before before 1961, Art. 10.2) would all become included by the Code in the species group, with the Principle of Homonymy applying throughout. Both malacologists introduced many varieties, such as *minor* and *major*, often several times for different species within the very same work. Pallary apparently considered some of the variety names he introduced as self-explanatory (e.g., Latin *minor* means “small”) and left them undescribed (see, e.g., Pallary 1904: 37). In order to avoid inflating the present catalogus with anyway unavailable and taxonomically hardly useful names, I chose not to include these nomina nuda.

Finally, Pérès (1938a, b, 1939, 1946) and Starmühlner (1970, 1973) applied elaborate systems to categorize morphologies and coloration patterns of recent melanopsids. Pérès introduced several “formae” and “modes” that he obviously did not use as




Figure 1. Numbers of melanopsid taxa described between 1758 and 2015 per year and per five years. The most important contributions, i.e., pioneer works and those with numerous new taxa described, are highlighted.

valid names but rather as descriptive terms to fit existing species into his morphological concept (see, e.g., Pérès 1939: 135–137, Pérès 1946: 166). He even assigned “types” to his morpho-units (Pérès 1939: 137). Although hardly any of his new names have any meaning in nomenclature or taxonomy, the available names are included and discussed herein.

Starmühlner (1970, 1973) in a similar way classified New Caledonian specimens of *Melanopsis frustulum* according to form and size of the shell (“forma”), type and height of the spire (“modus”) and coloration (“coloratus”). None of the names used by Starmühlner are available in nomenclature because they were introduced after 1961 and are of infrasubspecific rank (Art. 10.2). They are not included in this catalogus.

Note on the information provided

The catalogue lists all names in alphabetical order in the original spelling and combination, with the necessary amendments required by the Code. The status of taxa that are invalid, unavailable or unresolved is denoted in square brackets after the taxon name; those without status declaration are available and nomenclaturally valid, irrespective of their taxonomic status. The first description, or alternatively the basis of record for unavailable names, is always indicated. Taxa solely found as fossils are marked by a dagger. Taxon authorities attributed to a person other than one of the authors in the original source are only accepted as such if there is clear evidence that the description derived

from that person (Art. 50.1.1). In such cases, the notation is given in accordance with Recommendation 51E. Information on type locality, type horizon (for fossils only) and type specimens are indicated as far as available. The exact spelling or phrase (given in quotes) provided for the type locality in the original source is denoted, along with an English translation if required. If the localities have been indicated indirectly (e.g., “same as for the previous species”), the phrase is given in square brackets. Old locality names have been matched with today’s geographic names as far as possible, mostly using the GeoNames geographical database (<http://www.geonames.org/v3/>). Places that could not be found on the map or where the matched name is uncertain are marked by a postposed question mark in the translation.

Note that Wenz (1929) in his *Fossilium Catalogus* quite often misstated type localities, which he indicated in spaced letters. For taxa where no type has been fixed and several localities have been mentioned in the original publication, he apparently preferred localities mentioned first or those indicated for illustrated specimens. After Art. 76.1 of the Code, in absence of a holotype or lectotype the type locality encompasses the localities of all syntypes. Illustrated specimens do not automatically qualify as holotypes or have any priority in this respect.

Information on type horizons follows the most recent age classifications found in the literature (e.g., Neubauer et al. 2015a, Georgopoulou et al. 2015 for the Neogene and Quaternary freshwater deposits of Europe). As far as possible, the present taxonomic status of a taxon is indicated, i.e., whether it has been considered a junior synonym by later authors. Data on type specimens are provided for all names where information is available, particularly for those published after 1999, when a type fixation became mandatory (Art. 16.4). If not indicated otherwise, the information provided regards the storage of the holotype. For names published by Spiridion Brusina and Petar S. Pavlović the inventory catalogues of the Croatian Natural History Museum in Zagreb (Milan et al. 1974) and the Natural History Museum in Belgrade (Milošević 1962) formed the basis. Note, however, that Milan et al. (1974) often wrongly indicated holotypes from syntype series, apparently considering illustrated specimens preferable to the rest of the material (see also discussion in Neubauer et al. 2016b). Milošević (1962) provided inventory numbers of the specimens illustrated by Pavlović; neither of both applied a type concept.

Nomenclatural notes

The following list of species-group names comprises several recurrent nomenclatural issues. In order to save space, to avoid multiple elaborate repetitions of the same rules, and to prevent that the reader needs to consult the Code constantly, I refer in text to two nomenclatural notes which are defined as follows:

Note 1: Because of the Principle of Coordination (Art. 46), homonymy in the species-group does not depend on a taxon’s original rank in the species-group. This also

encompasses variety and forma names published before 1961 (Art. 10.2). Many authors have been unaware that species names can constitute junior homonyms of subspecies, variety or forma names. Only in the case of simultaneously published names the taxon of higher rank takes precedence (Art. 24.1).

Note 2: Several names of Melanopsidae first occurred in synonymy lists of other names (see, e.g., Brot 1874–1879). According to Art. 11.6 of the Code, “a name which when first published in an available work was treated as a junior synonym of a name then used as valid is not thereby made available”. Some of the names, however, have been made indeed available following the provisions of Art. 11.6.1: “However, if such a name published as a junior synonym had been treated before 1961 as an available name and either adopted as the name of a taxon or treated as a senior homonym, it is made available thereby but dates from its first publication as a synonym [...]”.

† *Melanopsis aaronsohni* Blanckenhorn in Blanckenhorn & Oppenheim, 1927

Original source. Blanckenhorn and Oppenheim 1927: 37, pl. 21, figs 14–15.

Type horizon. Early Pleistocene.

Type locality. “10 Minuten nördlich von der Station Djisr et-Medjâmi” (p. 34) [10 minutes north of the station Djisr et-Medjâmi (= ‘Erq el-Ahmar, also known as Gesher)], Israel.

† *Melanopsis costata* var. *abbreviata* Brusina, 1874

Original source. Brusina 1874: 41, pl. 7, fig. 10.

Type horizon. Cernikian, Pliocene.

Type locality. “Podvinje (Čaplja) [Čaplja trench near Slavonski Brod]; Novska; Farkašić”, Croatia.

Types. Milan et al. (1974: 89) indicated a holotype, but it is uncertain whether the specimen was the only one Brusina had at hand (holotype by monotypy, Art. 73.1.2). The specimen is stored in the Croatian Natural History Museum, Zagreb, coll. no. 3717-1357/1.

† *Melanopsis abbreviata* Pallary, 1916 [invalid]

Original source. Pallary 1916: 81.

Type horizon. Middle Lutetian, Eocene.

Type locality. “Au Nord d’Albas” (Doncieux 1908: 204), France.

Remarks. Replacement name for *M. brevis* Doncieux, 1908, non Sowerby, 1826. Junior homonym of *M. abbreviata* Brusina, 1874 (see *M. atacica* Wenz, 1928).

† *Melanopsis abichi* Calvert & Neumayr, 1880

Original source. Calvert and Neumayr 1880: 376, pl. 2, figs 20–22.

Type horizon. Late Sarmatian, Khersonian, late Miocene.

Type locality. “Renkiöi” [north of İntepe], Turkey.

Melanopsis (Mesopotamia) mesopotamica var. *abrahamiana* Pallary, 1939

Original source. Pallary 1939: 100, pl. 5, figs 1–2.

Type locality. “Ain Arouss” [‘Ayn al ‘Arūs, near Tall Abyaḍ], Syria.

† “*Melanopsis martiniana abscissa* Handmann” mentioned in Fischer (1996b: 20, fig. 2) [unavailable]

Horizon. Pannonian, zone B–D, late Miocene.

Locality. “Leobersdorf”, Austria.

Remarks. Nomen nudum, based on an “in schedis” determination by Handmann (the corresponding label is illustrated in Fischer 1996b: 21, fig. 2).

† *Melanopsis absyrtidum* Stache in Sandberger, 1871

Original source. Sandberger 1870–1875: 134, pl. 19, figs 13–13a.

Type horizon. Cretaceous.

Type locality. “Insel Unie bei Lussin” [Unije island], Croatia.

† *Melanopsis (Canthidomus) acanthica* Neumayr, 1869

Original source. Neumayr 1869: 357, pl. 11, figs 6–7.

Type horizon. Langhian, middle Miocene.

Type locality. “Miocic” [Miočić], Croatia.

Types. Illustrated syntypes are stored at the Geological Survey Austria, Vienna, coll. no. 1869/01/4/1-2.

† *Melanopsis acanthicoides* Hoernes, 1876

Original source. Hoernes 1876: 14, figs 1–5.

Type horizon. Late Sarmatian, Khersonian, late Miocene.

Type locality. “Renkiöi” [north of İntepe], Turkey.

† *Melanoptychia acanthicula* Bourguignat, 1880

Original source. Bourguignat 1880: 31.

Type horizon. Langhian, middle Miocene.

Type locality. “Vallée de la Cettina” [Cetina river valley], Croatia.

Remarks. The name “*acanthinula*” as mentioned in Pallary (1926a: 74) is an incorrect subsequent spelling. Wenz (1929: 2650) considered the taxon as a junior synonym of *Melanopsis acanthica* Neumayr, 1869.

† *Melanopsis (Martinia) martiniana* var. *accedens* Handmann, 1887

Original source. Handmann 1887: 24, pl. 3, figs 15–16.

Type horizon. Pannonian, zone B–D, late Miocene.

Type locality. “Leobersdorf”, Austria.

Remarks. Wenz (1929: 2719) considered this taxon as a junior synonym of *Melanopsis fossilis* (Gmelin, 1791).

Melanopsis acicularis Férussac, 1823

Original source. Férussac 1823: 160.

Type locality. “La rivière de Laybach. [...] Les eaux thermales de Weslau, près de Vienne. [...] Le Danube, à Wissegrad et à Bude. [...] de l’île de Wight” [Ljubljanica river; in the thermal water of Vöslau near Vienna; in the Danube, at Visegrád and Budapest; from the Isle of Wight], Slovenia.

Remarks. Partly a junior objective synonym of *Melanopsis subulata* Sowerby, 1822 (regarding the specimen of the Isle of Wight) and *Melanopsis daudebartii* [Prevost], 1821 (regarding the Vöslau material), which Férussac considered varieties of *M. acicularis* and listed in synonymy. Currently considered as a junior synonym or subspecies of *Microcolpia daudebartii* [Prevost], 1821, respectively (Welter-Schultes 2012: 36; Neubauer et al. 2014d).

Melanopsis aciculella Potiez & Michaud, 1838

Original source. Potiez and Michaud 1838: 346, pl. 31, figs 3–4.

Type locality. “Les rivières de la Carniole” [rivers of Carniola, a historical region that comprised parts of present-day Slovenia].

Remarks. Starobogatov et al. (1992: 65) considered the species as a junior synonym of *Microcolpia cornea* (Pfeiffer, 1828).

† *Turripontica aciculina* Anistratenko, 1993

Original source. Anistratenko 1993: 75, textfig. 3.

Type horizon. Duab Beds, middle to late Kimmerian, Pliocene.

Type locality. “Окр. с. Мокви, Очамчирский р-н” [near the village Mok’vi, Ochamchirskiy rayon], Georgia.

Types. Schmalhausen Institute of Zoology of National Academy of Sciences of Ukraine, Kiev; no number indicated.

Remarks. Type species of the genus *Turripontica* Anistratenko, 1993.

† *Melanopsis acrolepta* Fontannes, 1884

Original source. Fontannes 1884: 29, pl. 4, figs 1–2.

Type horizon. Early Rupelian, Oligocene.

Type locality. “Barjac, Roméjac, Avéjan, Saint-Jean-de-Maruéjols, Célas, Issirac, Galès, près de Montclus (Gard)”, France.

Fagotia acroxia Bourguignat, 1884

Original source. Bourguignat 1884: 48.

Type locality. “Dans la rivière au-dessous de Krapina-Toeplitz, en Croatie” [in the river below Krapinske toplice], Croatia.

Remarks. Note that Bourguignat denoted the authority as “Bourguignat, 1879”.

† *Melanopsis acuminata* Gümbel, 1861

Original source. Gümbel 1861: 753.

Type horizon. Chattian, Oligocene.

Type locality. “In der oberen Leizach, an der Schlierach, im Sulzgraben, bei Pensberg, Rimselrain, im Höllbache, am hohen Peissenberge” [all localities are near Miesbach, southern Bavaria], Germany.

Remarks. Gümbel attributed the authority to Sandberger, but there is no evidence that the description really derived from that author.

† *Melanopsis bouei* var. *acuminata* Handmann, 1882 [invalid]

Original source. Handmann 1882: 557.

Type horizon. Pannonian, zone D, late Miocene.

Type locality. “Kottingbrunn [...] Ziegelei a”, Austria.

Remarks. Junior homonym of *Melanopsis acuminata* Gümbel, 1861. Not included in the Fossilium Catalogus of Wenz (1929).

***Melanopsis costata* var. *acuminata* Bourguignat, 1884 [invalid]**

Original source. Bourguignat 1884: 140.

Type locality. “Le Jourdain, à 4 kilomètres au-dessus de la Mer Morte” [Jordan river, 4 km north of the Dead Sea], Israel/Jordan.

Remarks. Junior homonym of *Melanopsis acuminata* Gümbel, 1861.

† *Melanopsis acuminata* Pallary, 1901 [invalid]

Original source. Pallary 1901a: 178, pl. 2, fig. 23.

Type horizon. Plio-Pleistocene.

Type locality. “Du puits Karoubi” [from the well Karoubi], Algeria.

Remarks. Junior homonym of *Melanopsis acuminata* Gümbel, 1861. Wenz (1919a: 73) introduced *Melanopsis pallaryi* as replacement name (see also *M. raphidia* Pallary, 1920).

† *Melanopsis acuminata* Seninski, 1905 [invalid]

Original source. Seninski 1905: 62, pl. 2, figs 3–4.

Type horizon. Duab Beds, middle to late Kimmerian, Pliocene.

Type locality. “Моквинские пласты, р. Дуабъ” [Mokvi layers at Duab river], Georgia.

Remarks. Junior homonym of *Melanopsis acuminata* Gümbel, 1861. Wenz (1928a) introduced *Melanopsis seninskii* as replacement name.

† *Melanopsis acuta* Handmann, 1882

Original source. Handmann 1882: 556.

Type horizon. Pannonian, zone D, late Miocene.

Type locality. “Kottingbrunn [...] Ziegelei a”, Austria.

***Melanopsis acutespira* Bourguignat, 1884**

Original source. Bourguignat 1884: 114.

Type locality. “Dans l’aqueduc de la Palafanga, près Almazora (Espagne), aux environs de Mascara (Algérie)” [in the aqueduct Palafanga near Almazora (Spain), near Mascara (Algeria)].

Remarks. Bourguignat introduced this species for formerly misidentified *M. dufouriei* [sic] *graellsii* sensu Rossmässler, 1854 and *M. maroccana* sensu Paladilhe, 1875 as well as a part of the material of the variety *M. maroccana subgraellsiana* Bourguignat, 1864. He apparently overlooked that Paladilhe (1875) had not referred to *M. maroccana* but had introduced the new variety *M. maroccana zonatosubcostata*. Since there was never a holotype or lectotype defined, the name *acutespira* comprises the syntypes of all three works; it remains nomenclaturally valid as name for the two specimens illustrated by Rossmässler (1854) and Bourguignat (1864).

***Melanopsis acutissima* Gassies, 1871**

Original source. Gassies 1871: 197, pl. 6, fig. 13.

Type locality. “Bélep (île Art)” [Bélep, Art Island], New Caledonia.

***Melanopsis acutula* Pallary, 1920**

Original source. Pallary 1920a: 31.

Type locality. “Dans un bassin entre le Mellah et le pont; vers Dar Mahrès; [...] Bahlil (28 kil. au sud de Fès)” [in a basin between the Mellah and the bridge; toward Dar Mahres; Bhalil (28 km south of Fes)], Morocco.

***Melanopsis adrarensis* Pallary, 1912**

Original source. Pallary 1912: 16, figs 8–9.

Type locality. “Dans les seguias des oasis du Touat, spécialement dans celles de l’Adrar” [in the irrigation channels of the oasis of Touat, especially those of Adrar], Algeria.

Remarks. Given as “*adparensis*” on p. 16, but “*adrarensis*” in plate captions. Since Pallary clearly denoted it from the locality Adrar, the name must be “*adrarensis*” (Art. 32.5.1).

† *Melanopsis aegaea* Tournouër, 1875

Original source. Tournouër 1875: 76.

Type horizon. Phoka to Elia Formation, Plio-Pleistocene.

Type locality. “[Prope vicum Antimaki et prope civitatem Cos], in loco Hagios-Foukas” [near near the village Antimácheia and near the city of Kos, in the locality Ágios Fokás], Greece.

Remarks. The name “*aegaea*” as mentioned in Tournouër (1876: 453), Wenz (1929: 2653) and Willmann (1981: 174) is an incorrect subsequent spelling.

† *Melanopsis aegyptiaca* Blanckenhorn, 1901

Original source. Blanckenhorn 1901: 414, pl. 15, figs 15–16.

Type horizon. Pleistocene.

Type locality. “In der Fossilienbank an der Tewfik-Moschee bei Kairo. [...] Wadi Urag, Sanur, Moschasch, Raijade” [in the fossil deposits at the Tewfik Mosque near Cairo. In the wadis Urag (?), Sannūr, Moschasch (?), Raijade (?)], Egypt.

† *Melanopsis aetolica* Neumayr, 1876

Original source. Neumayr 1876: 368.

Type horizon. Gelasian, early Pleistocene.

Type locality. “Stamna, nordwestlich von Missolonghi” [Stamná, NW of Mesolóngi], Greece.

Types. Papp (1955: pl. 20, fig. 9) designated a specimen from the type material of Neumayr (1880a) as lectotype, but did not indicate whether it had been illustrated by Neumayr or not. A comparison of the images in both works was inconclusive. However, since it is likely that Neumayr (1880a) used the material reported in 1876 for the description, this designation is considered to be based on the original material and is thus valid. The specimen is stored in the Institute of Paleontology, University of Vienna; no number indicated.

Remarks. The name became available from Neumayr (1876), where he briefly described the species. He nevertheless described the species as “new” in Neumayr (1880a: 126), providing a detailed description and illustrations (pl. 6, figs 13–17; note that the plate is inserted at the end of the previous article of the same volume).

† *Melanopsis affinis* Handmann, 1882

Original source. Handmann 1882: 558.

Type horizon. Pannonian, zone D, late Miocene.

Type locality. “Kottingbrunn [...] Ziegelei a”, Austria.

Remarks. Pallary (1916) considered the species a junior homonym of “*M. affinis* Férussac” and introduced *M. subaffinis* as replacement name. “*M. affinis* Férussac” is, however, not an available name (see below) and *M. subaffinis* is thus a junior objective synonym of *M. affinis* Handmann, 1882. Wenz (1929: 2674, 2681) synonymized both Handmann’s and Pallary’s species with *M. bouei* Férussac, 1823.

† *Melanopsis affinis* Pallary, 1916 [invalid]

Original source. Pallary 1916: 78.

Type horizon. Late Villafranchian, early Pleistocene.

Type locality. “D’Italie” (Férussac 1823 had given the locality more precisely as “entre Otricoli et le Vigne, route de Rome à Foligno” []), Italy.

Remarks. The name often appears as “*M. affinis* Férussac” in the literature (e.g., Wenz 1929: 2653; Esu and Girotti 1975: 251). It was first mentioned in Férussac (1814) as nomen nudum for a specimen from the Euphrates River. Later, Férussac (1823) listed the name in synonymy of *M. nodosa* Férussac, 1822. Unlike Pallary (1916) claimed and followed by Wenz (1929), *affinis* is not to be preferred over *nodosa*, because at that time *affinis* was not an available name – it has never been described, indicated or illustrated (as, for example, wrongly presumed by Esu and Girotti 1975) and so does not fulfill the requirements of availability (Art. 11 and 12; see also Art. 11.6). Pallary (1916) was the first to affiliate the name with an illustration, namely the one of *M. nodosa* provided by Férussac, 1823 (pl. 7, fig. 13), and made the name thereby available. The illustrated specimen is, however, the holotype (by monotypy) of *M. nodosa* Férussac, 1822 (Férussac used the same plates in both of his works and thus Pallary actually referred to the specimen illustrated as *M. nodosa* in the “Histoire naturelle”; see also introduction and Table 1 for details of Férussac’s publications). Consequently, *Melanopsis affinis* Pallary, 1916 is a junior objective synonym of *M. nodosa*. Moreover, the name is a junior homonym of *M. affinis* Handmann, 1882.

***Melania afra* Rossmässler, 1839**

Original source. Rossmässler 1838–1844: 38, pl. 50, fig. 665.

Type locality. “Aus der Ringelsza bei Laibach” [Ringelsza (?) brook near Ljubljana], Slovenia.

Remarks. Introduced in synonymy of *M. holandri laevigata*. It was made available at least by Brusina (1867) who treated it as a valid name (see Note 2).

† *Melanopsis matheroni* var. *agatensis* Pantanelli, 1886

Original source. Pantanelli 1886a: 68 or 1886b: 78, pl. 3, figs 1–4 (precedence not established).

Type horizon. Tortonian–early Messinian, late Miocene.

Type locality. “S. Valentino e S. Agata” [San Valentino and Sant’Agata Fossili], Italy.

Remarks. Harzhauser et al. (2015: 9) considered the taxon as a junior synonym of *Melanopsis narzolina* d’Archiac in Viquesnel, 1846.

***Melania holandrii* var. *agnata* Pfeiffer, 1828**

Original source. Pfeiffer 1828: 47, pl. 8, fig. 8.

Type locality. “In der Muhr” [in the river Mur], Austria or Slovenia.

Remarks. Note that Bourguignat denoted the authority incorrectly as “Bourguignat, 1877”. The name “*aequata*” as mentioned in Clessin (1890: 676) is an incorrect subsequent spelling.

***Melanella agnatella* Bourguignat, 1884**

Original source. Bourguignat 1884: 14.

Type locality. “Rivière près Zenica, en Bosnie” [river near Zenica], Bosnia and Herzegovina.

Remarks. Appeared first as a nomen nudum in Servain (1884: 379).

“*Melanopsis agoroea*” mentioned in Bourguignat (1884: 83) [unavailable]

Locality. Not indicated.

Remarks. Nomen nudum: Bourguignat (1884) listed the name in synonymy of *M. laevigata*, perhaps referring to an unused manuscript name. Probably the name was intended as “*agoraea*”, but the ligature was mixed up during typesetting.

***Melanopsis ahui* Ahuir Galindo, 2014**

Original source. Ahuir Galindo 2014b: 7, unnumbered figure.

Type locality. “Around Guefait, at the Northeastern of Morocco”, Morocco.

Types. Museo Malacologico di Cupra Marittima, Italy; no number indicated.

† *Melanopsis ajkaensis* Tausch, 1886

Original source. Tausch 1886: 9, pl. 1, figs 35a–c.

Type horizon. Late Santonian–early Campanian, late Cretaceous.

Type locality. “Csingerthal bei Ajka” [Csinger valley near Ajka], Hungary.

† *Melanopsis albasensis* Wenz, 1919 [invalid]

Original source. Wenz 1919a: 73.

Type horizon. Middle Lutetian, Eocene.

Type locality. “Au Nord d’Albas” (Doncieux 1908: 204), France.

Remarks. Introduced as replacement name for the junior homonym *Melanopsis nodosa* Doncieux, 1908, non Férussac, 1822, for which Pallary (1916) had already

introduced *M. doncieuxi* as replacement name. Thus, *M. albasensis* is a junior objective synonym of *M. doncieuxi* Pallary, 1916.

***Melanopsis alepensis* Pallary, 1939 [invalid]**

Original source. Pallary 1939: 86.

Remarks. Unjustified emendation and therefore junior objective synonym of *M. alepi* Bourguignat, 1884.

***Melanopsis alepi* Bourguignat, 1884**

Original source. Bourguignat 1884: 119.

Type locality. “Environs d’Alep” [surroundings of Aleppo], Syria.

Remarks. Heller et al. (2005: 243) tentatively considered the species as a junior synonym of *Melanopsis costata* (Olivier, 1804).

***Melanopsis algerica* Pallary, 1904**

Original source. Pallary 1904: 35.

Type locality. “Oran; [...] Mostaghanem” (Bourguignat 1864: caption of pl. 15; the exact localities where the illustrated specimens had been found are unknown), Algeria.

Remarks. Introduced by indication of illustrations in Bourguignat (1864: pl. 15, figs 12–14, 17–18).

***Melanopsis algericensis* Pallary, 1922 [invalid]**

Original source. Pallary 1922: 207.

Remarks. Unjustified emendation and therefore junior objective synonym of *M. algerica* Pallary, 1904.

† *Melania* (*Campylostylus*) *allobroga* Oppenheim, 1892

Original source. Oppenheim 1892: 766, pl. 30, figs 5–7.

Type horizon. Calcaire de Rognac, Maastrichtian, Cretaceous.

Type locality. “Les Pennes, Valon du Duc bei Rognac” [Les Pennes, Valon du Duc near Rognac], France.

Remarks. Originally the gender was incorrectly given as neutrum (“*allobrogum*”), but *Melania* is feminine.

† *Melanopsis (Lyrcea) almerai* Cossmann, 1909

Original source. Cossmann 1909: 175.

Type horizon. Eocene.

Type locality. “Pyrénées catalanes”, Spain.

Remarks. Cossmann (1909) gave a very brief description (“little keeled”) and made the name available.

† *Melanopsis ancillaroides* var. *alpina* Traub, 1938

Original source. Traub 1938: 76, pl. 6, figs 8a–b.

Type horizon. Paleocene.

Type locality. “Aus Kch Fi1 [etwa 20 m unterhalb der großen betonierten Bachverbauung südöstlich von Kleinoiching am linken Ufer des Kroisbaches]” [from block Kch Fi1, ca. 20 m below the large concrete structure southeast of Kleinoiching, at the left bank of the Kroisbach brook], Austria.

† *Melanopsis alutensis* Stefanescu, 1896

Original source. Stefanescu 1896: 128, pl. 11, figs 12–14.

Type horizon. Early Pliocene.

Type locality. “À Milcov, près de Slatina, dans la vallée de l’Oltu” [at Milcov, near Slatina, in the valley of the river Olt], Romania.

Melanopsis amabilis Pallary, 1928

Original source. Pallary 1928a: 261, pl. 5, figs 19–20.

Type locality. “Mechera Kredar, sur la route de Media à Larache; Bou Hellou (secteur ouest de Taza)” [Mechera Kredar (?), at the road from Mehdy to Larache; Bou Hellou, western part of prov. Taza], Morocco.

† *Melanopsis hybostoma* var. *amaradica* Fontannes, 1887

Original source. Fontannes 1887: 337, pl. 26, figs 20–22.

Type horizon. Early Cernikian, early Pliocene.

Type locality. “Caprenu, val. Amaradii (Jud. Gorjiu)” [Căpreni], Romania.

† “*Melanopsis ambigua* Gaudry, 1862” mentioned in Neumayr (1869: 372) [unavailable]

Horizon. Pliocene?

Locality. Not indicated.

Remarks. Status unclear: the name was mentioned in Neumayr (1869: 372) and Pallary (1926a: 75), but the original description could not be found. Perhaps it is a lapsus calami of *Melanopsis anceps* Gaudry & P. Fischer in Gaudry, 1867.

***Melanella ambly* Bourguignat, 1884**

Original source. Bourguignat 1884: 24.

Type locality. “La Save à Sissek, en Slavonie” [in the Sava river near Sisak in Slavonia], Croatia.

Remarks. Note that Bourguignat denoted the authority as “Bourguignat, 1879”.

† *Melanopsis americana* White, 1883

Original source. White 1883: 96, pl. 4, figs 9–10.

Type horizon. Laramie Group, Cretaceous.

Type locality. “Valley of South Platte River, Northeastern Colorado”, United States.

Remarks. Probably not a Melanopsidae.

***Melanopsis ammanensis* Pallary, 1939 [invalid]**

Original source. Pallary 1939: 88, pl. 4, fig. 3.

Remarks. Unjustified emendation and therefore junior objective synonym of *M. ammonis* Tristram, 1865.

***Melanopsis ammonis* Tristram, 1865**

Original source. Tristram 1865: 542.

Type locality. “Streams at Heshbon and Ammon, east of Jordan”, Jordan.

“*Melanopsis amori*” mentioned in Azpeitia Moros (1929: 40, 250) [unavailable]

Locality. “Pedroche (arroyo de), en el partido judicial de Pozoblanco (Córdoba)” [Pedroche stream, district Pozoblanco, Córdoba], Spain.

Remarks. Based on a manuscript name from Hidalgo and introduced in synonymy of *M. etrusca* (see Art. 11.6).

† *Melanopsis amphora* Oppenheim, 1890

Original source. Oppenheim 1890a: 136, pl. 5, figs 2–2b.

Type horizon. Ronca Beds, Bartonian, Eocene.

Type locality. “Lovara di Tressino, Monte Pulli, Mussolon” [Lovara, Monte Pulli (near Valdagno), Muzzolon], Italy.

Remarks. Wenz (1929: 2846) considered this taxon as a junior synonym of *Melanopsis vicentina* Oppenheim, 1890.

† *Melanopsis ampla* Pallary, 1920

Original source. Pallary 1920b: 109.

Type horizon. Middle Pannonian, late Miocene.

Type locality. “Sulzlacke bei Margarethen nächst Oedenburg [...]. Tinnye bei Ofen” (Fuchs 1873: 20) [Sulzlacke near St. Margarethen (Burgenland, Austria); Tinnye (Hungary)].

Remarks. Replacement name for *M. avellana* Fuchs, 1873, non Sandberger, 1870. Wenz (1929: 2760) considered the taxon as a junior synonym of *M. inermis* Handmann, 1882.

***Melanopsis marocana* [sic] var. *ampla* Pallary, 1920**

Original source. Pallary 1920c: 145.

Type locality. “Tazouta”, Morocco.

Remarks. This name, published in November 1920, is a junior homonym of *M. ampla* Pallary, 1920 [July].

† *Melanopsis textilis* var. *ampullacea* Handmann, 1887

Original source. Handmann 1887: 16.

Type horizon. Pannonian, zone B–D, late Miocene.

Type locality. “Leobersdorf”, Austria.

***Fagotia anatolica* Bourguignat, 1884**

Original source. Bourguignat 1884: 43.

Type locality. “Rivières près Ismidt (Anatolie)” [rivers near İzmit], Turkey.

Remarks. Note that Bourguignat denoted the authority as “Bourguignat, 1880”. Starobogatov et al. (1992: 63) considered the species as a junior synonym of *Fagotia* [= *Esperiana*] *gallandi* Bourguignat, 1884.

† *Melanopsis anceps* Gaudry & P. Fischer in Gaudry, 1867

Original source. Gaudry 1862–1867: 446, pl. 62, figs 1–6.

Type horizon. Pliocene.

Type locality. “Mégare” (p. 444), Greece.

Fagotia anceyana Bourguignat, 1884

Original source. Bourguignat 1884: 36.

Type locality. “Dans la Save au-dessous d’Agram, dans la rivière de Krapina (Croatie); enfin, dans le lac Sabandja près d’Ismidt (Anatolie)” [in the Sava river below Zagreb, in the river Krapina (Croatia); finally, in Lake Sapanca near İzmit (Turkey)].

Remarks. Starobogatov et al. (1992: 60) considered the species as a junior synonym of *Fagotia* [= *Esperiana*] *esper* (Férussac, 1823).

† *Melanopsis ancillaroides* Deshayes, 1825

Original source. Deshayes 1824–1837: 121, pl. 15, figs 1–2.

Type horizon. Cuisian, late Ypresian, Eocene.

Type locality. “Les environs de Meaux” [surroundings of Meaux], France.

Melanopsis andersoni Pallary, 1923

Original source. Pallary 1923: 41, [unnumbered plate], figs 1–2.

Type locality. “Dans l’Ain-Touagha, à Fatnassa dans le Nefzaoua” [Ain Touagha (?) at Fatnassa in Nefzaoua], Tunisia.

† *Melanopsis andrussowi* Brusina, 1885

Original source. Brusina 1885: 160.

Type horizon. *Spaniodon* Beds, Karaganian, middle Miocene.

Type locality. “Лонушны” (Sinzov 1884: 4) [Lăpușna], Moldova.

Remarks. Introduced for *M. lanzaeana* sensu Sinzov, 1884, non Brusina, 1874.

† *Melanopsis angulata* Neumayr, 1880

Original source. Neumayr 1880c: 479, pl. 7, fig. 8.

Type horizon. Langhian, middle Miocene.

Type locality. “Žepj” [Džepi], Bosnia and Herzegovina.

Types. The type material, with all specimens studied by Neumayr (1880), is lost. Neubauer et al. (2016c: 275) defined a neotype based on material from the type locality. The specimen is stored in the Geological-Paleontological Department, Natural History Museum Vienna, coll. no. 2014/0364/0012.

† *Melanopsis angusta* Pallary, 1916

Original source. Pallary 1916: 82.

Type horizon. Pannonian, zone D, late Miocene.

Type locality. “Kottingbrunn [...] Ziegelei a” (Handmann 1882: 561), Austria.

Remarks. Replacement name for *M. fusiformis* Handmann, 1882, non Sowerby, 1822. Wenz (1929: 2742) considered the taxon as a junior synonym of *Melanopsis haueri* Handmann, 1882.

† *Melanopsis (Canthidomus) hybostoma anili* Taner, 1997

Original source. Taner 1997: 39.

Type horizon. Early Romanian, Pliocene.

Type locality. “Südöstlich vom Hügel Başaltı, 2,2 km W Musaköy, 15 km NE Çanakkale, W-Anatolien” [southeast of hill Başaltı, 2.2 km W of Musaköy, 15 km NE Çanakkale, W Anatolia], Turkey.

Types. Geological-Paleontological Department, Natural History Museum Vienna, Austria, coll. no. 1996/0053/0001.

† *Melanopsis anistratenkoi* Neubauer, Harzhauser, Georgopoulou, Mandic & Kroh, 2014

Original source. Neubauer et al. 2014a: 456.

Type horizon. Duab Beds, middle to late Kimmerian, Pliocene.

Type locality. “Окр. с. Мокви, Очамчирский р-н” (Anistratenko 1993: 69) [near the village Mok’vi, Ochamchirskiy rayon], Georgia.

Types. Schmalhausen Institute of Zoology of National Academy of Sciences of Ukraine, Kiev, coll. no. 22/VI 1989.

Remarks. Replacement name for the junior secondary homonym *Melanopsis cylindrica* Anistratenko, 1993, non *Lyrcea cylindrica* Stoliczka, 1862.

† *Melanopsis anita* Aldrich, 1886

Original source. Aldrich 1886: 35, pl. 5, fig. 12.

Type horizon. Early Eocene.

Type locality. “Gregg’s Landing, Alabama”; Harris (1899, p. 77, pl. 10, fig. 11) designated a lectotype, United States.

Types. Harris (1899: 77) designated a lectotype, which is stored in the collection of the National Museum of Natural History, Smithsonian Institution, Washington, DC, coll. no. 638955 (Palmer and Brann 1966: 755).

Remarks. The taxon does not belong to the Melanopsidae after Allmon (1990: 63).

***Melanopsis nodosa* var. *anodifera* Cerulli-Irelli, 1914**

Original source. Cerulli-Irelli 1914: 185, pl. 15 (47), fig. 7.

Type locality. “M. Mario”, Italy.

Remarks. Girotti (1972: 232) considered the taxon as a junior synonym of “*Melanopsis affinis* Férussac”, which is not an available name.

† *Bulimus antediluvianus* Poiret, 1801

Original source. Poiret 1801: 36, 37.

Type horizon. Sparnacian, early Ypresian, Eocene.

Type locality. “Chemin de Soissons à Château-Thierry” [way of Soissons at Château-Thierry], France.

Remarks. The name “*antediluviana*” as mentioned in Wenz (1929: 2658) is an incorrect subsequent spelling.

† *Melanopsis antiqua* Pallary, 1916

Original source. Pallary 1916: 78.

Type horizon. Late Villafranchian, Pleistocene.

Type locality. “D’Italie” (Férussac 1823 had given the locality more precisely as “entre St.-Germini et Carsoli” [between San Gemini and Carsoli, Italy]).

Remarks. The name “*Melanopsis antiqua* Férussac, 1823 has been used as valid name by several authors, but obviously was not intended as species-group name by Férussac (1823). The name became available from Pallary (1916) who associated the name with an illustration in Férussac (1823: pl. 8, fig. 2). See introduction for detailed discussion about the names used by Férussac (1823).

***Melanopsis aperta* Gassies, 1861**

Original source. Gassies 1861: 291, pl. 7, fig. 11.

Type locality. “Jengen, dans les ruisseaux” [in streams at Hienghène], New Caledonia.

† *Melanopsis bleicheri* var. *apicula* Pallary, 1901

Original source. Pallary 1901a: 178, pl. 2, fig. 26.

Type horizon. Plio-Pleistocene.

Type locality. “Puits Karoubi” [from the well Karoubi], Algeria.

Melanopsis graellsii var. *apiculata* Pallary, 1924

Original source. Pallary 1924: 250, pl. 25, fig. 15.

Type locality. “Almenara (entre Castellón et Valencia)” [Almenara, between Castellón and Valencia], Spain.

† *Melanopsis turrisformis* *applanata* Schütt in Schütt & Ortal, 1993

Original source. Schütt and Ortal 1993: 92, pl. 2, figs 21–23.

Type horizon. Pleistocene, Mindel glacial epoch.

Type locality. “Galilee, ‘Ubeidiya [El ‘Ubeidiya], 3 km SE of the Sea of Galilee”, Israel.

Types. Paleontology Collection of the Hebrew University of Jerusalem; no number indicated.

Melanopsis cossoni var. *aprica* Bourguignat, 1884

Original source. Bourguignat 1884: 112.

Type locality. “Dans les Seguia des jardins de Miliana, premier ksar au nord de l’oasis d’Insalah, dans le Sahara” [in the irrigation channel (?) in the gardens of Miliana, first ksar (= fortified village, castle) north of the oasis of In Salah in the Sahara], Algeria.

† *Melanopsis aquensis* Grateloup, 1838

Original source. Grateloup 1838: 139, pl. 4, figs 48–49.

Type horizon. Burdigalian, early Miocene.

Type locality. “Dax; Mandillot, à Saint-Paul”, France.

Remarks. Wenz (1929) synonymized the name with “*M. major* Férussac, 1823”, which is not an available name.

† *Melanopsis aquitanica* Pallary, 1916

Original source. Pallary 1916: 79.

Type horizon. Burdigalian, early Miocene.

Type locality. “Dax, St-Paul, Mandillot” (Grateloup 1840: captions of the plate “Mollusques terrestres et fluviatiles fossiles de Dax”), France.

Remarks. Introduced for *M. dufourii* sensu Grateloup, 1840, non Férussac, 1822. Wenz (1929: 2662) considered the taxon as a junior synonym of “*Melanopsis major* Férussac, 1823”, which is not an available name.

***Melanopsis scalaris* var. *arbalensis* Pérès, 1939**

Original source. Pérès 1939: 144, pl. 4, figs 15, 19–20.

Type locality. “Stations 3, 4, 1 3, 29, 35, 143. Oued Tizguit; [...] Station 109. Ain Aghbal près d’Azrou” [Stations 3, 4, 1 3, 29, 35, 143 at Oued Tizguit; station 109 at Ain Aghbal near Azrou], Morocco.

† ***Melanopsis arcuata* Brusina, 1878**

Original source. Brusina 1878: 348.

Type horizon. Cernikian, Pliocene.

Type locality. “Malino”, Croatia.

Types. The syntypes are stored in the Croatian Natural History Museum, Zagreb; no number indicated (Milan et al. 1974: 85).

Remarks. Pallary (1925) introduced *M. simulata* as replacement name for the “junior homonym” *M. arcuata* Brusina, 1878, non Matheron, 1842, which is not available name but an incorrect subsequent spelling of *M. armata* Matheron, 1842.

† ***Melanopsis armata* Matheron, 1842**

Original source. Matheron 1842: 294, pl. 37, figs 12–14.

Type horizon. Calcaire de Rognac, Maastrichtian, Cretaceous.

Type locality. “Rognac et St. Victoret”, France.

Remarks. The name “*arcuata*” as mentioned in Pallary (1925: 257) is an incorrect subsequent spelling. Based on this error, Pallary (1925) considered *M. arcuata* Brusina, 1897 a junior homonym and introduced the replacement name *M. simulata* (junior objective synonym). Sandberger (1871: 101) attributed the species to the genus *Paludomus* Swainson, 1840 (Paludomidae). After Calzada and Urquiola (1994: 152) this species belongs in the genus *Cosinia* Stache, 1880 (Paludomidae).

***Melanopsis (Mesopotamia) aroussiana* Pallary, 1939**

Original source. Pallary 1939: 100, pl. 5, figs 22–25, 56.

Type locality. “Ain Arouss” [‘Ayn al ‘Arūs, near Tall Abyad], Syria.

† *Melanopsis arsinovi* Brusina, 1902

Original source. Brusina 1902: pl. 6, figs 2–4.

Type horizon. Early Langhian, middle Miocene.

Type locality. “Zvezdanski ključ” [village Zvezdan], Serbia.

Types. The illustrated syntypes are stored in the Croatian Natural History Museum, Zagreb, coll. no. 2503-149/1-3 (Milan et al. 1974: 86).

***Fagotia ascanica* Bourguignat, 1884**

Original source. Bourguignat 1884: 39.

Type locality. “Lac Sabandja” [Lake Sapanca], Turkey.

Remarks. Note that Bourguignat denoted the authority as “Bourguignat, 1880”.

***Melanopsis ascanica* Bourguignat, 1884**

Original source. Bourguignat 1884: 96.

Type locality. “Lac Sabandja, près d’İsmidt (Anatolie)” [Lake Sapanca near İzmit], Turkey.

Remarks. Appeared first as a nomen nudum in Locard (1883a: 201). Note that Bourguignat denoted the authority as “Bourguignat, 1880”.

***Melanopsis ashkhabadensis* Izzatullaev & Starobogatov, 1984**

Original source. Izzatullaev and Starobogatov 1984: 1480, fig. 1 (14), fig. 3 (8).

Type locality. “Ашхабад, горные быстро текущие ручьи - Ферюза, Гули и др.” [Ashgabat, fast mountain streams - Firyuza, Guli and others], Turkmenistan.

Types. Zoological Institute of Russian Academy of Sciences, St.-Petersburg; no number indicated.

† *Melanopsis arbalensis* f. *assakaensis* Jodot, 1965 [unavailable]

Original source. Jodot 1965: 104, fig. 2.

Type horizon. Quaternary.

Type locality. “Oued Assaka” [Oued Asaca], Morocco.

Remarks. Unavailable for two reasons: First, the original work lacks a verbal description of the taxon which is required for names published after 1930 (Art. 13.1.1). Second, the taxon was introduced after 1961 as “forma” which is deemed to be infrasubspecific after Art. 15.2.

† *Melanopsis astathmeta* Brusina, 1897

Original source. Brusina 1897: 8, pl. 6, figs 13–16.

Type horizon. Cernikian, Pliocene.

Type locality. “Malino”, Croatia.

Types. Milan et al. (1974: 86) indicated a holotype, but it is uncertain whether the specimen was the only one Brusina had at hand (holotype by monotypy, Art. 73.1.2). The specimen is stored in the Croatian Natural History Museum, Zagreb, coll. no. 3000-646.

† *Melanopsis astrapaea* Brusina, 1876

Original source. Brusina 1876: 110.

Type horizon. Early Langhian, middle Miocene.

Type locality. Originally given as “Sinj” and later specified as “Župića potok” in Brusina (1884b), Croatia.

Types. Milan et al. (1974: 86) stated that Brusina (1897) had indicated one of the specimens illustrated by him (pl. 4, fig. 1) as type. However, it is uncertain whether the specimen actually derives from the original type series. The specimen is stored in the Croatian Natural History Museum, Zagreb, coll. no. 2968-614/1-2.

Melanopsis praemorsa var. *astropaliae* Gambetta, 1929

Original source. Gambetta 1929: 104.

Type locality. “Stampalia” [Astypalaia Island], Greece.

† *Melanopsis atacica* Wenz, 1928

Original source. Wenz 1928a: 119.

Type horizon. Lutetian, Eocene.

Type locality. “Au Nord d’Albas” (Doncieux 1908: 204), France.

Remarks. Replacement name for *M. doncieuxi* Wenz, 1919, non Pallary, 1916, which in turn was introduced as replacement name for *M. brevis* Doncieux, 1908, non Sowerby, 1826 (see also *M. abbreviata* Pallary, 1916).

† *Melanopsis (Lyrcaea) slavonica* var. *atanasiui* Şoverth, 1953

Original source. Şoverth 1953: 206, pl. 3, figs d-d”.

Type horizon. Romanian, Pliocene–early Pleistocene.

Type locality. “Hurezanii-de-Sus - Hurezanii-de-jos”, Romania.

***Melanopsis aterrima* Bourguignat, 1884**

Original source. Bourguignat 1884: 127.

Type locality. “Dans deux ou trois sources de la plaine de Jéricho (Syrie)” [in two or three sources of the plain of Jericho], Palestine.

Remarks. Heller et al. (2005: 248) considered the species as a junior synonym of *Melanopsis saulcyi* Bourguignat, 1853.

***Melanopsis chehirensis* var. *atra* Pallary, 1939**

Original source. Pallary 1939: 94.

Type locality. Not explicitly stated but probably the same as for the species (“Dans le source de Yeni Chehir, [...] entre Antioche et Alep, à l’intersection de la route d’Alexandrette” [at the source of Yenişehir, between Antakya and Aleppo, at the intersection of the road from İskenderun], Turkey).

***Melanopsis atramentaria* Pallary, 1939**

Original source. Pallary 1939: 91, pl. 5, fig. 29.

Type locality. “D’Acharné, sur l’Oronte, entre Hama et Kalâat el Moudik” [from Acharne at the Orontes, between Hama and Qal’at al Maḏīq], Syria.

† ?*Melanopsis attenuata* Sowerby in Fitton, 1836

Original source. Fitton 1836: 228, 346, pl. 22, fig. 5.

Type horizon. Weald Clay, early Cretaceous.

Type locality. “Punfield”, United Kingdom.

Remarks. After Sandberger (1870: 57) this species belongs in the genus *Goniobasis* Lea, 1862 (Pleuroceridae).

***Melanopsis nobilis* var. *attenuata* Pallary, 1912 [invalid]**

Original source. Pallary 1912: 21, fig. 43.

Type locality. “Des bords du Chott Djerid, à Tozeur” [banks of the Chott el Djerid at Tozeur], Tunisia.

Remarks. Junior homonym of *Melanopsis attenuata* Sowerby in Fitton, 1836.

***Melanopsis attenuata* Pallary, 1920 [invalid]**

Original source. Pallary 1920a: 29.

Type locality. “Tétouan”, Morocco.

Remarks. Junior homonym of *Melanopsis attenuata* Sowerby in Fitton, 1836.

***Melanopsis aurantiaca* Gassies, 1874**

Original source. Gassies 1874: 383.

Type locality. “Bourail et Nékété” [Bourail and Nakéty], New Caledonia.

† *Melanopsis austriaca* Handmann, 1882

Original source. Handmann 1882: 560.

Type horizon. Pannonian, zone D, late Miocene.

Type locality. “Kottingbrunn [...] Ziegelei a”, Austria.

Remarks. Wenz (1929: 2741) considered the taxon as a junior synonym of *M. haueri* Handmann, 1882.

† *Melanopsis avellana* Sandberger, 1870

Original source. Sandberger 1870–1875: pl. 5, figs 15–15a.

Type horizon. Maastrichtian, Cretaceous.

Type locality. “Auzas”, France.

Remarks. Plate 5 of Sandberger’s monograph was issued in 1870, while the description on p. 110 appeared in 1871 (Woodward 1906).

† *Melanopsis avellana* Fuchs, 1873 [invalid]

Original source. Fuchs 1873: 20, pl. 4, figs 16–17.

Type horizon. Middle Pannonian, late Miocene.

Type locality. “Sulzlacke bei Margarethen nächst Oedenburg [...]. Tinnye bei Ofen” [Sulzlacke near St. Margarethen (Burgenland, Austria); Tinnye (Hungary)].

Remarks. Junior homonym of *Melanopsis avellana* Sandberger, 1870. Pallary (1920b) introduced *M. ampla* as replacement name.

***Melanopsis lorcana* var. *aynensis* Azpeitia Moros, 1929**

Original source. Azpeitia Moros 1929: 146, pl. 2, figs 33–35.

Type locality. “Del río Mundo, cerca de Ayna (Albacete)” [in Mundo river, near Ayna, prov. Albacete], Spain.

† *Melanopsis baconica* Oppenheim, 1892

Original source. Oppenheim 1892: 770, pl. 34, figs 10–10b.

Type horizon. Late Santonian–early Campanian, late Cretaceous.

Type locality. “Csingerthal” (locality only given in plate captions) [near Ajka], Hungary.

Remarks. Bandel and Riedel (1994: 19) considered this taxon as a junior synonym of *Melanopsis ajkaensis* Tausch, 1886.

† *Melanopsis (Canthidomus) balatonensis* Cossmann, 1909

Original source. Cossmann 1909: 178.

Type horizon. Transdanubian, Pannonian, late Miocene.

Type locality. “Vörös-Bereny im Hohlweg nächst des Füzfő-major und in Kenese [...] Fonyód; [...] Szt-György-hegy in Hegymagys” (Halaváts 1903: 49) [Vörösberény, in the hollow-way near the Füzfő-major (?) and in Balatonkenese; Fonyód; Szent György-hegy in Hegymagas], Hungary.

Remarks. Replacement name for *M. boettgeri* Halaváts, 1903, non Klika, 1891.

† *Melanopsis banatica* Jekelius, 1944

Original source. Jekelius 1944: 132, pl. 50, figs 3–8, 12–14.

Type horizon. Early Pannonian, late Miocene.

Type locality. “Turislav-Tal bei Soceni” [Turislav valley near Soceni], Romania.

† *Melanoptychia banatica* Jekelius, 1944 [invalid]

Original source. Jekelius 1944: 139, pl. 58, figs 11–12.

Type horizon. Early Pannonian, late Miocene.

Type locality. “Turislav-Tal bei Soceni” [Turislav valley near Soceni], Romania.

Remarks. Junior secondary homonym of *Melanopsis banatica* Jekelius, 1944 (same work). Neubauer et al. (2014a: 462) considered both synonymous and gave priority to *Melanopsis banatica* (since *Melanoptychia* is considered as a synonym of *Melanopsis*).

***Melanopsis praemorsa bandeli* Schütt, 1988**

Original source. Schütt 1988b: 216, pl. 1, fig. 7.

Type locality. Unclear: Schütt referred to two different localities: “Wasserfall 1 km oberhalb der Straßenbrücke Jerash - Amman über den Zarqa” (p. 216) [waterfall 1 km above the bridge over the Zarqa’ along the road Jerash to Amman] and “Fluß Zerqa bei der alten Brücke am King Talal-See” (p. 219) [Zarqa’ river at the old bridge at the King Talal Dam], Jordan.

Types. Natural History Museum Vienna, Austria, coll. no. 85.544.

Remarks. Heller et al. (2005: 248) considered the variety as a junior synonym of *Melanopsis saulcyi* Bourguignat, 1853.

† ***Melanopsis banovaci* Brusina, 1902**

Original source. Brusina 1902: pl. 6, figs 67–68.

Type horizon. Middle Pannonian, late Miocene.

Type locality. “Kúp”, Hungary.

Types. Milan et al. (1974: 87) indicated a holotype, but it is uncertain whether the specimen was the only one Brusina had at hand (holotype by monotypy, Art. 73.1.2). The specimen is stored in the Croatian Natural History Museum, Zagreb, coll. no. 2527-173.

***Melanopsis barbini* Pallary, 1911**

Original source. Pallary 1911: 129, [unnumbered plate], figs 4–5.

Type locality. “Tout près d’Oudjda, à 4 kilom. S.-E., sourdent les belles sources de Sidi-Yahia qui alimentent une véritable oasis, puis la ville d’Oudjda, et vont finalement se déverser dans l’oued Isly” [near Oujda, 4 km southeast, at the sources of Sidi Yahya that feed an oasis and the city of Oujda, and ultimately will flow into the Oued Isly], Morocco.

† ***Melanopsis barthai* Bandel, 2000**

Original source. Bandel 2000: 160, figs 43–44, 46–51.

Type horizon. Late Pannonian, late Miocene.

Type locality. “Sand pit near Papkesi [Papkeszi]”, Hungary.

Types. Geological-Palaeontological Institute and Museum University of Hamburg, coll. no. 4268.

† *Melanopsis bartolinii* Capellini, 1873

Original source. Capellini 1873: 550, pl. 8, figs 1–4.

Type horizon. Messinian, late Miocene.

Type locality. “Sterza di Laiatico”, Italy.

Remarks. Illustration not on pl. 7 as indicated by Capellini (1873). The name “*bartolini*” as mentioned in Ligios et al. (2012: 358) is an incorrect subsequent spelling.

Melanopsis bedei Pallary, 1933

Original source. Pallary 1933: 248.

Type locality. “L’Oued Bou Regreg, au pont des Seouls” [in the Oued Bou Regreg, at the bridge of the Séouls (?)], Morocco.

† *Melanopsis belestensis* Villatte, 1952

Original source. Villatte 1952: 329, pl. 2, figs 16–21.

Type horizon. Campanian, Cretaceous.

Type locality. “Bélesta (Ariège)”, France.

Melanopsis belonidaea Bourguignat, 1884

Original source. Bourguignat 1884: 110.

Type locality. “Ruisseau d’eau chaude à Ouargla (prov. de Constantine) et eaux thermales du Djérid, au nord du chott Tiraoun (sud de la Tunisie)” [in a warm water stream in Ouargla (Algeria) and thermal waters of Djerid, north of chott Tiraoun (southern Tunisia)].

Remarks. Multiple spellings occur in the original work: “*belonidoea*” on p. 110 but “*belonidaea*” on p. 75. Apparently, the spelling on p. 110 is based on a typesetting mistake regarding the ligature (“œ” instead of “æ”). Letourneux & Bourguignat (1887) acted as First Reviser sensu Art. 24.2.2, giving the name as “*belonidaea*”. The spellings “*balonidaea*” mentioned in Westerlund (1886: 125) and “*belonidae*” given in Pallary (1912: 14, 19, 22; 1926: 75) are incorrect spellings.

Melanopsis belusi Bourguignat, 1884

Original source. Bourguignat 1884: 134.

Type locality. “Du Bélus, près de Saint-Jean-d’Acre (Syrie)” [in the Na’aman river, near Acre], Israel.

Remarks. Bourguignat (1884) denoted the authority as “Letourneux, 1882”, but there is no evidence that the description really derived from that author. Heller et al. (2002: 596) considered the species as a junior synonym of *Melanopsis costata* (Olivier, 1804). Heller et al. (2005: 244) in turn treated it as a junior synonym of *Melanopsis lamprea* Bourguignat, 1884.

***Melanopsis belusiensis* Germain, 1921 [invalid]**

Original source. Germain 1921: 463.

Remarks. Unjustified emendation and therefore junior objective synonym of *M. belusi* Bourguignat, 1884.

† *Melanopsis (Calodiona) bergeroni* Stefanescu, 1896

Original source. Stefanescu 1896: 131, pl. 11, figs 32–37.

Type horizon. Sienisian to Pelendavian, Pliocene.

Type locality. “À Gura-Motrului et à Bocovatz, dans la vallée de Jiu” [at Gura-Motrului and at Bucovăț, in the valley of the river Jiu], Romania.

***Melanopsis magnifica* var. *berkanensis* Pallary, 1911**

Original source. Pallary 1911: 128, [unnumbered plate], fig. 12.

Type locality. “Près du village de Berkane, à la lisière Sud-Ouest de la plaine des Triffas, tout au Nord des Beni-Znassen, [...] source connue sous le nom berbère d’Aoûllout” [near Berkane, to the southwestern edge of the plain of Triffa, just north of Beni Snassene, in a spring known as Aïn Aoullout], Morocco.

***Fagotia berlani* Bourguignat, 1884**

Original source. Bourguignat 1884: 34.

Type locality. “Le Danube à Ibraila; la Save entre Agram et Sissek” [Danube river at Brăila (Romania); Sava river between Zagreb and Sisak (Croatia)].

Remarks. Note that Bourguignat denoted the authority as “Bourguignat, 1880”.

***Melanella berlani* Bourguignat, 1884**

Original source. Bourguignat 1884: 22.

Type locality. “Le Danube près Ibraila” [Danube river at Brăila], Romania.

Remarks. Note that Bourguignat denoted the authority as “Bourguignat, 1879”.

† *Melanopsis textilis* var. *bicarinata* Handmann, 1887

Original source. Handmann 1887: 16.

Type horizon. Pannonian, zone B–D, late Miocene.

Type locality. “Leobersdorf”, Austria.

Melanopsis mellalensis var. *bicarinata* Pallary, 1928 [invalid]

Original source. Pallary 1928b: 19, pl. 2, figs 22–23.

Type locality. “De Béni Mellal; de l’oued Dai” [from Beni Mellal; from Oued Dai (?)], Morocco.

Remarks. Junior homonym of *Melanopsis textilis bicarinata* Handmann, 1882 (see Note 1).

† *Melanopsis doboi bicarinata* Schréter, 1975 [invalid]

Original source. Schréter 1975: 8, pl. 2, figs 7–10.

Type horizon. Riss/Würm end to early Würm Ice Age, Pleistocene.

Type locality. “Eger, az egri vár Zárkándy bástyájának átmetszése” [Eger, section at the Zarkandy bastion of the fortress Eger], Hungary.

Types. Magyar Állami Földtani Intézet (Hungarian Geological Museum), Budapest; no number indicated.

Remarks. Junior homonym of *Melanopsis textilis bicarinata* Handmann, 1882 (see Note 1). Neubauer et al. (2016a) attributed the species *M. doboi* to the genus *Microcolpia*.

† *Melanopsis bicincta* Blanckenhorn, 1897

Original source. Blanckenhorn 1897: 119, pl. 9, figs 35–40.

Type horizon. Plio-Pleistocene.

Type locality. “In der zweiten Thonbank des linken Orontesufers und der ersten und zweiten des rechten Ufers” [in the second clay bank at the left riverside of the Orontes and the first and second bank of the right riverside], Syria?

Remarks. Introduced as “n. mut.” but clearly as a binomen and hence not infrasubspecific in the sense of ICZN Art. 45.6.

† *Melanopsis tothi* var. *bicingulata* Brusina, 1903

Original source. Brusina 1903: 115.

Type horizon. Late Pleistocene–early Holocene.

Type locality. “Bischofsbad” [Püspökfürdő, Băile 1 Mai, Lake Peșea], Romania.

Remarks. Neubauer et al. (2014d: 125) considered this taxon as a junior synonym of *Microcolpia parreyssii sikorai* (Brusina, 1903).

***Melanopsis praeros* [sic] var. *bicolorata* Paetel, 1888**

Original source. Paetel 1888: 403.

Type locality. “W. of Shiraz” (Nevill 1884: 209), Iran.

Remarks. Originally introduced as infrasubspecific taxon (“subvariety”) by Nevill (1884), but made available by Paetel (1888) who treated it as variety (Art. 45.5.1). Paetel clearly referred to the description of Nevill.

† *Melanopsis (Martinia) capulus* var. *biconica* Handmann, 1887

Original source. Handmann 1887: 21, pl. 2, figs 16–18.

Type horizon. Pannonian, zone B–D, late Miocene.

Type locality. “Leobersdorf”, Austria.

† *Melanopsis bicoronata* Brusina, 1884

Original source. Brusina 1884b: 56.

Type horizon. Early Langhian, middle Miocene.

Type locality. “Stuparuša” (p. 47) [near Sinj], Croatia.

Types. Milan et al. (1974: 87) stated that Brusina (1897) had indicated one of the specimens illustrated by him (pl. 4, fig. 15) as type. However, it is uncertain whether the specimen actually derives from the original type series. The specimen is stored in the Croatian Natural History Museum, Zagreb, coll. no. 2975-621/1.

† *Melanopsis hazayi* var. *bifilosa* Brusina, 1903 [unresolved]

Original source. Brusina 1903: 113.

Type horizon. Late Pleistocene–early Holocene.

Type locality. “Bischofsbad” [Püspökfürdő, Băile 1 Mai, Lake Peșea], Romania.

Remarks. Brusina (1903) introduced several varieties with this name, apparently considering it only a descriptive term; the homonymy issue needs to be solved by a First Reviser. Currently, all of them are considered junior synonyms of *Microcolpia parreyssii sikorai* (Brusina, 1903) (Neubauer et al. 2014d: 125).

† *Melanopsis sikorai* var. *bifilosa* Brusina, 1903 [unresolved]

Original source. Brusina 1903: 112.

Type horizon. Late Pleistocene–early Holocene.

Type locality. “Bischofsbad” [Püspökfürdő, Băile 1 Mai, Lake Pețea], Romania.

Remarks. Brusina (1903) introduced several varieties with this name, apparently considering it only a descriptive term; the homonymy issue needs to be solved by a First Reviser. Currently, all of them are considered junior synonyms of *Microcolpia parreyssii sikorai* (Brusina, 1903) (Neubauer et al. 2014d: 125).

† *Melanopsis themaki* var. *bifilosa* Brusina, 1903 [unresolved]

Original source. Brusina 1903: 110.

Type horizon. Late Pleistocene–early Holocene.

Type locality. “Bischofsbad” [Püspökfürdő, Băile 1 Mai, Lake Pețea], Romania.

Remarks. Brusina (1903) introduced several varieties with this name, apparently considering it only a descriptive term; the homonymy issue needs to be solved by a First Reviser. Currently, all of them are considered junior synonyms of *Microcolpia parreyssii sikorai* (Brusina, 1903) (Neubauer et al. 2014d: 125).

† “*Melanopsis tothi* var. *bifilosa*” mentioned in Brusina (1903: 114) [unavailable]

Horizon. Late Pleistocene–early Holocene.

Locality. “Bischofsbad” [Püspökfürdő, Băile 1 Mai, Lake Pețea], Romania.

Remarks. Nomen nudum (Brusina apparently considered the term self-explanatory). Brusina (1903) introduced several varieties with this name, apparently considering it only a descriptive term. Neubauer et al. (2014d: 125) considered this taxon as a junior synonym of *Microcolpia parreyssii sikorai* (Brusina, 1903).

Hemisinus acicularis var. *biharensis* Clessin, 1890

Original source. Clessin 1887–1890: 683, fig. 467.

Type locality. “In der warmen Quelle bei Robogany im Bihargebirge in Ungarn” [in the thermal spring near Răbăgani in the Bihar Mts], Romania.

Remarks. Based on an “in schedis” name from Hazay.

***Melanopsis crenocarinata* var. *bilineata* Moricand, 1841**

Original source. Moricand 1841: 62.

Type locality. “Rio de Pedra Branca, procince de Bahia” [Pedra Branca river, province Bahia], Brazil.

Remarks. Although not explicitly stated, this variety was apparently considered to belong to the new genus *Verena* by Adams and Adams (1854) (Thiaridae), of which *Melanopsis crenocarinata* is the type species (see Nuttall 1990: 253).

† ***Melanopsis biliottii* Bukowski, 1892**

Original source. Bukowski 1892: 249.

Type horizon. Apolakkia/Monolithos Formation, Pliocene.

Type locality. “Rhodos” (locality specified as “Monolithos” in Bukowski 1893), Greece.

Remarks. The names “*billiottii*” and “*biolittii*” as mentioned in Pallary (1926a: 76) and Wenz (1929: 2667) are incorrect subsequent spellings.

† ***Melanopsis binodosa* Blanckenhorn, 1897**

Original source. Blanckenhorn 1897: 117, pl. 9, figs 28–34.

Type horizon. Plio-Pleistocene.

Type locality. “Aus der zweiten Thonbank des rechten und linken Orontesufers; obere Thonbank des rechten Ufers” [from the second clay bank of the right and left riversides of the Orontes; upper clay bank of the right riverside], Syria?

Remarks. Introduced as “n. mut.” but clearly as a binomen and hence not infraspecific in the sense of ICZN Art. 45.6.

***Melanopsis bullio* var. *bipartita* Dautzenberg, 1894**

Original source. Dautzenberg 1894: 344.

Type locality. “Lac de Homs”, Syria.

† ***Melanopsis bittneri* Fuchs, 1877**

Original source. Fuchs 1877: 40, pl. 4, figs 12–17.

Type horizon. Chaudian, late Pleistocene.

Type locality. “Livonates bei Talandi” (p. 36) [Livanates], Greece.

† *Melanoptychia bittneri* Neumayr, 1880 [invalid]

Original source. Neumayr 1880c: 480, pl. 7, fig. 11.

Type horizon. Langhian, middle Miocene.

Type locality. “Žepj” [Džepi], Bosnia and Herzegovina.

Remarks. Neubauer et al. (2013: 135) considered *Melanoptychia* as a junior synonym of *Melanopsis* and this species as a junior secondary homonym of *Melanopsis bittneri* Fuchs, 1877. They introduced *M. medinae* as replacement name (see also *Melanoptychia carusi* Brusina, 1902).

† *Melanopsis bittneri* Brusina, 1902 [invalid]

Original source. Brusina 1902: pl. 7, figs 18–21.

Type horizon. Langhian, middle Miocene.

Type locality. “Džepe” [Džepi], Bosnia and Herzegovina.

Remarks. Junior homonym of *Melanopsis bittneri* Fuchs, 1877. Neubauer et al. (2013: 134) considered it as a junior synonym of *M. cvijici* Brusina, 1902.

† *Melanopsis blanchardi* Brusina, 1892

Original source. Brusina 1892: 143.

Type horizon. Middle Pannonian, late Miocene.

Type locality. “Markuševac”, Croatia.

Types. Milan et al. (1974: 100) indicated a holotype, but it is uncertain whether the specimen actually derives from the original type series and whether it was the only specimen Brusina had at hand. The specimen is stored in the Croatian Natural History Museum, Zagreb, coll. no. 2539-185.

† *Melanopsis blanckenhorni* Wenz, 1929

Original source. Wenz 1929: 2667.

Type horizon. Pleistocene.

Type locality. “Dans le lac d’Antioche, dans l’oued Baradah près Aïn Fidji, et dans l’Aïn Plaça, fontaine de la plaine du Bahr-el-Houlé” (Bourguignat 1884: 82) [in Lake Anuk (also as Amik), in the river Barada near Aïn al-Fiji, and in Aïn el Placa, a spring in the plains of the Hula valley], Syria.

Remarks. Replacement name for *M. prophetarum minor* Bourguignat, 1884, non Grateloup, 1838 (see Note 1). The type locality given by Wenz (“U. Orontestal, 7 km unterhalb Antâkîje”) is incorrect.

† *Melanopsis blanckenhorni* Schütt, 1988 [invalid]

Original source. Schütt 1988a: 137, pl. 3, figs 18–20.

Type horizon. Plio-Pleistocene.

Type locality. “Höhere Lage am westlichen Orontes-Ufer 12 km S Ġisir aš-Šugur” [western riverside of the Orontes, 12 km south of Jisr Ash-Shughur], Syria.

Types. Senckenberg Forschungsinstitut und Naturmuseum Frankfurt, coll. no. SMF 307206.

Remarks. Junior homonym of *Melanopsis blanckenhorni* Wenz, 1929. Heller and Sivan (2001: 131) considered the species as junior synonym of *Melanopsis costata* (Olivier, 1804).

† *Purpuroidea blaschkei* Schenk, 1969

Original source. Schenk 1969: 39, pl. 1, fig. 15.

Type horizon. Late Santonian to early Campanian, late Cretaceous.

Type locality. “Zöttbachalm bei Brandenburg, Tirol” [Zöttbachalm near Brandenburg, Tyrol], Austria.

Remarks. The species was originally attributed to the marine genus *Purpuroidea* Lyett, 1848 (Purpuroideidae), but considered as a junior synonym of the melanopsid *Megalonoda spiniger* (Sowerby in Sedgwick & Murchison, 1832) by Kollmann (1984: 58).

Melanopsis bleicheri Paladilhe, 1874

Original source. Paladilhe 1874: 403, pl. 8, figs 23–25.

Type locality. “Environs d’Oran” [surroundings of Oran], Algeria.

† *Melanopsis bleunardi* Porumbaru, 1881

Original source. Porumbaru 1881: 29, pl. 9, fig. 6.

Type horizon. Early Romanian, Pliocene.

Type locality. “Bucovatzu” [Bucovăț], Romania.

Remarks. Wenz (1929) erroneously gave the town of Craiova as type locality, near which Bucovăț lies.

† *Hydrobia bodeica* Tausch, 1886

Original source. Tausch 1886: 12, pl. 1, figs 47a–c.

Type horizon. Late Santonian–early Campanian, late Cretaceous.

Type locality. “Csingerthal bei Ajka” [Csinger valley near Ajka], Hungary.

Remarks. Bandel and Riedel (1994: 20) considered this taxon as a juvenile specimen and as a junior synonym of *Melanopsis ajkaensis* Tausch, 1886.

† *Melanopsis boettgeri* Klika, 1891

Original source. Klika 1891: 112, fig. 111a–d.

Type horizon. Oligocene.

Type locality. “Wärzen; [...] Tuchořic” [Dvërce; Tuchořice], Czech Republic.

† *Melanopsis conemenosiana* var. *boettgeri* Oppenheim, 1891 [invalid]

Original source. Oppenheim 1891: 470.

Type horizon. Plio-Pleistocene.

Type locality. “Preveza in Epirus”, Greece.

Remarks. Junior homonym of *Melanopsis boettgeri* Klika, 1891. Pallary (1920b: 112) introduced *M. conemenosiana* var. *turritella* as replacement name.

† *Melanopsis (Pauluccia) boettgeri* Brusina, 1902 [invalid]

Original source. Brusina 1902: pl. 29, figs 5–8.

Type horizon. Transdanubian, Pannonian, late Miocene.

Type locality. “Radmanest” [Rădmănești], Romania.

Types. The syntypes are stored in the Croatian Natural History Museum, Zagreb; no number indicated (Milan et al. 1974: 99).

Remarks. Junior homonym of *Melanopsis boettgeri* Klika, 1891. Pallary (1920b) introduced *M. delicata* as replacement name.

† *Melanopsis boettgeri* Halaváts, 1903 [invalid]

Original source. Halaváts 1903: 49, pl. 2, fig. 14.

Type horizon. Transdanubian, Pannonian, late Miocene.

Type locality. “Vörös-Bereny im Hohlweg nächst des Füzfő-major und in Kenese [...] Fonyód; [...] Szt-György-hegy in Hegymagyos” [Vörösberény, in the hollow-way near the Füzfő-major (?) and in Balatonkenese; Fonyód; Szent György-hegy in Hegymagas], Hungary.

Remarks. Junior homonym of *Melanopsis boettgeri* Klika, 1891. Cossmann (1909) introduced *Canthidomus balatonensis* as replacement name.

***Melanopsis bofilliana* Bourguignat, 1884**

Original source. Bourguignat 1884: 101.

Type locality. “Environs de Lorca, en Espagne” [surroundings of Lorca], Spain.

Remarks. The name “*bofilliana*” as mentioned in Westerlund (1886: 126) is an incorrect subsequent spelling.

† ***Melanopsis bogdanowi* Brusina, 1892**

Original source. Brusina 1892: 142.

Type horizon. Middle Pannonian, late Miocene.

Type locality. “Markuševac”, Croatia.

Types. The illustrated syntypes are stored in the Croatian Natural History Museum, Zagreb, coll. no. 2538-184/1-3 (Milan et al. 1974: 100).

Remarks. The name “*bogdanovi*” as mentioned in Wenz (1929: 2868) is an incorrect subsequent spelling.

“*Melanopsis bondeensis* Gassies” mentioned in Paetel (1888: 399) and Pallary (1926a: 76) [unavailable]

Locality. “Bondé”, New Caledonia.

Remarks. Nomen nudum, found only in species lists by Paetel (1888) and Pallary (1926a). Perhaps confused with “*Bulimus bondeensis*” from the same locality, which Gassies (1871: 203) listed in the plate captions right below the *Melanopsis* species.

† ***Melanopsis bonellii* Manzoni, 1870**

Original source. Manzoni 1870: 498, pl. 3, figs 8–9.

Type horizon. Late Miocene to Pliocene.

Type locality. “Mte. Gibio nel Modanese ed a St. Agata nel Tortonese; [...] a Sogliano” [Mt. Gibio, Sant’Agata Fossili, Sogliano al Rubicone], Italy.

Remarks. In the old literature the name appears frequently as “*M. bonellii* Sismonda”. The name is not available from Sismonda (1847: 55), however, who solely referred to a misidentified name (*M. carinata* non Sowerby) in the unpublished museum catalogue of Bonelli (1827) but did not supply a description. It appears as a nomen nudum also in d’Orbigny (1852: 28) and Doderlein (1863: 16). Hörnes (1856: 595) mentioned the name along with two other melanopsids from Italy and gave a collective description for all of them (translated from German: “all individuals are smaller as the Viennese [*M. martiniana*], as well as more elongate, but they have the characteristic marginal bulge and the typical keel below the suture”). Following Art. 12.1, every

name [...] must be accompanied by a description or a definition of the taxon that it denotes [...], which is not the case for the mentioning of *M. bonellii* in Hörnes (1856) because the description given there refers to a group of taxa and not a single taxon. Manzoni (1870) made the name available by providing an illustration (Art. 12.2.7).

The name “*bonelli*” as mentioned in Sacco (1889: 65) and Syrides (1998: 173) is an incorrect subsequent spelling.

***Melanella letourneuxi* var. *bosnica* Servain, 1884**

Original source. Servain 1884: 379.

Type locality. “Migliaska” [Miljacka river near Sarajevo], Bosnia and Herzegovina.

† ***Melanopsis bouei* Férussac, 1823**

Original source. Férussac 1819–1851: *Mélanopsides fossiles*, pl. 2 (1823), figs 9–10 or Férussac 1823: 159, pl. 8, figs 9–10 (precedence not established; see introduction for discussion).

Type horizon. Pannonian, late Miocene.

Type locality. “Près de Bisentz, dans la vallée de la Marsch, en Moravie; [...] près de Scharditz” [near Bzenec, in the valley of the March river, in Moravia; near Šardice], Czech Republic.

Types. Papp (1953b) designated the specimen illustrated in Férussac (1823, pl. 8, fig. 9) as lectotype (after Art. 74.5). Whereabouts of type specimen unknown.

***Fagotia bourguignati* Bourguignat, 1884**

Original source. Bourguignat 1884: 48.

Type locality. “Dans la rivière de Krapina et dans celle entre Plaski et Ostaria, en Croatie” [in the Krapina river and that between Plaški and Oštarije], Croatia.

Remarks. Bourguignat denoted the authority as “Letourneux, 1884”, but there is no evidence that the description really derived from that author. Starobogatov et al. (1992: 60) considered the species as a junior synonym of *Fagotia* [= *Esperiana*] *esperi* (Férussac, 1823).

***Melanella bourguignati* Bourguignat, 1884**

Original source. Bourguignat 1884: 26.

Type locality. “Rivière d’Ogulin (Croatie); la Save, à Sissek (Slavonie)” [river at Ogulin; Sava river at Sisak], Croatia.

Remarks. Bourguignat denoted the authority as “Letourneux, 1879”, but there is no evidence that the description really derived from that author.

***Melanopsis bovieri* Pallary in Germain, 1921**

Original source. Germain 1921: 504, pl. 20, figs 7–8.

Type locality. “Le Nahr ez Zaïr (Liban)” [locality not found], Lebanon.

Remarks. Heller et al. (2005: 245) considered the species as a junior synonym of *Melanopsis lamprea* Bourguignat, 1884.

***Melanopsis brachymorpha* Pallary, 1936**

Original source. Pallary 1936: 58, pl. 3, fig. 4.

Type locality. “Aïn Akseri Ifesfassen à Imouzer d’Agadir” [Aïn Akseri Ifesfassen at Imouzzer near Agadir], Morocco.

† *Melanopsis brachyptycha* Neumayr, 1880

Original source. Neumayr 1880c: 478, pl. 7, fig. 3.

Type horizon. Pannonian, late Miocene.

Type locality. “Posušje”, Bosnia and Herzegovina.

Remarks. The name “*brachyptychia*” as mentioned in Wenz (1929: 2682) is an incorrect subsequent spelling.

***Melanopsis brasiliensis* Moricand, 1838**

Original source. Moricand 1838: 144, pl. 3, figs 12–13.

Type locality. “Près de Villa de Barra” [near Villa de Barra, province Bahia], Brazil.

Remarks. Currently considered to belong in the genus *Hemisinus* Swainson, 1840 (Thiaridae) (Nuttall 1990: 244).

† *Melanopsis braueri* Neumayr in Neumayr & Paul, 1875

Original source. Neumayr and Paul 1875: 43, pl. 8, figs 26–27.

Type horizon. Cernikian, Pliocene.

Type locality. “Graben hinter der Kirche von Podwin; [...] Graben zwischen der Čapla und Podwin; [...] Čapla; [...] Gromačnik; [...] Strassengraben zwischen Gromačnik und Sibin; [...] Cigelnik” [ditch behind the church of Podvinje; Čaplja

trench near Slavonski Brod; Čaplja; Gromačnik; roadside ditch between Gromačnik and Sibinj; Ciglenik], Croatia.

† *Melanopsis pterochila* var. *breastensis* Stefanescu, 1896

Original source. Stefanescu 1896: 130, pl. 11, figs 26, 28, 30–31.

Type horizon. Early Cernikian, early Pliocene.

Type locality. “À Breasta, à Bocovatz, à Glodu et à Bâzdâna, dans la vallée de Jiu, et à Beceni, dans la vallée de Slanic, district de Buzau” [at Breasta, at Bucovăț, at Glodu and at Bâzdâna, in the valley of the river Jiu, and at Beceni, in the valley of the river Slănic, Buzau county], Romania.

Melanopsis brevicula Pallary, 1918

Original source. Pallary 1918: 150.

Type locality. “Agourai, Aïn Mahrouf” [Agourai, Oued Ain Maarouf], Morocco.

† *Melanopsis brevis* Sowerby, 1826

Original source. Sowerby 1826–1829: 42, pl. 523, fig. 2.

Type horizon. Eocene.

Type locality. “Upon the Hampshire coast”, United Kingdom.

Remarks. After Cossmann (1909: 153) this species belongs in the genus *Coptostylus* Sandberger, 1872 (Thiaridae).

Melanopsis brevis Parreyss in Mousson, 1854 [invalid]

Original source. Mousson 1854: 51.

Type locality. “Dans les eaux de l’ancien Léonthes” [in the Litani river], Lebanon.

Remarks. Junior homonym of *Melanopsis brevis* Sowerby, 1826. Blanckenhorn (1897) introduced *M. maroccana media* as replacement name (see also *M. moussoni* Pallary, 1916). The authority is clear from the discussion provided in Mousson (1854); in fact, Mousson even seemed to doubt the specific separation of this taxon from other melanopsids.

Melanopsis brevis Morelet, 1857 [invalid]

Original source. Morelet 1857: 32.

Type locality. “[Ad Sanctam-Mariam de Balade]” [Balade], New Caledonia.

Remarks. Junior homonym of *Melanopsis brevis* Sowerby, 1826. Pallary (1916: 81) introduced *M. moreleti* as replacement name.

† *Melanopsis* (?*Macrospira*) *brevis* Doncieux, 1908 [invalid]

Original source. Doncieux 1908: 205, pl. 11, figs 13a–b.

Type horizon. Middle Lutetian, Eocene.

Type locality. “Au Nord d’Albas”, France.

Remarks. Junior homonym of *Melanopsis brevis* Sowerby, 1826. Pallary (1916: 81) introduced *M. abbreviata* as replacement name, which is itself a junior homonym of *M. abbreviata* Brusina, 1874 (see *M. doncieuxi* Wenz, 1919 and *M. atacica* Wenz, 1928).

Melanopsis graellsii var. *brevis* Pallary, 1924 [invalid]

Original source. Pallary 1924: 250.

Type locality. “Navajas (Castellón), Alcudia, Alberrique, Jativa (Valencia)” [Navajas, L’Alcúdia, Alberic, Xàtiva (prov. Valencia)], Spain.

Remarks. Junior homonym of *Melanopsis brevis* Sowerby, 1826. Not available from Rossmässler (1854), to whom Pallary referred to, because he did not use the term “*brevis*” to denote a species-group taxon but only cited Férussac’s (1823) description of an unnamed variety of *M. dufourii*.

† *Melanopsis* (*Melanosteira*) *aetolica brevisesta* [sic] Papp & Thenius, 1952

Original source. Papp and Thenius 1952: 4.

Type horizon. Gelasian, early Pleistocene.

Type locality. Not stated in Papp and Thenius (1952), but specified by Papp (1955) as “Angelocastron, (Agrinion)” [Angelókastro], Greece.

Types. Museum of Palaeontology and Geology of the University of Athens; no number indicated (Papp 1955).

Remarks. Although probably unintentionally, Papp and Thenius (1952) briefly described and thus validly introduced the taxon. The spelling “*brevisesta*” is probably a lapsus calami for “*brevitesta*” (Latin for “short shell”), as used in the re-description by Papp (1955: 126, pl. 20, figs 10–14). Nevertheless, “*brevisesta*” is the correct name following Art. 32.5.1 and “*brevitesta*” is an incorrect subsequent spelling.

Melanopsis jordanica var. *breviuscula* Bourguignat, 1884

Original source. Bourguignat 1884: 142.

Type locality. “Lac de Tibériade” [Sea of Galilee], Israel.

† *Melanopsis briarti* Cossmann, 1888

Original source. Cossmann 1888: 282.

Type horizon. Late Danian, Paleocene.

Type locality. “Mons” (Briart and Cornet 1873: 71), Belgium.

Remarks. Introduced for *M. buccinoidea* sensu Briart & Cornet, 1873, “non Fé-russac” (actually it should read “non Olivier, 1801”).

† *Melanopsis briarti* Munier-Chalmas, 1897 [invalid]

Original source. Munier-Chalmas 1897: 86.

Type horizon. Late Danian, Paleocene.

Type locality. “Mons” (Briart and Cornet 1873: 71), Belgium.

Remarks. Introduced for *M. buccinoidea* sensu Briart & Cornet, 1873, “non Fé-russac” (actually it should read “non Olivier, 1801”). Junior homonym, as well as junior objective synonym of *Melanopsis briarti* Cossmann, 1888. Obviously, Munier-Chalmas overlooked that the name had already been introduced for the very same misidentified taxon by Cossmann.

† *Melania Brocchii* Michelotti, 1847

Original source. Michelotti 1847: 189.

Type horizon. Late Miocene.

Type locality. “Près de Tortone” [near Tortona], Italy.

Remarks. Wenz (1929: 2879) considered it a junior synonym of *Ptychomelania buccinella* Sacco, 1895 (Thiaridae).

† *Melanopsis brongniarti* Locard, 1883

Original source. Locard 1883b: 99.

Type horizon. Mammal zone MN 15, Pliocene.

Type locality. “Montgardon”, France.

† *Melanopsis broti* Neumayr, 1880

Original source. Neumayr 1880b: 295, pl. 1, fig. 29.

Type horizon. Plio-Pleistocene.

Type locality. “Zwischen Pyllé und Antimachia” [between Pýli and Antimácheia, Kos Island], Greece.

Remarks. Pallary (1925) and Wenz (1929) erroneously referred to it as homonym of “*M. broti*” Gassies, 1874, which was actually introduced as *M. brotiana*. The replacement name *M. cosiana* Pallary, 1925 is thus invalid as it is a junior objective synonym (see also Willmann 1981).

***Melanopsis brotiana* Gassies, 1874**

Original source. Gassies 1874: 386.

Type locality. “La Conception, prope Noumea” [Conception, near Nouméa], New Caledonia.

† ?*Melanopsis brownii* Etheridge, 1879

Original source. Etheridge 1879: 87, pl. 7, fig. 4.

Type horizon. Pebasian, late Cenozoic.

Type locality. “Canama”, Peru.

Remarks. Currently considered to belong in the genus *Verena* H. Adams & A. Adams, 1854 (Thiaridae) (Nuttall 1990: 256). The name “*browni*” as given by Nuttall (1990) is an incorrect subsequent spelling.

† *Melanopsis brusinai* Lörenthey, 1902

Original source. Lörenthey 1902: 223, pl. 16, fig. 7.

Type horizon. Middle Pannonian, late Miocene.

Type locality. “Tinnye”, Hungary.

† *Melanoptychia brusinai* Jekelius, 1944 [invalid]

Original source. Jekelius 1944: 137, pl. 56, figs 1–23.

Type horizon. Early Pannonian, late Miocene.

Type locality. “Turislav-Tal bei Soceni” [Turislav valley near Soceni], Romania.

Remarks. Junior secondary homonym and junior synonym of *Melanopsis brusinai* Lörenthey, 1902 (see Neubauer et al. 2014a: 462).

† *Melanopsis (Homalia) bucciniformis* Handmann, 1887

Original source. Handmann 1887: 14, pl. 1, figs 8–9.

Type horizon. Pannonian, zone B–D, late Miocene.

Type locality. “Leobersdorf”, Austria.

Remarks. Wenz (1929: 2813) considered the taxon as a junior synonym of *Melanopsis pygmaea* Hörnes, 1856.

Melania buccinoidea Olivier, 1801

Original source. Olivier 1801: 297, pl. 17, fig. 8.

Type locality. “De Scio, de presque toutes les îles de l’Archipel, de Crète, de Syrie” [from Chios, from almost all islands of the Archipelago, from Crete (Greece), from Syria].

Remarks. Today combined as *Melanopsis buccinoidea*, the species is one of the few stable taxa in melanopsid taxonomy (e.g., Glaubrecht 1996).

† *Melanopsis buccinulum* Melleville, 1843

Original source. Melleville 1843: 95, pl. 4, figs 13–15.

Type horizon. Thanetian, Paleocene.

Type locality. “Châlons” [Châlons-sur-Vesle], France.

Remarks. Perhaps unaware of Melleville’s publication, Deshayes (1862: 469, pl. 31, figs 11–13) erroneously described this species as new: he marked the name with “Desh.”, provided a Latin diagnosis and did not list any citations. Judging from the illustrations in both works, Deshayes’ species is undoubtedly the same as Melleville’s. Moreover, both were recorded from the same locality (Châlons-sur-Vesles).

Melanopsis tingitana var. *bucheti* Pallary, 1899

Original source. Pallary 1899: 147, pl. 9, fig. 6.

Type locality. “L’O.[ued] Ida ou Guert, près de Mogador” [Wadi Ida Ou Gourdh at Essaouira], Morocco.

† *Melanopsis bukovaci* Brusina, 1902

Original source. Brusina 1902: pl. 6, figs 8–10.

Type horizon. Middle–late Cernikian, late Pliocene–early Pleistocene.

Type locality. “Gromačnik”, Croatia.

Types. The illustrated syntypes are stored in the Croatian Natural History Museum, Zagreb, coll. no. 2506-152/1-3 (Milan et al. 1974: 87).

***Bithynia bulgarica* Drensky, 1947**

Original source. Drensky 1947: 45.

Type locality. “Познат за сега само от р. Дунав, северно от гр. Лом” [from the river Danube, near the city of Lom], Bulgaria.

Remarks. Angelov (2000: 15), while incorrectly treating the name as nomen nudum, considered the species as a junior synonym of *Amphimelania holandri* [sic] (Pfeiffer, 1828).

***Melanopsis bullio* Kobelt, 1879**

Original source. Kobelt 1879–1880: 17, pl. 188, figs 1902–1903.

Type locality. Not indicated, probably in the Middle East.

Remarks. Introduced in synonymy of *M. costata* by Kobelt (1879), referring to an “in schedis” name by Parreyss. Dautzenberg (1894) made the name available by treating it as valid name (see Note 2).

† *Melanopsis calamonensis* Magrograssi, 1928

Original source. Magrograssi 1928: 261, pl. 6, fig. 22.

Type horizon. Plio-Pleistocene.

Type locality. “Rodi: colline sulla sinistra del fiume Dimilia” [Rhodes island: hills on the left bank of the river Dimilia], Greece.

***Melanopsis callichroa* Bourguignat, 1884**

Original source. Bourguignat 1884: 91.

Type locality. “Dans l’intérieur de la grotte du Nahr-el-Kelb, près de Beyrouth (Syrie)” [in the cave of Nahr el-Kalb near Beirut], Lebanon.

***Melanopsis callista* Bourguignat, 1884**

Original source. Bourguignat 1884: 118.

Type locality. “Sadjour-Sou entre Ain-Taïb et Alep” [at Sadjour-Sou between Gaziantep (Turkey) and Aleppo (Syria)].

Remarks. Heller et al. (2005: 243) tentatively considered the species as a junior synonym of *Melanopsis costata* (Olivier, 1804).

† *Melanopsis callosa* Voltz, 1852

Original source. Voltz 1852: viii, pl. 3, fig. 10.

Type horizon. Mammal zone MN 1, early Miocene.

Type locality. “Weisenau”, Germany.

Remarks. The name appeared first as nomen nudum in Genth (1848: 198–199), Sandberger (1850: 18; 1851: 676; 1852: 684) and Braun in Walchner (1851: 1126). Wenz (1929: 2764) considered this taxon as a junior synonym of *Melanopsis fritzei* Thomä, 1845.

† *Melanopsis camptogramma* Brusina, 1876

Original source. Brusina 1876: 109.

Type horizon. Early Langhian, middle Miocene.

Type locality. Originally given as “Sinj” and later specified as “Župića potok” in Brusina (1884b), Croatia.

Types. Milan et al. (1974: 87) stated that Brusina (1897) had indicated one of the specimens illustrated by him (pl. 5, figs 1–2) as type. However, it is uncertain whether the specimen actually derives from the original type series. The specimen is stored in the Croatian Natural History Museum, Zagreb, coll. no. 2980-626/1.

Microcolpia canaliculata Bourguignat, 1884

Original source. Bourguignat 1884: 62.

Type locality. “Le Danube à Ibraïla” [Danube river at Brăila], Romania.

† *Melanopsis capelliniana* Pallary, 1920

Original source. Pallary 1920b: 115.

Type horizon. Messinian, late Miocene.

Type locality. “Colognole e al casino Cubbe; [...] presso il casino Sant’Andrea sotto Colognole” (Capellini 1880: 397) [at Colognole and at Casa Cubbe; near Casa Sant’Andrea below Colognole], Italy.

Remarks. Introduced for “*M. dufourii* var. a” in Capellini, 1880. Pallary used multiple original spellings (“*capelliniana*”, “*capelliana*”) but according to 32.5.1 the name must be *capelliniana*. Wenz (1929: 2785) considered Capellini’s material synonymous with *M. narzolina* Manzoni, 1870.

† *Melanopsis harpula* var. *capreniensis* Fontannes, 1887

Original source. Fontannes 1887: 335.

Type horizon. Early Cernikian, early Pliocene.

Type locality. “Căpreni, val. Amaradii (Jud. Gorjiu)” [Căpreni], Romania.

† *Melanopsis (Martinia) vindobonensis* var. *capuliformis* Handmann, 1887

Original source. Handmann 1887: 29, pl. 6, figs 5–6.

Type horizon. Pannonian, zone B–D, late Miocene.

Type locality. “Leobersdorf”, Austria.

† *Melanopsis capulus* Handmann, 1882

Original source. Handmann 1882: 554.

Type horizon. Pannonian, zone D, late Miocene.

Type locality. “Kottingbrunn [...] Ziegelei a”, Austria.

Remarks. The species epithet is a noun in apposition and needs not to agree in gender with the generic name (Art. 31.2.1). Wenz (1929: 2718) considered this taxon as a junior synonym of *Melanopsis fossilis* (Gmelin, 1791).

† *Melanopsis caputinensis* Stefanescu, 1897

Original source. Stefanescu 1897: 311, pl. 8, figs 8–10.

Type horizon. Pliocene.

Type locality. “Au ravin de Tura, à Capatzineni; [...] aussi à Salatrucu-Mare” [in the ravine of Tura in Căpățânenii; also at Sălătrucu], Romania.

† *Melanopsis carasiensis* Jekelius, 1944

Original source. Jekelius 1944: 131, pl. 49, figs 7–10.

Type horizon. Early Pannonian, late Miocene.

Type locality. “Turislav-Tal bei Soceni” [Turislav valley near Soceni], Romania.

† *Melanopsis (Melanoptychia) cari* Pavlović, 1927

Original source. Pavlović 1927: 61, pl. 7, figs 3–4.

Type horizon. Pannonian, zone D–E, late Miocene.

Type locality. “Из Панахе” [from the village Ramača (near Ripanj)], Serbia.

Types. The illustrated syntype is stored in the Natural History Museum, Belgrade, coll. no. 796 (Milošević 1962: 23).

† *Melanopsis carinata* Sowerby, 1826

Original source. Sowerby 1826–1829: 41, pl. 523, fig. 1.

Type horizon. Eocene.

Type locality. “In a well near Newport, Isle of Wight; [...] from Hampstead Cliff to Cowes, and [...] on the opposite Cliffs of Hampshire”, United Kingdom.

Melanopsis carinata Gassies, 1861 [invalid]

Original source. Gassies 1861: 289, pl. 7, fig. 13.

Type locality. “Dans le Diahot, à Balade; à Jengen, à Kanala, dans les marais et les petits ruisseaux” [in the Diahot river at Balade; at Hienghène, at Canala, in swamps and small streams], New Caledonia.

Remarks. Junior homonym of *Melanopsis carinata* Sowerby, 1826. Pallary (1916: 82) introduced *M. ducosi* as replacement name.

Melanopsis mingrelica var. *carinata* Issel, 1865 [invalid]

Original source. Issel 1865 [page unknown; original work not seen; described on p. 400 of the re-published version from 1866].

Type locality. “Nel lago di Paleaston presso Poti” [Paliastomi Lake near Poti], Georgia.

Remarks. Junior homonym of *Melanopsis carinata* Sowerby, 1826.

Melanopsis dufourii var. *carinata* Gentiluomo, 1868 [invalid]

Original source. Gentiluomo 1868: 97, pl. 6, figs 10–11.

Type locality. “Lago dell’Accesa, Toscana”, Italy.

Remarks. Junior homonym of *Melanopsis carinata* Sowerby, 1826.

† *Melanopsis hazayi* var. *carinata* Brusina, 1903 [invalid]

Original source. Brusina 1903: 113.

Type horizon. Late Pleistocene–early Holocene.

Type locality. “Bischofsbad” [Püspökfürdő, Băile 1 Mai, Lake Pețea], Romania.

Remarks. Junior homonym of *Melanopsis carinata* Sowerby, 1826. Neubauer et al. (2014d: 125) considered this taxon as a junior synonym of *Microcolpia parreyssii sikorai* (Brusina, 1903).

† *Melanopsis sikorai* var. *carinata* Kormos, 1903 [invalid]

Original source. Kormos 1903: 456, 501, pl. 13, fig. 1.

Type horizon. Late Pleistocene–early Holocene.

Type locality. “Püspökfürdő” [Băile 1 Mai, Lake Pețea], Romania.

Remarks. Obviously unaware of the fact that variety names are available in nomenclature as species-group names, Kormos (1903) stated that, if raised to species, he suggests “*M. mucronifera*” as name for the taxon. Both names were simultaneously published and are junior objective synonyms. Since *carinata* sensu Kormos is a junior homonym of *Melanopsis carinata* Sowerby, 1826, *M. mucronifera* is the valid name of the taxon.

† *Melanopsis staubi* var. *carinata* Brusina, 1903 [invalid]

Original source. Brusina 1903: 115.

Type horizon. Late Pleistocene–early Holocene.

Type locality. “Bischofsbad” [Püspökfürdő, Băile 1 Mai, Lake Pețea], Romania.

Remarks. Junior objective synonym of *M. staubi*: Brusina (1903) indicated it as the typical form of the species. Junior homonym of *M. carinata* Sowerby, 1826. Neubauer et al. (2014d: 125) considered this taxon as a junior synonym of *Microcolpia parreyssii sikorai* (Brusina, 1903).

Melanopsis doriae var. *carinata* Biggs, 1937 [invalid]

Original source. Biggs 1937: 248.

Type locality. “From Jelalabad” [Jalalabad], Iran.

Remarks. Junior homonym of *Melanopsis carinata* Sowerby, 1826.

† *Melanopsis impressa* var. *carinatissima* Sacco, 1889

Original source. Sacco 1889: 65, pl. 2, figs 24–25.

Type horizon. Tortonian–early Messinian, late Miocene.

Type locality. “Delle colline tortonesi presso S. Agata” [from the Torino hills near Sant’Agata Fossili], Italy.

† *Melanopsis aetolica* var. *carinatocostata* Oppenheim, 1891

Original source. Oppenheim 1891: 468, pl. 27, figs 1–2.

Type horizon. Gelasian, early Pleistocene.

Type locality. “Stamna”, Greece.

Remarks. Not available from Oppenheim (1890b), where he gave it as “mutation”, which is not ruled by the provisions of the Code. In 1891 he introduced the name as “*carinato-costata*”.

Melanopsis praerosa [sic] var. *carinifera* Paetel, 1888

Original source. Paetel 1888: 403.

Type locality. “Kerman” (Nevill 1884: 209), Iran.

Remarks. Originally introduced as infrasubspecific taxon (“subvariety”) by Nevill (1884), but made available by Paetel (1888) who treated it as variety (Art. 45.5.1). Paetel clearly referred to the description of Nevill.

Murex cariosus Linnaeus, 1767

Original source. Linnaeus 1767: 1220.

Type locality. “In Aqaeductu ad Sevillam” [in an aqueduct near Sevilla], Spain.

Remarks. Today combined as *Melanopsis cariosa*, the species is one of the few stable taxa in melanopsid taxonomy (e.g., Glaubrecht 1996).

† *Melanoptychia carusi* Brusina, 1902

Original source. Brusina 1902: pl. 30, figs 6–7.

Type horizon. Langhian, middle Miocene.

Type locality. “Džepe” [Džepi], Bosnia and Herzegovina.

Types. The original type material of Brusina (1902) is lost. Neubauer et al. (2016c: 275–276) defined a neotype from the type locality. The specimen is stored in the Geological Survey Austria, Vienna, coll. no. 2014/010/0002.

† *Lyrcaea* [sic] *caryota* Brusina, 1902

Original source. Brusina 1902: pl. 5, figs 21–25.

Type horizon. Transdanubian, Pannonian, late Miocene.

Type locality. “Kenese” [Balatonkenese], Hungary.

Types. The illustrated syntypes are stored in the Croatian Natural History Museum, Zagreb, coll. no. 2485-131/1-5 (Milan et al. 1974: 83).

“*Melanopsis castanea*” mentioned in Férussac (1814: 54) [unavailable]

Locality. Not indicated.

Remarks. *Nomen nudum*. Férussac (1823: 149) listed it in synonymy of *M. buccinoidea* var. α [= *M. buccinoidea* (Olivier, 1801)].

***Melanella castanea* Bourguignat, 1884**

Original source. Bourguignat 1884: 27.

Type locality. “Rivière à Ostaria et à Ogulin (Croatie)” [river at Oštarije and at Ogulin], Croatia.

Remarks. Note that Bourguignat denoted the authority as “Bourguignat, 1879”.

† *Melanopsis castrensis* Noulet, 1854

Original source. Noulet 1854: 50.

Type horizon. Bartonian, Eocene.

Type locality. “À Labruguière; [...] à Augmontel (Tarn)”, France.

***Melanella coronata* var. *catoleia* Bourguignat, 1884**

Original source. Bourguignat 1884: 12.

Type locality. “De la Savina” [from the Savinja river], Slovenia.

***Melanopsis ceardi* Pallary, 1928**

Original source. Pallary 1928a: 268, pl. 6, figs 4–5.

Type locality. “Ouakda, à 4 kil. N.-E. de Colomb-Béchar” [Ouakda, 4 km north-east of Bechar], Algeria.

† *Melanopsis cepula* Guppy, 1866

Original source. Guppy 1866: 580, pl. 26, fig. 14.

Type horizon. Late Miocene.

Type locality. “Cumana”, Venezuela.

Remarks. Given as “*capula*” on p. 580, but as “*cepula*” in the plate captions. Guppy (1867) acted as First Reviser and fixed the correct original spelling as “*cepula*” (Art. 24.2.3, 32.2.1). Guppy (1867) did not consider the species a melanopsid anymore and introduced the new genus *Crepitacella* (Rissoinidae) with *M. cepula* as type species.

† *Melanopsis cerithiformis* Watelet, 1851

Original source. Watelet 1851: 121, pl. 1, figs 1–2.

Type horizon. Cuisian, late Ypresian, Eocene.

Type locality. “Mercin” [Mercin-et-Vaux], France.

Remarks. Cossmann (1888: 279) classified the species in the genus *Faunus* Montfort, 1810 (Pachychilidae). Wenz (1929: 2637) apparently agreed with that classification but listed it a second time (p. 2563) in synonymy of *Semisinus* [= *Hemisinus*] *resectus* (Deshayes, 1834).

Melanopsis cerithiopsis Bourguignat, 1884

Original source. Bourguignat 1884: 130.

Type locality. “Plaine du Bahr-el-Houlé (haut Jourdain) dans l’Aïn-el-Mellaha” [in the plains of the Hula valley (upper Jordan), in Aïn Mallahah], Israel.

Remarks. Given as “*cerithiopsis*” in heading of description, but as “*cerithiopsis*” throughout the rest of the work. Heller et al. (2005: 248) considered the species as a junior synonym of *Melanopsis saulcyi* Bourguignat, 1853.

Melanopsis cerullii Pallary, 1920

Original source. Pallary 1920b: 110.

Type locality. “M. Mario: Farnesina” (Cerulli-Irelli 1914: 186), Italy.

Remarks. Replacement name for *M. transiens* Cerulli-Irelli, 1914, non Blanckenhorn, 1897. Girotti (1972: 232) considered *M. transiens* Cerulli-Irelli, 1914 a junior synonym of “*Melanopsis affinis* Férussac”, which is not an available name.

Melanopsis cesari Pallary, 1920

Original source. Pallary 1920a: 30.

Type locality. “Beni Abbès” [prov. Béchar], Algeria.

† *Melanopsis clavigera* f. *cecticillus* Brusina, 1897

Original source. Brusina 1897: 7, pl. 5, fig. 20.

Type horizon. Cernikian, Pliocene.

Type locality. “Kozarica” [Kozarice], Croatia.

Types. Milan et al. (1974: 88) indicated a holotype, but it is uncertain whether the specimen was the only one Brusina had at hand (holotype by monotypy, Art. 73.1.2). The specimen is stored in the Croatian Natural History Museum, Zagreb, coll. no. 2991-637.

Remarks. Mandić et al. (2015: 194) considered this taxon as a junior synonym of *Melanopsis clavigera* Neumayr in Neumayr & Paul, 1875.

***Melanopsis chantrei* Locard, 1883**

Original source. Locard 1883a: 268, pl. 23, figs 44–49.

Type locality. “Lac d’Antioche” [Lake Anuk (also as Amik)], Turkey.

***Melanopsis charpentieri* Brot, 1879**

Original source. Brot 1874–1879: 430, pl. 46, fig. 8.

Type locality. “Schiraz”, Iran.

Remarks. Brot (1879) attributed the name to Parreyss, based on his “in schedis” determination.

***Melanopsis chehirensis* Pallary, 1939**

Original source. Pallary 1939: 94, pl. 6, figs 51–58, 75.

Type locality. “Dans le source de Yeni Chehir, [...] entre Antioche et Alep, à l’intersection de la route d’Alexandrette” [at the source of Yenişehir, between Antakya and Aleppo, at the intersection of the road from İskenderun], Turkey.

***Melanopsis (Mesopotamia) cheragragensis* Pallary, 1939**

Original source. Pallary 1939: 104, pl. 5, figs 8–9, 11.

Type locality. “Cheragrag, en Mésopotamie, entre Rakka, au Sud et Tell Abiad au Nord, sur la rive gauche du Karamouk Sou, affluent de la rive droite du Nahr Bâhlik” [Sharakrak in Mesopotamia, between Ar Raqqa in the south and Tall Abyad in the north, on the left bank of the Qarah Mûkh, tributary of the right side of the Nahr al Balikh], Syria.

Remarks. Heller et al. (2005: 254) considered the species as a junior synonym of *Melanopsis infracincta* Martens, 1874.

***Melanopsis buccinoidea* var. *chlorotica* Pallary, 1913**

Original source. Pallary 1913: 364.

Type locality. “L’Abreuvoir de Dar-Beida” [water trough of Dar Beïda], southern Morocco.

***Melanopsis chobauti* Nicolas, 1898**

Original source. Nicolas 1898: 530.

Type locality. “Aux environs de Biskra, à Ain-Oumach [...] dans une source d’eau chaude” [around Biskra, at Ain Oumache; in a hot spring], Algeria.

Remarks. Pallary (1912: 14) considered the taxon as a junior synonym of *Melanopsis saharica* Bourguignat, 1864.

† *Melanopsis choctavensis* Aldrich, 1886

Original source. Aldrich 1886: 35, pl. 3, fig. 8.

Type horizon. Early Eocene.

Type locality. “Hatchetigbee, Butler, Choctaw County, Alabama”, United States.

Types. Palmer and Brann (1966: 546) stated the the figured syntype was missing. A remaining syntype is stored in the collection of the National Museum of Natural History, Smithsonian Institution, Washington, DC, coll. no. 638787.

Remarks. Classified within the marine genus *Bulliopsis* Conrad, 1862 (Nassariidae) by Harris (1899: 58) and Allmon (1990: 53).

† *Melanopsis cikosi* Brusina, 1902

Original source. Brusina 1902: pl. 6, fig. 5.

Type horizon. Early Cernikian, early Pliocene.

Type locality. “Čerević”, Serbia.

Types. Milan et al. (1974: 88) indicated a holotype, but it is uncertain whether the specimen was the only one Brusina had at hand (holotype by monotypy, Art. 73.1.2). The specimen is stored in the Croatian Natural History Museum, Zagreb, coll. no. 2504-150.

† *Melanopsis cincta* Neumayr, 1880

Original source. Neumayr 1880b: 290, pl. 1, fig. 10.

Type horizon. Plio-Pleistocene.

Type locality. “Zwischen Pyllé und Antimachia” [between Pýli and Antimácheia, Kos Island], Greece.

Remarks. Willmann (1981: 179) considered this taxon as a junior synonym of *Melanopsis delessei* Tournouër, 1875.

† *Melanopsis citharella* Merian, 1849

Original source. Merian 1849: 31.

Type horizon. Burdigalian, early Miocene.

Type locality. “Nahe am Gipfel des Plateaus des Randens” [near the top of the plateau of Mt. Randen], Switzerland.

† *Melanopsis clava* Sandberger, 1872

Original source. Sandberger 1870–1875: pl. 25, fig. 31.

Type horizon. Badenian, middle Miocene.

Type locality. “Grund [...], Vöslau” (Hörnes 1856: 597), Austria.

Remarks. Plate 25 of Sandberger’s monograph was issued in 1872, while the description on pp. 512, 521 appeared in 1875 (Woodward 1906). Introduced for *M. aquensis* sensu Hörnes, 1856, non Grateloup, 1838.

† *Melanopsis clavigera* Neumayr in Neumayr & Paul, 1875

Original source. Neumayr and Paul 1875: 41, pl. 7, figs 13–14.

Type horizon. Cernikian, Pliocene.

Type locality. “Ciglenik; [...] Graben zwischen Podwin und der Čapla; [...] Čapla; [...] An der Strasse von Sibin nach Gromačnik; [...] Gromačnik” [Ciglenik; Čaplja trench near Slavonski Brod; Čaplja; at the road from Sibin to Gromačnik; Gromačnik], Croatia.

† *Melanopsis clementina* Michelin, 1833

Original source. Michelin 1833: pl. 29.

Type horizon. Cretaceous?

Type locality. “À Gérodot, département de l’Aube”, France.

Remarks. Probably not a Melanopsidae.

† *Melanopsis (Martinia) martiniana* var. *coaequata* Handmann, 1887

Original source. Handmann 1887: 25, pl. 4, figs 8–9.

Type horizon. Pannonian, zone B–D, late Miocene.

Type locality. “Leobersdorf”, Austria.

Types. A lectotype was designated by Fischer (1996b: 20). It is stored in the collection of the Geological Survey Austria, Vienna; no number indicated.

***Melanella codiella* Servain, 1884**

Original source. Bourguignat 1884: 27.

Type locality. “La Migliaska à Serajewo (Bosnie); Ostaria (Croatie)” [in the Miljacka river at Sarajevo (Bosnia and Herzegovina); Oštarije (Croatia)].

Remarks. Appeared first as a nomen nudum in Servain (1884).

† *Melanopsis cognata* Brusina, 1878

Original source. Brusina 1878: 349.

Type horizon. Late Portaferrian, late Miocene–early Pliocene.

Type locality. “Karlowitz [...] Görgetek in Syrmien” (Neumayr and Paul 1875: 49) [Karlovcı; Grgeteg], Serbia.

Types. Milan et al. (1974: 88) indicated a holotype, but it is uncertain whether the specimen was the only one Brusina had at hand (holotype by monotypy, Art. 73.1.2). The specimen is stored in the Croatian Natural History Museum, Zagreb, coll. no. 3001-647.

Remarks. Introduced for “*M. cf. visianiana*” in Neumayr and Paul, 1875, non Brusina, 1874.

***Melanopsis guiraoi* var. *communis* Pallary, 1924**

Original source. Pallary 1924: 251, pl. 25, fig. 20.

Type locality. “Dans les cours d’eau de la province de Murcie” [in rivers of the province Murcia], Spain.

† *Melanopsis neumayri* var. *compacta* Fontannes, 1880 [invalid]

Original source. Fontannes 1879–1882: 174.

Type horizon. Mammal zone MN 11, late Miocene.

Type locality. “Les marnes à *Potamides Basteroti* de Visan (Vaucluse)” [in the marls with *P. basteroti* in Visan, Dép. Vaucluse], France.

Remarks. Wenz (1929: 2793) considered the variety as a junior synonym *M. neumayri* Tournouër, 1874.

***Melanopsis compacta* Pallary, 1920 [invalid]**

Original source. Pallary 1920a: 31.

Type locality. “Aït Taleb sur le Sefrou près d’el Menzel, avant l’oued Sebou” [Douar Ait Taleb at Sefrou near El Menzel, before the Sebou river], Morocco.

Remarks. Junior homonym of *Melanopsis compacta* Fontannes, 1880.

† *Lyrcaea narzolina* var. *compressoides* Sacco, 1895

Original source. Sacco 1895: 14, pl. 1, fig. 26.

Type horizon. Tortonian–early Messinian, late Miocene.

Type locality. “S. Agata fossili” [Sant’Agata Fossili], Italy.

† *Melanopsis conemenosiana* Oppenheim, 1891

Original source. Oppenheim 1891: 469, pl. 27, figs 7–8.

Type horizon. Late Pliocene–early Pleistocene.

Type locality. “Preveza”, Greece.

Remarks. Appeared first as a nomen nudum (as “*M. Conemenosi* Bttg. in litt.”) in Oppenheim (1890b).

† *Melanopsis confusa* Strausz, 1941

Original source. Strausz 1941: 143.

Type horizon. Transdanubian, Pannonian, late Miocene.

Type locality. “Radmanest” (Fuchs 1870: 353) [Rădmănești], Romania.

Remarks. Replacement name for *M. fuchsi* Brusina, 1884, non Handmann, 1882 and *M. hungarica* Pallary, 1916, non Kormos, 1904, which were in turn introduced for *M. costata* sensu Fuchs, 1870, non Olivier, 1804 (see also Strausz 1942; Neubauer et al. 2014a).

Melanopsis praerosa var. *conica* Bronn, 1848 [invalid]

Original source. Bronn 1848: 718.

Type horizon. Not stated, probably fossil.

Type locality. Not indicated.

Remarks. Junior objective synonym of *Melanopsis antediluviana* (Poirer, 1801), which Bronn (1848) gave in synonymy.

† *Pseudofagotia conica* Anistratenko, 1993

Original source. Anistratenko 1993: 74, textfig. 2.

Type horizon. Duab Beds, middle to late Kimmerian, Pliocene.

Type locality. “Окр. с. Мокви, Очамчирский р-н” [near the village Mok’vi, Ochamchirskiy rayon], Georgia.

Types. Schmalhausen Institute of Zoology of National Academy of Sciences of Ukraine, Kiev; no number indicated.

† *Pseudofagotia coniformis* Anistratenko, 1993

Original source. Anistratenko 1993: 74, textfig. 2.

Type horizon. Duab Beds, middle to late Kimmerian, Pliocene.

Type locality. “Окр. с. Мокѳу, Очамчирский р-н” [near the village Mok’vi, Ochamchirskiy rayon], Georgia.

Types. Schmalhausen Institute of Zoology of National Academy of Sciences of Ukraine, Kiev; no number indicated.

† *Melanopsis coniungens* Sacco, 1886

Original source. Sacco 1886: 450, pl. 1, fig. 9.

Type horizon. Late Burdigalian, early Miocene.

Type locality. “Collina di Torino” [Torino hills], Italy.

Remarks. The name “*conjungens*” as mentioned in Sacco (1889: 68) is an incorrect subsequent spelling.

Melanopsis jordanica var. *conoidea* Bourguignat, 1884

Original source. Bourguignat 1884: 142.

Type locality. “Lac de Tibériade” [Sea of Galilee], Israel.

† *Melanopsis conoidea* Pană, 2003

Original source. Pană 2003: 319, pl. 9, figs 6–13.

Type horizon. Parscovian–Pelendavian, Pliocene.

Type locality. “Forage Mihăița, profundeur 109 m” [borehole Mihăița, at a depth of 109 m], Romania.

Types. Laboratory of Paleontology, Bucharest, coll. no. 651.

Remarks. Pană denoted the authorship as “Pană, 1989”. Originally the gender was indicated as masculine (“*conoideus*”), but *Melanopsis* is feminine, which is why the name must be “*conoidea*”.

† *Melanopsis (Martinia) vindobonensis* var. *consimilis* Handmann, 1887

Original source. Handmann 1887: 29, pl. 6, figs 7–10.

Type horizon. Pannonian, zone B–D, late Miocene.

Type locality. “Leobersdorf”, Austria.

† *Lyrcaea megacantha* f. *conspicua* Brusina, 1897

Original source. Brusina 1897: 13, pl. 7, figs 3–4.

Type horizon. Pannonian, zone D–E, late Miocene.

Type locality. “Begaljica”, Serbia.

Types. The syntypes are stored in the Croatian Natural History Museum, Zagreb; no number indicated (Milan et al. 1974: 84).

Remarks. Wenz (1929: 2682) considered this taxon as a junior synonym of *Melanopsis bouei megacantha* Handmann, 1887.

† *Melanopsis conspicua* Pallary, 1916 [invalid]

Original source. Pallary 1916: 80.

Type horizon. Chattian, Oligocene.

Type locality. “Dax. St-Geours, Abesse” (Grateloup 1840: captions of the plate “Mollusques terrestres et fluviatiles fossiles de Dax”), France.

Remarks. Introduced for *M. costata* sensu Grateloup, 1840, for which d’Orbigny (1852) had already introduced *M. nereis* as new name. Thus, *M. conspicua* is a junior objective synonym of *M. nereis*. Moreover, it is a junior secondary homonym of *Lyrcaea conspicua* Brusina, 1897.

† *Melanopsis constricta* Brusina, 1878

Original source. Brusina 1878: 348.

Type horizon. Cernikian, Pliocene.

Type locality. “Kozarica” [Kozarice], Croatia.

Types. Milan et al. (1974: 88) stated that Brusina (1897) had indicated one of the specimens illustrated by him (pl. 5, fig. 21) as type. However, it is uncertain whether the specimen actually derives from the original type series. The specimen is stored in the Croatian Natural History Museum, Zagreb, coll. no. 2992-638/1.

† *Melanopsis (Martinia) martiniana* var. *constricta* Handmann, 1887 [invalid]

Original source. Handmann 1887: 26, pl. 5, figs 1–2.

Type horizon. Pannonian, zone B–D, late Miocene.

Type locality. “Leobersdorf”, Austria.

Remarks. Junior homonym of *Melanopsis constricta* Brusina, 1878. Fischer (1996a) introduced *M. handmanniana* as replacement name.

† *Melanopsis vindobonensis* var. *contecta* Handmann, 1882

Original source. Handmann 1882: 554.

Type horizon. Pannonian, zone D, late Miocene.

Type locality. “Kottingbrunn”, Austria.

Remarks. Wenz (1929: 2848) considered the taxon as a junior synonym of *Melanopsis vindobonensis* Fuchs, 1870. Note that Wenz gave the name as “*costata*”, which is an incorrect subsequent spelling.

† *Melanopsis (Martinia) vindobonensis* var. *contigua* Handmann, 1887

Original source. Handmann 1887: 29, pl. 6, figs 11–12.

Type horizon. Pannonian, zone B–D, late Miocene.

Type locality. “Leobersdorf”, Austria.

† *Melanopsis hazayi* var. *contracta* Brusina, 1903 [invalid]

Original source. Brusina 1903: 112.

Type horizon. Late Pleistocene–early Holocene.

Type locality. “Bischofsbad” [Püspökfürdő, Băile 1 Mai, Lake Peșea], Romania.

Remarks. Junior objective synonym of *M. hazayi*: Brusina (1903) indicated it as the typical form of the species. Neubauer et al. (2014d: 125) considered this taxon as a junior synonym of *Microcolpia parreyssii sikorai* (Brusina, 1903).

† *Melanopsis convexa* Doncieux, 1908

Original source. Doncieux 1908: 202, pl. 11, fig. 10.

Type horizon. Lutetian, Eocene.

Type locality. “Au Nord d’Albas”, France.

Melanopsis cookiana Pallary, 1916 [invalid]

Original source. Pallary 1916: 82.

Type locality. “Prope Kanala [...]; insula Ouen” (Gassies 1870: 148) [near Canala; Île Ouen], New Caledonia.

Remarks. Replacement name for the junior homonym *M. fusiformis* Gassies, 1870, non Sowerby, 1822, for which Gassies (1880) had already introduced *M. rossiteri* as replacement name. Thus, *M. cookiana* is a junior objective synonym of *M. rossiteri*.

† *Melanopsis corici* Neubauer, Mandić, Harzhauser & Hrvatović, 2013

Original source. Neubauer et al. 2013: 134, figs 4E–H.

Type horizon. Langhian, middle Miocene.

Type locality. “Fatelj section”, Bosnia and Herzegovina.

Types. Geological-Paleontological Department, Natural History Museum Vienna, Austria, coll. no. 2011/0138/0100.

Melanopsis cornea Pfeiffer, 1828

Original source. Pfeiffer 1828: 50, pl. 8, figs 22–23.

Type locality. “In der Donau bei Pesth” [in the Danube river near Budapest], Hungary.

Melania cornea Reeve, 1860

Original source. Reeve 1860: Section *Melania*, pl. 34, fig. 233.

Type locality. “Dalmatia” [no locality indicated], Croatia.

Remarks. Based on a manuscript or “in schedis” name from Küster in the museum of Von dem Busch (see Reeve 1860).

Melania coronata Reeve, 1860 [invalid]

Original source. Reeve 1860: Section *Melania*, pl. 34, fig. 228.

Type locality. “Römerbad in Steiermark” [Rimske Toplice], Slovenia (after Clessin 1890, no locality given in Reeve 1860).

Remarks. Based on a manuscript name by Küster and introduced in synonymy of *Melania hollandri* [sic] Pfeiffer, 1828 (see also Brot 1874 and Clessin 1890). It was made available by Bourguignat (1884), who treated it as valid name (although he denoted the authority with “Bourguignat, 1877”; see Note 2). Junior homonym of *Melania coronata* Von dem Busch in Philippi, 1845 (Bengal).

† *Melanopsis (Lyrcea) coronata* Brusina, 1878

Original source. Brusina 1878: 348.

Type horizon. Cernikian, Pliocene.

Type locality. “Repusnica, Slobodnica”, Croatia.

Types. The syntype (?) illustrated in Brusina (1897: 12, pl. 5, figs 15–16) is stored in the Croatian Natural History Museum, Zagreb, coll. no. 2987-633 (Milan et al. 1974: 84).

***Melanopsis tricarinata coronata* Ahuir Galindo, 2015 [invalid]**

Original source. Ahuir Galindo 2015: 22, unnumbered figure.

Type locality. “In brooks and ‘gorgos’ from Anna, Valencia”, Spain.

Types. Museo Malacologico di Cupra Marittima, Italy; no number indicated.

Remarks. Junior homonym of *Melanopsis coronata* Brusina, 1878.

***Melanopsis obediensis* var. *coroniformis* Picard, 1934**

Original source. Picard 1934: 121, pl. 7, figs 45–52.

Type locality. “Jarmukmündung” [Yarmouk river mouth], Jordan/Israel.

† *Melanopsis rumana* var. *correcta* Stefanescu, 1896

Original source. Stefanescu 1896: 128, pl. 11, figs 4–6.

Type horizon. Pliocene.

Type locality. “À Mosculesti, dans la vallée de Gilortu, à Gura-Motrului et à Breasta, dans la vallée de Jiu” [at Musculești in the valley of the river Gilortu, at Gura Motrului and at Breasta in the valley of the river Jiu], Romania.

† *Melanopsis corrugata* Schütt in Schütt & Ortal, 1993

Original source. Schütt and Ortal 1993: 91, pl. 2, figs 15–16.

Type horizon. Pleistocene, Riss glacial epoch.

Type locality. “Galilee, southern Hula basin, about 500 m north of the new bridge over the river Jordan at Benot Ya’Aqov”, Israel.

Types. Paleontology Collection of the Hebrew University of Jerusalem; no number indicated.

† *Melanopsis cosiana* Pallary, 1925 [invalid]

Original source. Pallary 1925: 257.

Type horizon. Plio-Pleistocene.

Type locality. “Zwischen Pylle und Antimachia” (Neumayr 1880b: 295) [between Pýli and Antimácheia, Kos Island], Greece.

Remarks. Invalid replacement name for *Melanopsis broti* Neumayr, 1880, “non Gassies, 1874”, but Gassies had actually introduced his species as “*M. brotiana*”.

† *Melanopsis cosmanni* [sic] Pallary, 1916 [invalid]

Original source. Pallary 1916: 80.

Type horizon. Cernikian, Pliocene.

Type locality. “De Slavonie” (Cossmann 1909: 176), Croatia.

Remarks. Introduced for *M. costata* sensu Cossmann, 1909, non Olivier, 1804. The name was corrected to “*cosmanni*” by Neubauer et al. (2014a: 461), which is a justified emendation according to Art. 32.5.1 and 33.2.2. Wenz (1929) considered all records of “*M. costata*” from the Pliocene of Slavonia to represent the same species and synonymized them with *M. abbreviata cosmanni*. Obviously, he was unaware that *M. croatica* Brusina, 1884 is the first available name for them (see also discussion of *M. pseudocostata* Oppenheim, 1890).

† *Melanopsis cosmanni* Pallary, 1916

Original source. Pallary 1916: 80 (as “*cosmanni*”).

Type horizon. Cernikian, Pliocene.

Type locality. “De Slavonie” (Cossmann 1909: 176), Croatia.

Remarks. Justified emendation of *M. cosmanni* Pallary, 1916 by Neubauer et al. (2014a: 461).

Melanopsis cossoni Bourguignat, 1884

Original source. Bourguignat 1884: 111.

Type locality. “Eaux thermales d’Ouargla et près du chott Tiraoun dans le sud de la province de Constantine et de la Tunisie” [in thermal waters at Ouargla (Algeria) and near chott Tiraoun (Tunisia)], Algeria.

Melania costata Olivier, 1804

Original source. Olivier 1804: 294 (footnote), pl. 31, fig. 3.

Type locality. “De Orontes [Gesser-Chourl]” [in the Orontes river, at Jisr Ash-Shughur], Syria.

Remarks. Type species of the genus *Melanopsis* Férussac in Férussac & Férussac, 1807.

† *Melanopsis costata* Ludwig, 1865 [invalid]

Original source. Ludwig 1865: 71, pl. 21, figs 7–7b.

Type horizon. Early Rupelian, Oligocene.

Type locality. “Grossalmerode”, Germany.

Remarks. Junior secondary homonym of *M. costata* (Olivier, 1804). Speyer (1870: 97) introduced *M. ludwigi* as replacement name.

† *Melanopsis bergeroni* var. *costata* Botez, 1914 [invalid]

Original source. Botez 1914: 240.

Type horizon. *Viviparus stricturatus* Zone, Cernikian, Pliocene.

Type locality. “Moreni”, Romania.

Remarks. Junior secondary homonym of *Melanopsis costata* (Olivier, 1804).

† *Melanopsis lanzae costata* Olujić, 1999 [invalid]

Original source. Olujić 1999: 21, 49, pl. 2, figs 19–24, pl. 3, figs 26–28.

Type horizon. Langhian, middle Miocene.

Type locality. It is unclear from the original work in which of the studied localities/sections along the valleys of the Sutina, Batarelov and Vojskava rivers (4 km W of Sinj) the taxon occurred and in which not, Croatia.

Remarks. Junior secondary homonym of *Melanopsis costata* (Olivier, 1804). Neubauer et al. (2011) considered it as a junior synonym of *M. lanzaeana*.

† *Melanoptychia lyrata costata* Olujić, 1999 [invalid]

Original source. Olujić 1999: 20, 48, pl. 1, fig. 5.

Type horizon. Langhian, middle Miocene.

Type locality. It is unclear from the original work in which of the studied localities/sections along the valleys of the Sutina, Batarelov and Vojskava rivers (4 km W of Sinj) the taxon occurred and in which not, Croatia.

Remarks. Junior secondary homonym of *Melanopsis costata* (Olivier, 1804). Considered as a junior synonym of *Melanopsis lyrata* Neumayr, 1869 by Neubauer et al. (2011: 207).

“*Mellanella hollandri* [sic] var. *costata* [sic] Kucik” mentioned in Brusina (1867: 85) [unavailable]

Locality. “U Savi i u potoku kod Susjeda, u Žirovcu, u Maksimiru kod Zagreba, u potoku Toplici kod Oroslavja i u Kutinji blizu Jastrebarskoga” [in the river Sava and a creek at Podsused in Zagreb, in Žirovac, in Maksimir at Zagreb, in the river Toplica at Oroslavje and in Kutinja (?) near Jastrebarsko], Croatia.

Remarks. Nomen nudum, based on an “in schedis” name from the collection of Kucik (also read as “Kutschig”). The name was probably a typesetting mistake for “*costata*”.

† *Melanopsis (Canthidomus) costatiformis* Papp, 1953

Original source. Papp 1953a: 108, pl. 23, figs 1–8.

Type horizon. Gelasian, early Pleistocene.

Type locality. “Aghios Georgios” [Agios Georgios, Elis], Greece.

Types. Museum of Palaeontology and Geology of the University of Athens; no number indicated.

Melanopsis costellata Férussac, 1823 [invalid]

Original source. Férussac 1823: 157.

Type locality. “Dans l’aqueduc de Séville [...] et dans les ruisseaux des environs. [...] Dans les lacs et les rivières du royaume de Maroc” [in the aqueduct of Sevilla and in the rivers of its surroundings (Spain); in the lakes and rivers of Morocco].

Remarks. Junior objective synonym of *M. cariosa* (Linnaeus, 1767), which Férussac listed in synonymy.

† *Melanopsis costellata* Douvillé, 1904 [invalid]

Original source. Douvillé 1904: 327, pl. 46, figs 7–11.

Type horizon. Maastrichtian, Cretaceous.

Type locality. “Du versant oriental du Kouh Mapeul” [eastern slope of mount Kuh-e Mapel, c. 60 km WNW of Khorramābād], Iran.

Remarks. Junior homonym of *Melanopsis costellata* Férussac, 1823. Pallary (1916: 83) introduced *M. douvillei* as replacement name.

† *Pseudofagotia costifera* Anistratenko, 1993

Original source. Anistratenko 1993: 74, textfig. 2.

Type horizon. Duab Beds, middle to late Kimmerian, Pliocene.

Type locality. “Okp. c. Mokvu, Очамчирский р-н” [near the village Mok’vi, Ochamchirskiy rayon], Georgia.

Types. Schmalhausen Institute of Zoology of National Academy of Sciences of Ukraine, Kiev; no number indicated.

***Melania holandri* var. *costulata* Schmidt, 1847**

Original source. Schmidt 1847: 25.

Type locality. “Aus einem Mühlbache bei Klinze ob Schischka” [from a mill creek at Glinica near Šiška], Slovenia.

† *Melanopsis inconstans* var. *costulata* Brusina, 1874 [invalid]

Original source. Brusina 1874: 39.

Type horizon. Langhian, middle Miocene.

Type locality. “Miočić”, Croatia.

Types. The syntypes are stored in the Croatian Natural History Museum, Zagreb; no number indicated (Milan et al. 1974: 92).

Remarks. Brusina (1874) divided the species *M. inconstans* Neumayr, 1869 into three varieties, none of which he termed “*inconstans*”. The first one, “var. *costulata*”, he referred to as the typical one, which makes it an objective synonym of the nominal subspecies and hence *M. inconstans*.

† *Melanopsis visianiana* f. *costulata* Brusina, 1897 [invalid]

Original source. Brusina 1897: 12, pl. 5, fig. 6.

Type horizon. Langhian, middle Miocene.

Type locality. “Miočić”, Croatia.

Types. Milan et al. (1974: 99) indicated a holotype, but it is uncertain whether the specimen was the only one Brusina had at hand (holotype by monotypy, Art. 73.1.2). The specimen is stored in the Croatian Natural History Museum, Zagreb, coll. no. 2981-627/2.

Remarks. Junior homonym of *Melanopsis inconstans costulata* Brusina, 1874 (see Note 1). Currently considered as a junior synonym of *M. visianiana* Brusina, 1874 (Neubauer et al. 2016b: 25).

***Melanopsis laevigata* var. *costulata* Pallary, 1899 [invalid]**

Original source. Pallary 1899: 137.

Type locality. “Souani près Tanger” [Souani near Tanger], Morocco.

Remarks. Junior homonym of *Melanopsis costulata* Brusina, 1897.

† *Melanopsis staudi* var. *costulata* Brusina, 1903 [invalid]

Original source. Brusina 1903: 115.

Type horizon. Late Pleistocene–early Holocene.

Type locality. “Bischofsbad” [Püspökfürdő, Băile 1 Mai, Lake Pețea], Romania.

Remarks. Junior homonym of *Melanopsis costulata* Brusina, 1897. Neubauer et al. (2014d: 125) considered this taxon as a junior synonym of *Microcolpia parreyssii sikorai* (Brusina, 1903).

† *Melanopsis (Coptostylus) costulata* Doncieux, 1908 [invalid]

Original source. Doncieux 1908: 205, pl. 11, figs 13a–13b.

Type horizon. Middle Lutetian, Eocene.

Type locality. “Au Nord d’Albas” (Doncieux 1908: 205), France.

Remarks. Junior homonym of *Melanopsis costulata* Brusina, 1897. Has been considered to belong to the genus *Coptostylus* Sandberger, 1872 (Thiaridae).

Melanopsis vondeli var. *costulata* Pallary, 1928 [invalid]

Original source. Pallary 1928b: 16, pl. 2, fig. 13.

Type locality. “O. Taguenout” [Oued Taguenout, said to be near Beni Mellal], Morocco.

Remarks. Junior homonym of *Melanopsis costulata* Brusina, 1897.

Melanopsis (Mesopotamia) infracincta var. *costulata* Pallary, 1939 [invalid]

Original source. Pallary 1939: 104, pl. 4, fig. 13.

Type locality. “Ras el ‘Ain du Khabour” [Chabur river near Ra’s al ‘Ayn], Syria.

Remarks. Junior homonym of *Melanopsis costulata* Brusina, 1897.

† *Melanopsis cotrocenensis* Cobălcescu, 1883

Original source. Cobălcescu 1883: 123, pl. 9, figs 8a–b.

Type horizon. Cernikian, Pliocene.

Type locality. “Cotroceni lângă București” [Cotroceni near Bucarest], Romania.

Melanopsis coupfa Bourguignat, 1884

Original source. Bourguignat 1884: 94.

Type locality. “Dans les eaux chaudes du Djerid, au nord du chott Tiraoun, dans le sud de la Tunisie” [in the warm water of Djérid, north of chott Tiraoun], Tunisia.

***Microcolpia coutagniana* Bourguignat, 1884**

Original source. Bourguignat 1884: 56.

Type locality. “Lac Sabandja, près d’Ismidt, en Anatolie” [Lake Sapanca near İzmit], Turkey.

† *Melanopsis covurluensis* Cobălcescu, 1883

Original source. Cobălcescu 1883: 123, pl. 9, figs 7a–d.

Type horizon. Pliocene.

Type locality. “Barboschi” (p. 156) [Barboși], Romania.

***Melania crassa* Brusina, 1866 [invalid]**

Original source. Brusina 1866: 106.

Type locality. “In der Muhr” (Pfeiffer 1828: 48) [in the river Mur], Austria.

Remarks. Brusina (1866) introduced the name for “var. A” in Pfeiffer (1828), which that author, however, had affiliated with the name “*Melania agnata* Ziegler” (see Pfeiffer 1828: 67). Therefore, *Melania crassa* Brusina, 1866 is a junior objective synonym of *Melania agnata*. Moreover, it is a junior homonym of *Melania crassa* Münster, 1841. Note that the latter species is certainly no Melanopsidae as it derives from the marine deposits of the Triassic St. Cassian Fm.

† *Melanopsis anceps* var. *crassicosta* Gaudry & P. Fischer in Gaudry, 1867

Original source. Gaudry 1862–1867: 446.

Type horizon. Pliocene.

Type locality. “Mégare” (p. 444), Greece.

***Melanella crassilabris* Bourguignat, 1884**

Original source. Bourguignat 1884: 16.

Type locality. “La Save à Agram et rivière d’Ostaria en Croatie” [Sava river at Zagreb and Oštarije river], Croatia.

Remarks. Note that Bourguignat denoted the authority as “Bourguignat, 1879”.

† *Melanopsis crassitesta* Blanckenhorn, 1897

Original source. Blanckenhorn 1897: 134, pl. 10, fig. 18.

Type horizon. Plio-Pleistocene.

Type locality. “In der pliocänen Dreissensiaschicht von Dschir esch-Schurr” [in the pliocene *Dreissena* layer at Jisr Ash-Shughur], Syria.

† *Melanopsis crastina* Vidal, 1874

Original source. Vidal 1874: 235, pl. 2, fig. 12, pl. 5, figs 32–34.

Type horizon. Maastrichtian, Cretaceous.

Type locality. “Isona”, Spain.

Melanopsis crenocarinata Moricand, 1841

Original source. Moricand 1841: 61, pl. 4, figs 10–11.

Type locality. “Rio de Pedra Branca, procince de Bahia” [Pedra Branca river, province Bahia], Brazil.

Remarks. Type species of the genus *Verena* H. Adams & A. Adams, 1854 (Thiariidae) (see Nuttall 1990: 253).

† *Melanopsis maresi* var. *crenulata* Pallary, 1901

Original source. Pallary 1901a: 180, pl. 2, fig. 22.

Type horizon. Pleistocene.

Type locality. “De Géryville” [Aïn Sefra], Algeria.

Melanopsis fasensis var. *cristata* Pallary, 1920

Original source. Pallary 1920c: 147, pl. 4, fig. 13.

Type locality. “Fès”, Morocco.

† *Melanopsis croatica* Brusina, 1884

Original source. Brusina 1884a: 168.

Type horizon. Cernikian, Pliocene.

Type locality. “Repušnica” (Neumayr 1869: 372), Croatia.

Types. Milan et al. (1974: 89) indicated a holotype, but it is uncertain whether the specimen actually derives from the original type series and whether it was the only specimen Brusina had at hand. The specimen is stored in the Croatian Natural History Museum, Zagreb, coll. no. 2988-634.

Remarks. Introduced for *M. costata* sensu Neumayr, 1969, non Olivier, 1804. Wenz (1929) considered all records of “*M. costata*” from the Pliocene of Slavonia to represent the same species and synonymized them with *M. abbreviata cosmanni*. Obviously, he was unaware that *M. croatica* Brusina, 1884 is the first available name for them (see also discussion of *M. pseudocostata* Oppenheim, 1890).

† *Melanopsis austriaca croatica* Brusina, 1902 [invalid]

Original source. Brusina 1902: pl. 6, figs 71–72.

Type horizon. Middle Pannonian, late Miocene.

Type locality. “Markuševac”, Croatia.

Types. The illustrated syntypes are stored in the Croatian Natural History Museum, Zagreb, coll. no. 2529-175/1-2 (Milan et al. 1974: 86).

Remarks. Junior homonym of *Melanopsis croatica* Brusina, 1884. Wenz (1930) introduced *M. haueri markusevecensis* as replacement name.

† *Melanopsis cuisienensis* Dominici & Kowalke, 2014

Original source. Dominici and Kowalke 2014: 147, pl. 2, figs 5a–b.

Type horizon. Castigaleu group, late Ypresian, Eocene.

Type locality. “Morillo de Lena (Esera valley, Tremp-Graus basin), [...] CG-A2, sample 53”, Spain.

Types. Museo di Storia Naturale, Università degli Studi di Firenze, coll. no. IGF 4334E.

Melanopsis curta Gassies, 1870

Original source. Gassies 1870: 146.

Type locality. “Tuo” [Touho], New Caledonia.

Remarks. Brot (1879: 444) considered the taxon as a junior synonym of *Melanopsis frustulum* Morelet, 1857.

Melanopsis turcica var. *curta* Locard, 1883 [invalid]

Original source. Locard 1883a: 270.

Type locality. “Lac d’Antioche” [Lake Anuk (also as Amik)], Turkey.

Remarks. Junior homonym of *Melanopsis curta* Gassies, 1870.

***Melanopsis cerithiopsis* var. *curta* Bourguignat, 1884 [invalid]**

Original source. Bourguignat 1884: 131.

Type locality. “Plaine du Bahr-el-Houlé (haut Jourdain) dans l’Aïn-el-Mellaha” [in the plains of the Hula valley (upper Jordan), in Aïn Mallahah], Israel.

Remarks. Junior homonym of *Melanopsis curta* Gassies, 1870. Heller et al. (2005: 248) considered the variety as a junior synonym of *Melanopsis saulcyi* Bourguignat, 1853.

† ***Melanopsis callosa* var. *curta* Locard, 1893 [invalid]**

Original source. Locard 1893: 181, pl. 9, fig. 20.

Type horizon. Late Burdigalian–Langhian, early–middle Miocene.

Type locality. “Le Locle; [...] Vermes”, France.

Remarks. Junior homonym of *Melanopsis curta* Gassies, 1870. Wenz (1929: 2764) considered the variety as a junior synonym of *Melanopsis kleini* Kurr, 1856.

“*Melanopsis praemorsa* f. *curta*” mentioned in Pérès (1939) [unavailable]

Locality. “Station 119. Aïn Attig. Source près de la route de Rabat à Casablanca à 13 kilomètres de Rabat” [station 119 at Ain Attig. A spring near the road from Rabat to Casablanca, 13 km from Rabat], Morocco.

Remarks. First of all, the name as given by Pérès (1939) is a nomen nudum – Pérès apparently considered the expression “*curta*” self-explanatory and did not describe it. Moreover, he obviously used the name not as separate taxon but rather as descriptive term to fit existing species into his morphological concept. He even indicated *M. brevis* Morelet, 1857 as its “type”.

“*Melania cuspidata* Parreyss” mentioned in Brot (1874–1879: 12) [unavailable]

Locality. Not indicated.

Remarks. Nomen nudum, “in schedis” name from Parreyss listed in the synonymy list of “*Melania holandri*” [sic] by Brot (1874).

† ***Melanopsis cvijici* Brusina, 1902**

Original source. Brusina 1902: pl. 29, figs 19–22.

Type horizon. Langhian, middle Miocene.

Type locality. “Vatelj” [Fatelj hill], Bosnia and Herzegovina.

Types. Milan et al. (1974: 89) stated that only one of the specimens illustrated by Brusina (1902: pl. 29, figs 19–20) has been preserved, which they designated as lectotype. The specimen is stored in the Croatian Natural History Museum, Zagreb, coll. no. 2900-546/1.

† *Melanopsis lyrata* var. *cylindracea* Brusina, 1874

Original source. Brusina 1874: 45.

Type horizon. Langhian, middle Miocene.

Type locality. “Ribarić”, Croatia.

Remarks. Currently considered as a junior synonym of *Melanopsis lyrata* Neumayr, 1869 (Neubauer et al. 2016b: 24; see also Wenz 1929: 2698).

† *Melanopsis cylindrata* Blanckenhorn, 1897

Original source. Blanckenhorn 1897: 136, pl. 10, figs 22–24.

Type horizon. Plio-Pleistocene.

Type locality. “Im Rab zwischen Dschir esch-Schurr und Kal ‘at el-Mdik; [...] in der pliocänen Dreissensiaschicht von Dschir esch-Schurr auf dem rechten Orontesufer” [in the Al Ghāb between Jisr Ash-Shughur and Qal’at al Maḍīq; in the Pliocene *Dreissena* layer at Jisr Ash-Shughur at the right riverbank of the Orontes], Syria.

† *Lyrcea cylindrica* Stoliczka, 1862

Original source. Stoliczka 1862: 537, pl. 17, fig. 9.

Type horizon. Transdanubian, Pannonian, late Miocene.

Type locality. “Bei Zala Apati; [...] in der Umgegend des Plattensees, wie auf der Halbinsel Tihany” [near Zalaapáti; in the surroundings of Lake Balaton, as well as on the Tihany peninsula], Hungary.

† *Melanopsis (Duabiana) cylindrica* Anistratenko, 1993 [invalid]

Original source. Anistratenko 1993: 69.

Type horizon. Duab Beds, middle to late Kimmerian, Pliocene.

Type locality. “Okp. c. Mokbu, Очамчирский р-н” [near the village Mok’vi, Ochamchirskiy rayon], Georgia.

Types. Schmalhausen Institute of Zoology of National Academy of Sciences of Ukraine, Kiev; no number indicated.

Remarks. Type species of *Duabiana* Starobogatov & Anistratenko in Anistratenko, 1993. Junior secondary homonym of *Lyrcea cylindrica* Stoliczka, 1862. Neubauer et al. (2014a: 456) introduced *M. anistratenkoi* as replacement name.

† *Melanopsis (Duabiana) cylindrospica* Anistratenko, 1993

Original source. Anistratenko 1993: 71, textfig. 1.

Type horizon. Duab Beds, middle to late Kimmerian, Pliocene.

Type locality. “Окр. с. Мокви, Очамчирский р-н” [near the village Mok’vi, Ochamchirskiy rayon], Georgia.

Types. Schmalhausen Institute of Zoology of National Academy of Sciences of Ukraine, Kiev; no number indicated.

† *Melanoptychia dalmatina* Bourguignat, 1880

Original source. Bourguignat 1880: 31.

Type horizon. Langhian, middle Miocene.

Type locality. “Vallée de la Cettina” [Cetina river valley], Croatia.

Remarks. The taxon is not included in the Fossilium Catalogus of Wenz (1929).

† *Melanopsis dalmatina* Brusina, 1884 [invalid]

Original source. Brusina 1884b: 55.

Type horizon. Early Langhian, middle Miocene.

Type locality. “Oberes Niveau v. Župića potok” (p. 47) [upper horizon of the Župića potok (near Sinj)], Croatia.

Types. Milan et al. (1974: 90) stated that Brusina (1897) had indicated one of the specimens illustrated by him (pl. 5, fig. 10) as type. However, it is uncertain whether the specimen actually derives from the original type series. The specimen is stored in the Croatian Natural History Museum, Zagreb, coll. no. 2984-630/1.

Remarks. Junior secondary homonym of *Melanoptychia dalmatina* Bourguignat, 1880.

Fagotia danubialis Bourguignat, 1884

Original source. Bourguignat 1884: 35.

Type locality. “Le Danube à Ibraila; la Save à Agram; la Krapina à Sused (Croatie)” [Danube river at Brăila (Romania); Sava river at Zagreb; Krapina river at Podused, in Zagreb (Croatia)].

Remarks. Note that Bourguignat denoted the authority as “Bourguignat, 1880”.

† *Melanopsis daphnes* Gaudry & P. Fischer in Gaudry, 1867

Original source. Gaudry 1862–1867: 407, pl. 62, figs 16–18.

Type horizon. Late Miocene.

Type locality. “Daphné”, Greece.

Remarks. The species epithet is a noun in apposition and needs not to agree in gender with the generic name (Art. 31.2.1). The name “*daphne*” as mentioned in Neumayr (1869: 369) and Hoernes (1876: 16) is an incorrect subsequent spelling. Wenz (1929: 2772) considered the taxon as a junior synonym of *Melanopsis longa* Deshayes in Férussac, 1839.

Melanopsis daudebartii [Prevost], 1821

Original source. [Prevost] 1821: 137.

Type locality. “Auprès de Baden, en Autriche, dans un bassin d’eau thermale sulfureuse” [near Baden, Austria, in a sulphurous thermal water basin], Austria.

Remarks. There is considerable uncertainty about the correct authority and spelling of this species in the literature. It was first mentioned and validly described in the year 1821 as “*Melanopsis Daudebartii*” in an article in the Bulletin des Sciences, par la Société philomatique de Paris. Constant Prevost was often considered to be the author of this article, but from the title and text it is obvious that the article is an “Extrait” of a talk given by Prevost earlier and summarized by an anonymous author. According to Art. 50.2 and Recommendation 51D, the correct citation should be *M. daudebartii* [Prevost], 1821. The names “*daudebarti*”, “*audebarti*” or “*audebardi*”, each occurring multiple times in the literature, are incorrect subsequent spellings. Currently, the species is classified within the genus *Microcolpia* (see also Neubauer et al. 2014d: 126).

† *Melanopsis dautzenbergi* Pallary, 1901

Original source. Pallary 1901a: 179, pl. 2, fig. 25.

Type horizon. Pleistocene.

Type locality. “De l’Oued Tiout (Sud-Oranais)” [Tiout], Algeria.

Melanopsis debilis Pallary, 1920

Original source. Pallary 1920a: 33.

Type locality. “Sidi Yahia, près d’Oudjda” [Sidi Yahya near Oujda], Morocco.

***Melanopsis (Mesopotamia) infracincta* var. *debilis* Pallary, 1939 [invalid]**

Original source. Pallary 1939: 104.

Type locality. “Ras el ‘Ain du Khabour” [Chabur river near Ra’s al ‘Ayn], Syria.

Remarks. Junior homonym of *Melanopsis debilis* Pallary, 1920.

† ***Melanopsis decipiens* P. Fischer, 1883**

Original source. Fischer 1883: 60, pl. 3, fig. 3.

Type horizon. Late Miocene.

Type locality. “Smendou, province de Constantine, Algérie” [Zighoud Youcef], Algeria.

Remarks. Fischer attributed the authority to Tournouër, but from the foregoing introduction it is clear that the species was described by Fischer.

† ***Melanopsis decollata* Stoliczka, 1862**

Original source. Stoliczka 1862: 536, pl. 17, fig. 8.

Type horizon. Late Transdanubian, Pannonian, late Miocene.

Type locality. “Bei Zala Apati am rechten Ufer der Zala und [...] im Gebiete des Plattensees” [near Zalaapáti at the right riverside of the Zala river and in the area around Lake Balaton], Hungary.

***Melanopsis praerosa* [sic] var. *decollata* Paetel, 1888 [invalid]**

Original source. Paetel 1888: 403.

Type locality. “Persia” (Nevill 1884: 209), Iran.

Remarks. Originally introduced as infrasubspecific taxon (“subvariety”) by Nevill (1884), but made available by Paetel (1888) who treated it as variety (Art. 45.5.1). Paetel clearly referred to the description of Nevill. Junior homonym of *M. decollata* Stoliczka, 1862.

† ***Melanopsis bouei* var. *decorata* Pallary, 1916**

Original source. Pallary 1916: 85.

Type horizon. Late Miocene or Pliocene.

Type locality. Sabba Stefanescu, to whom Pallary referred, did not denote the localities of the figured specimens. He reported *M. bouei* from many localities in Romania.

Remarks. Introduced for a part of Stefanescu’s material of *M. bouei* (Stefanescu 1896: pl. 11, figs 63–64). Wenz (1929: 2681) considered the taxon as a junior synonym of *Melanopsis bouei* Férussac, 1823.

***Melanopsis douttei* var. *decorata* Pallary, 1920 [invalid]**

Original source. Pallary 1920c: 150, pl. 4, fig. 2.

Type locality. “La Makina”, Morocco.

Remarks. Junior homonym of *Melanopsis decorata* Pallary, 1916.

† *Melanopsis decostata* Penecke, 1884

Original source. Penecke 1884: 22, pl. 10, fig. 7.

Type horizon. Cernikian, Pliocene.

Type locality. “Repusnica” [Repušnica], Croatia.

Remarks. The name was also marked as new taxon by Brusina (1902: vii). It is uncertain whether Brusina really intended to introduce a new subspecies or actually referred to the species of Penecke (1884), which he had ranked as subspecies of *M. croatica* earlier (Brusina 1897).

***Melanopsis decussata* Férussac, 1823**

Original source. Férussac 1823: 159.

Type locality. “Plattensée, en Hongrie [...] ; à Sary Maydan Zakrzewski, dans le gouvernement de Podolie, non loin de Kamieniec-Podolsk” [Lake Balaton; at Saryy Zakrevskiy Maydan, not far from Kam’yanets’-Podil’s’kyi], Hungary.

Remarks. Considered as a junior synonym of “*Hemisinus Esperi*” Férussac (1823) by Brot (1878: 372).

† *Melanopsis defensa* Fuchs, 1870

Original source. Fuchs 1870a: 353, pl. 14, figs 77–79.

Type horizon. Transdanubian, Pannonian, late Miocene.

Type locality. “Radmanest” [Rădmănești], Romania.

***Melanopsis costata* var. *degenerata* Preston, 1914**

Original source. Preston 1914: 467, pl. 27, fig. 9.

Type locality. “Lake of Tiberias [Sea of Galilee] at the exit of the Jordan”, Israel.

† *Melanopsis (Lyrcea) delessei* Tournouër, 1875

Original source. Tournouër 1875: 77.

Type horizon. Tafi Formation, early Pleistocene.

Type locality. “Prope Antimaki” [near Antimácheia, Kos Island], Greece.

† *Melanopsis delicata* Pallary, 1920

Original source. Pallary 1920b: 112.

Type horizon. Transdanubian, Pannonian, late Miocene.

Type locality. “Radmanest” (Brusina 1902: captions of pl. 29) [Rădmănești], Romania.

Remarks. Replacement name for *M. boettgeri* Brusina, 1902, non Klika, 1891. Jekeilius (1944: 74) considered this species as a junior synonym of *M. sturii* Fuchs, 1873.

Melanopsis delmasi Pallary, 1936

Original source. Pallary 1936: 55, pl. 3, fig. 3.

Type locality. “Dans les canaux de l’Aguedal, à Marrakech” [in the channels of Aguedal in Marrakech], Morocco.

Melanopsis denegabilis Pallary, 1939

Original source. Pallary 1939: 85, textfig. 11, pl. 6, figs 9–13.

Type locality. “En Iraq; [...] à Rahalya Springs, à Kani Seip et à Karsī” [in Iraq, at Rahalya springs (?), at Kānī Seip (?) and at Karsī], Iraq.

Remarks. Pallary attributed the authority to Férussac based on a manuscript name.

† *Melanopsis deperdita* de Serres, 1829

Original source. De Serres 1829: 101.

Type horizon. Early Campanian, Cretaceous.

Type locality. “Martigues”, France.

† *Melanopsis depereti* Boistel, 1898

Original source. Boistel 1898: 28 (footnote), figs 3A–D.

Type horizon. Mammal zone MN 10–12, late Miocene.

Type locality. “D’Ambronay [Vallon de Jurancieu]” [from Ambronay, in the valley of Jurancieu], France.

† *Melanopsis depressa* Pallary, 1916

Original source. Pallary 1916: 77.

Type horizon. Eocene?

Type locality. “L’île de Wight” [Isle of Wight], United Kingdom.

Remarks. Based on the record of “*Melanopsis buccinoidea* var. γ) antiqua; *elongata*” sensu Férussac, 1823 (pl. 7, fig. 6).

† *Melanopsis bouei* var. *depressa* Pallary, 1916 [invalid]

Original source. Pallary 1916: 78.

Type horizon. Pannonian, late Miocene.

Type locality. “De la Moravie” (Férussac 1823: 164; no precise locality indicated), Czech Republic.

Remarks. Based on a part of Férussac’s (1823: pl. 8, fig. 10) material of *M. bouei*. Homonym of the simultaneously published *Melanopsis depressa* Pallary, 1916, which has the higher rank in the species group and thus takes precedence (Art. 24.1).

† *Lyrcaea* [sic] *narzolina* var. *dertocylindrica* Sacco, 1895

Original source. Sacco 1895: 14, pl. 1, fig. 25.

Type horizon. Tortonian–early Messinian, late Miocene.

Type locality. “S. Agata fossili” [Sant’Agata Fossili], Italy.

Remarks. Harzhauser et al. (2015: 9) considered the taxon as a junior synonym of *Melanopsis narzolina* d’Archiac in Viquesnel, 1846.

† *Lyrcaea* [sic] *pedemontana* var. *dertoliva* Sacco, 1895

Original source. Sacco 1895: 11, pl. 1, fig. 19.

Type horizon. Tortonian–early Messinian, late Miocene.

Type locality. “S. Agata fossili” [Sant’Agata Fossili], Italy.

Remarks. The name “*dertolina*” as mentioned in Wenz (1929: 2802) is an incorrect subsequent spelling.

***Melanopsis deserticola* Annandale & Prashad, 1919**

Original source. Annandale and Prashad 1919: 37, pl. 3, fig. 8.

Type locality. “Kaindak (long. 60°48'E., lat. 29°48'N.), Persian Baluchistan”, Iran.

Types. Indian Museum, Calcutta, coll. no. 11535/2.

***Melanopsis desertorum* Bourguignat, 1884**

Original source. Bourguignat 1884: 134.

Type locality. “Ruisseaux entre Tarsous et Mersina (Anatolie)” [streams between Tarsus and Mersin], Turkey.

Remarks. Heller et al. (1999: 56) considered the species as a junior synonym of *Melanopsis buccinoidea* (Olivier, 1801). In Heller et al. (2005: 248) in turn it is treated as a junior synonym of *Melanopsis saulcyi* Bourguignat, 1853.

***Melanopsis deshayesiana* Gassies, 1861**

Original source. Gassies 1861: 292, pl. 7, fig. 12.

Type locality. “La Nouvelle-Calédonie, dans l’intérieur” [inland of New Caledonia], New Caledonia.

† *Melanopsis esperi* var. *desori* De Stefani, 1877

Original source. De Stefani 1877: 310, pl. 18, fig. 9.

Type horizon. Villafranchian, Plio-Pleistocene.

Type locality. “Spoleto”, Italy.

“*Melanella hollandri* [sic] var. *detrita* Kucik” mentioned in Brusina (1867: 85) [unavailable]

Locality. “U Savi kod Zagreba” [from the Sava river at Zagreb], Croatia.

Remarks. Nomen nudum, based on an “in schedis” name in the collection of Kucik (also read as “Kutschig”). Brot (1874: 11) listed it in synonymy of *Melania holandri* [sic].

***Melanopsis diabetensis* Pallary, 1915**

Original source. Pallary 1915: 28.

Type locality. “Dans la source du jardin du Sultan, à Diabet, près Mogador” [in the springs in the garden of the Sultan at Douar Dyabat, near Essaouira], Morocco.

***Melanopsis sesteri* var. *diadema* Bourguignat, 1884**

Original source. Bourguignat 1884: 119.

Type locality. “Petit cours d’eau à Sadjour-Sou, entre Aïn-Taïb et Alep [...]; Aïn-el-Bass, dans la plaine du Bahr-el-Houlé (Syrie)” [small brook at Sadjour-Sou between Gaziantep (Turkey) and Aleppo (Syria) [...]; Aïn el Bass, in the plains of the Hula valley (Israel)].

† *Melanopsis dianaeformis* Andrusov, 1909

Original source. Andrusov 1909: 84, 159, pl. 4, figs 21–27.

Type horizon. Pontian (sensu stricto), late Miocene.

Type locality. “Babadjan, Sundi, Meissary und Chilaalidasch” [Babadzhan, Syundi, Meysary canyon, Mount Chila-alidasch], Azerbaijan.

† *Melanopsis dichtli* Handmann, 1882

Original source. Handmann 1882: 555.

Type horizon. Pannonian, zone D, late Miocene.

Type locality. “Kottingbrunn”, Austria.

Melanella letourneuxi var. *dilatata* Bourguignat, 1884

Original source. Bourguignat 1884: 26.

Type locality. “Ogulin”, Croatia.

Melanopsis buccinoidea var. *dilatata* Pallary, 1899

Original source. Pallary 1899: 138, pl. 9, fig. 10.

Type locality. “Dans la Souani, à Tanger” [in Souani at Tanger], Morocco.

Melanopsis dircaeana Pallary, 1939

Original source. Pallary 1939: 87, pl. 6, figs 31–35.

Type locality. Unclear: given as “Dans l’Oronte” [in the Orontes river] in text but as “Du lac de Homs” [Lake Homs (through which the Orontes flows)] in plate captions, Syria.

Remarks. The name “*dircaeana*” as mentioned in Heller et al. (2005: 238) is an incorrect subsequent spelling.

† *Melanopsis dispar* Deshayes, 1862

Original source. Deshayes 1861–1864: 473, pl. 31, figs 29–30.

Type horizon. Lutetian, Eocene.

Type locality. “Brasles”, France.

Remarks. Cossmann (1888: 279) and Wenz (1929: 2639) classified the species in the genus *Faunus* Montfort, 1810 (Pachychilidae).

† *Melanopsis dissimilis* Pallary, 1916

Original source. Pallary 1916: 85.

Type horizon. Langhian, middle Miocene.

Type locality. “Ribarić” (Neumayr 1869: 358), Croatia.

Remarks. Replacement name for *M. lyrata* Neumayr, 1869 [June], non *M. lirata* Gassies, 1869 [January]. Both names are deemed to be identical after Art. 58.2.

Melanella divina Bourguignat, 1884

Original source. Bourguignat 1884: 11.

Type locality. “Mare du moulin de la Cettina, près Almissa, en Dalmatie” [mill-pond at the river Cetina, near Omiš], Croatia.

Remarks. Bourguignat denoted the authority as “Letourneux, 1879”, but there is no evidence that the description really derived from that author.

Fagotia (Dneprifagotia) dnezensis Starobogatov, Alexenko & Levina, 1992

Original source. Starobogatov et al. 1992: 63, figs 1 (2), 2 (4), 3 (11).

Type locality. “Из Днепра у Херсона” [from the Dniepr river at Kherson], Ukraine.

Types. Zoological Institute of Russian Academy of Sciences, St.-Petersburg; no number indicated.

† *Melanopsis doboi* Schréter, 1975

Original source. Schréter 1975: 7, textfig. 1, pl. 1, figs 3–6.

Type horizon. Riss/Würm end to early Würm Ice Age, Pleistocene.

Type locality. “Eger, az egri vár Zárkándy bástyájának átmetszése” [Eger, section at the Zarkandy bastion of the fortress Eger], Hungary.

Types. Magyar Állami Földtani Intézet (Hungarian Geological Museum), Budapest; no number indicated.

Remarks. Neubauer et al. (2016a) attributed the species to the genus *Microcolpia*.

† *Melanopsis matheroni* var. *doderleini* Pantanelli, 1886

Original source. Pantanelli 1886a: 69 or 1886b: 78, pl. 3, figs 5, 8 (precedence not established).

Type horizon. Tortonian–early Messinian, late Miocene.

Type locality. “S. Valentino, S. Agata, Boggione” [San Valentino, Sant’Agata Fossili, Boggione near Siena], Italy.

Remarks. Harzhauser et al. (2015: 9) considered the taxon as a junior synonym of *Melanopsis narzolina* d’Archiac in Viquesnel, 1846.

Melanopsis (Mesopotamia) khabourensis var. *dolichosoma* Pallary, 1939

Original source. Pallary 1939: 103, pl. 5, fig. 18.

Type locality. “Ras el ‘Ain du Khabour” [Chabur river near Ra’s al ‘Ayn], Syria.

† “*Melanopsis doliolum*” mentioned in Graves (1847: 600) [unavailable]

Horizon. Eocene.

Locality. “Cuisse-Lamotte, Jaulzy, Tiverny, Saint-Vaast-de-Longmont”, France.

Remarks. Nomen nudum. Graves attributed the authority to Defrance.

† *Melanopsis (Canthidomus) bouei* var. *doliolum* Handmann, 1887

Original source. Handmann 1887: 35, pl. 7, figs 6–7.

Type horizon. Pannonian, zone B–D, late Miocene.

Type locality. “Leobersdorf”, Austria.

† *Melanopsis doncieuxi* Pallary, 1916

Original source. Pallary 1916: 80.

Type horizon. Lutetian, Eocene.

Type locality. “Au Nord d’Albas” (Doncieux 1908: 204), France.

Remarks. Replacement name for *M. nodosa* Doncieux, 1908, non Férussac, 1822.

† *Melanopsis (Stylospirula) doncieuxi* Wenz, 1919 [invalid]

Original source. Wenz 1919b: 65.

Type horizon. Middle Lutetian, Eocene.

Type locality. “Au Nord d’Albas” (Doncieux 1908: 204), France.

Remarks. Replacement name for *M. brevis* Doncieux, 1908, non Sowerby, 1826. For that homonym, Pallary (1916) had introduced the replacement name *M. abbreviata*, which is a junior homonym of *M. abbreviata* Brusina, 1874. (Note that Wenz 1919 was unaware of that name.) *M. doncieuxi* Wenz, 1919 is, however, invalid too it is a junior homonym of *M. doncieuxi* Pallary, 1916 (see *M. atatica* Wenz, 1928).

***Melanopsis doriae* Issel, 1865**

Original source. Issel 1865 [page unknown; original work not seen; described on p. 400 of the re-published version from 1866].

Type locality. “Di Kerman nella Persia meridionale” [Kerman], Iran.

Remarks. Considered a junior synonym of *M. ammonis* Tristram, 1865 by Heller et al. (2005). Considered a synonym of *M. variabilis* v.d. Busch in Philippi, 1847 by Martens (1874).

† ***Melanopsis (Macrospira) doroghensis* Oppenheim, 1892**

Original source. Oppenheim 1892: 705, pl. 33, figs 7–11.

Type horizon. Paleocene.

Type locality. “Dorogh, Annathal, Nagy Kovacs” [Dorog, Annavölgy, Nagykovácsi], Hungary.

***Melanopsis dos* Preston, 1913**

Original source. Preston 1913: 436.

Type locality. “Island of Beilan-Beilan, to the north of the Obi Islands, Dutch East Indies” [Belangbelang Island], Indonesia.

Remarks. Probably not a Melanopsidae.

† ***Melanopsis (Smendovia) doumerguei* Pallary, 1901**

Original source. Pallary 1901a: 177, pl. 2, fig. 24.

Type horizon. Late Miocene.

Type locality. “Smendou” [Zighoud Youcef], Algeria.

***Melanopsis doumeti* Letourneux & Bourguignat, 1887**

Original source. Letourneux and Bourguignat 1887: 157.

Type locality. “De Gafsa; [...] de Tozer et de Nefta” [from Gafsa; in Tozeur and Nafta], Tunisia.

***Melanopsis douttei* Pallary, 1911**

Original source. Pallary 1911: 133, [unnumbered plate], figs 21–22.

Type locality. “Fez” [Fes], Morocco.

Remarks. Given as “*doutte*” on p. 133, but as “*douttei*” in plate caption. Since Pallary explicitly named the species after E. Doutté, the name must read “*douttei*” (Art. 32.5.1).

† ***Melanopsis douvillei* Pallary, 1916**

Original source. Pallary 1916: 83.

Type horizon. Maastrichtian, Cretaceous.

Type locality. “Du versant oriental du Kouh Mapeul” (Douvillé 1904: captions of pl. 46) [eastern slope of mount Kuh-e Mapel, c. 60 km WNW of Khorramābād], Iran.

Remarks. Replacement name for *M. costellata* Douvillé, 1904, non Férussac, 1823.

† ***Melanopsis draghicheniani* Cobălcescu, 1883**

Original source. Cobălcescu 1883: 124, pl. 9, figs 9a–b.

Type horizon. Dacian, Pliocene.

Type locality. “De Bécéni” [Beceni], Romania.

Remarks. The name “*draghicheniana*” as mentioned in Wenz (1929: 2709) is an incorrect subsequent spelling.

***Melanopsis (Sistaniana) drangianensis* Starobogatov & Izzatullaev, 1985**

Original source. Starobogatov and Izzatullaev 1985: 34, fig. 4.

Type locality. “Cucmah” [Sistan region], Iran.

Types. Zoological Institute of Russian Academy of Sciences, St.-Petersburg; no number indicated.

† ***Melanopsis dubia* Stoliczka, 1860**

Original source. Stoliczka 1860: 486, pl. 1, figs 14, 15a–b.

Type horizon. Late Turonian, late Cretaceous.

Type locality. “Neualpe im Russbachthal” [Neualm near Russbach am Pass Gschütt], Austria.

† *Melanopsis dubiosa* Matheron, 1879

Original source. Matheron 1878–1880: pl. O1, fig. 14.

Type horizon. Cuisian, late Ypresian, early Eocene.

Type locality. “Brignan, couches inférieures du Montaiguët et de Grabels; lignite de la Caunette” (it is unclear if this statement describes a single locality or several and, if so, from which the species derives), France.

† *Melanopsis (Canthidomus) dubiosa* Kühn, 1963 [invalid]

Original source. Kühn 1963: 387, pl. 3, figs 12–13.

Type horizon. Mammal zone MN 9–10, late Miocene.

Type locality. “Daphni” [Dafní valley near Athens], Greece.

Types. Museum of Palaeontology and Geology of the University of Athens, coll. no. 1963/84.

Remarks. Junior homonym of *Melanopsis dubiosa* Matheron, 1878. Papp (1979: 668) considered the species as a junior synonym of *Melanopsis longa* Deshayes in Férussac, 1839.

Melanopsis ducosi Pallary, 1916

Original source. Pallary 1916: 82.

Type locality. “Dans le Diahot, à Balade; à Jengen, à Kanala, dans les marais et les petits ruisseaux” (Gassies 1861: 289) [in the Diahot river at Balade; at Hienghène, at Canala, in swamps and small streams], New Caledonia.

Remarks. Replacement name for *M. carinata* Gassies, 1861, non Sowerby, 1826.

† *Melanopsis dufourii* Férussac, 1822

Original source. Férussac 1819–1851: Mélanopsides fossiles, pl. 1 (1822), fig. 16.

Type horizon. Burdigalian, early Miocene (?).

Type locality. “Des environs de Dax” [surroundings of Dax], France.

Remarks. The name first appeared in 1822 on the captions for plate 1 of the “Mélanopsides fossiles” in Férussac’s “Histoire naturelle” (see also introduction for details). While Férussac (1823) included also recent specimens under that name in his monograph on the Melanopsidae, the only specimen illustrated in 1822 in the “Histoire naturelle” was a fossil one from the Miocene of France. This fact remained widely unknown to biologists and paleontologists alike. Only few malacologists, such as Grateloup (1828, 1838, 1840), used the name correctly and attributed it to the French fossils. Consequently, none of the specimens referred to as *M. dufourii* in the biological literature actually corresponds to the real *M. dufourii*. Very likely, some of

the other melanopsids from Dax (see, e.g., Grateloup 1838, d’Orbigny 1852, Pallary 1916) are unrecognized junior synonyms of this species.

The names “*dufouri*” as mentioned by numerous authors and databases and “*dufoureri*” as given by Morelet (1853: 297) and Bourguignat (1884: 114) are incorrect subsequent spellings.

† *Melanopsis dufresnii* Deshayes, 1825

Original source. Deshayes 1824–1837: 120, pl. 12, figs 3–4.

Type horizon. Sparnacian, early Ypresian, Eocene.

Type locality. “Les environs de Soissons” [surroundings of Soisson], France.

Remarks. Cossmann (1888: 280) classified the species in the genus *Faunus* Montfort, 1810, Wenz (1929: 2623) affiliated it with *Melanatria* Bowdich, 1822 (both Pachychilidae). Note that both authors gave the name as “*dufresnei*”, which is an incorrect subsequent spelling. Moreover, *Melanatria* Bowdich, 1822 [February] is not an available name because it is a replacement name for the vernacular “Pyrene Lamarck” which was not available in nomenclature before April 1822.

Melanopsis dumbeensis Crosse, 1869

Original source. Crosse 1869a: 70 (Latin) [January].

Type locality. “Dumbea”, New Caledonia.

Remarks. Re-described in French by Crosse (1869b: 281, pl. 8, fig. 4) [October].

† *Melanopsis nodosa* var. *duonodulosa* Ippolito, 1947

Original source. Ippolito 1947: [p. 4 of the Separatum], [unnumbered plate], figs 32–35.

Type horizon. Late Villafranchian, early Pleistocene.

Type locality. “Colle dell’Oro presso Terni”, Italy.

† *Melanopsis bouei* var. *duplicata* Handmann, 1882

Original source. Handmann 1882: 557.

Type horizon. Pannonian, zone D, late Miocene.

Type locality. “Kottingbrunn [...] Ziegelei a”, Austria.

Remarks. The taxon is not included in the Fossilium Catalogus of Wenz (1929).

† *Melanopsis dutemplei* Deshayes, 1862

Original source. Deshayes 1861–1864: 427, pl. 21, fig. 31.

Type horizon. Sparnagian, early Ypresian, Eocene.

Type locality. “Saran, Cramant, Damery”, France.

Remarks. Wenz (1929: 2637) considered the taxon as a junior synonym of *Faunus cerithiformis* (Watelet, 1851) (Pachychilidae).

Melanopsis duveyrieri Bourguignat in Letourneux & Bourguignat, 1887

Original source. Letourneux and Bourguignat 1887: 160.

Type locality. “El-Hammam, près Tozer” [El-Hamma-Djerid near Tozeur], Tunisia.

† *Melanopsis matheroni* var. *ecarinata* Fontannes, 1880

Original source. Fontannes 1879–1882: 176, pl. 10, fig. 3.

Type horizon. Miocene or Pliocene.

Type locality. Rhône Basin (no exact locality given), France.

Remarks. Based on material from the Rhône Basin, not Italy as claimed by Wenz (1929); probably the taxon was mixed up by Italian authors cited by Wenz (1929). Harzhauser et al. (2015: 9) considered the taxon as a junior synonym of *Melanopsis narzolina* d’Archiac in Viquesnel, 1846.

Melanopsis edrissiana Pallary, 1918

Original source. Pallary 1918: 151.

Type locality. “Ras el Mâ et Fès” [Ras el Ma], Morocco.

Melanopsis egregia Bourguignat, 1884

Original source. Bourguignat 1884: 146.

Type locality. “Dans le Jourdain; [...] dans le Belus près de Saint-Jean-d’Acre” [in the Jordan river; in the Na’aman river, near Acre], Israel.

Remarks. Heller et al. (2005: 247) considered the species as a junior synonym of *Melanopsis obliqua* Bourguignat, 1884.

***Melanopsis dufourii* var. *elata* Issel, 1866**

Original source. Issel 1866: 33.

Type locality. “Nella Maremma Toscana in un fumicello d’acqua calda detto Caldana di Ravi, presso Campiglia” [in the Tuscan Maremma, in a stream of warm water called Caldana di Ravi, at Campiglia Marittima], Italy.

***Melanopsis graellsii* var. *elator* Pallary, 1924**

Original source. Pallary 1924: 250.

Type locality. “Gandia (Valencia)”, Spain.

Remarks. Not available from Rossmässler (1854), to whom Pallary referred, because Rossmässler did not use the term “*elator*” to denote a species-group taxon but only cited Férussac’s (1823) description of an unnamed variety of *M. dufourii*. Pallary briefly described the taxon (by indicating its size) and therefore made the name available.

***Melanopsis elegans* Gassies, 1869**

Original source. Gassies 1869: 76.

Type locality. “Noumea”, New Caledonia.

“*Melanella elegans*” mentioned in Bourguignat (1884: 15) [unavailable]

Locality. “De la Save et de la Savina; [...] du Danube, près de Belgrade (Serbie), de Zenica (Bosnie), de la rivière d’Ostaria et d’un ruisseau sur là route de Pregrada, non loin de Krapina-Toeplitz, en Croatie; [...] au pont de la Save, près d’Agram” [in the Sava and Savinja rivers (Croatia, Slovenia); [...] Danube river near Belgrade (Serbia), Zenica (Bosnia and Herzegovina), in the river at Oštarije and a stream at the road to Pregrada, near Krapinske toplice (Croatia); [...] at the Sava bridge near Zagreb].

Remarks. The name “Var. *elegans*” as mentioned by Rossmässler (1839) is a nomen nudum, referring to the personal opinion of Schmidt, with whom Rossmässler collected the material. If available, it would be a junior objective synonym of *M. holandrii legitima* Rossmässler, 1839, under which “*elegans*” was listed as a synonym by Rossmässler.

† *Melanopsis eleis* Oppenheim, 1891

Original source. Oppenheim 1891: 465, pl. 26, fig. 5.

Type horizon. Pliocene.

Type locality. “Bizerè, nördlich von Pyrgos (Elis)” [Bizeré, north of Pyrgos], Greece.

***Melanopsis orontis* var. *elevata* Pallary, 1939 [unavailable]**

Original source. Pallary 1939: 92, pl. 6, fig. 74.

Type locality. Unclear: given as “L’Oronte à Djishr ech Chegour” [Orontes river at Jisr Ash-Shughur] in text but as “L’Oronte à Homs” [Orontes river at Homs (which is far to the south of the former)] in plate captions, Syria.

Remarks. Unavailable according to Art. 13.1, because it lacks a verbal description or definition. Heller et al. (2005: 244) considered it as a junior synonym of *Melanopsis costata* (Olivier, 1804).

† ***Melanopsis impressa* var. *elliptica* Handmann, 1887**

Original source. Handmann 1887: 23, pl. 3, fig. 9.

Type horizon. Pannonian, zone B–D, late Miocene.

Type locality. “Leobersdorf”, Austria.

† ***Melanopsis buccinoidea* var. γ) *antiquua elongata* Férussac, 1822**

Original source. Férussac 1819–1851: *Mélanopsides fossiles*, pl. 1 (1822), figs 5–7.

Type horizon. Eocene.

Type locality. “Des environs d’Epernay. [...] De l’île de Wight” [surroundings of Épernay (France); Isle of Wight (United Kingdom)].

Remarks. Introduced within the var. γ , to which Férussac attributed the term “antiquua”, which was probably not intended as species-group name (see introduction for a detailed discussion of the names introduced by Férussac 1823). Note that for figs 6 and 7 of pl. 7 in Férussac (1823), which is the same as pl. 1 of the “*Mélanopsides fossiles*” in Férussac (1819–1851), Pallary (1916) introduced two new names, i.e., *M. depressa* and *M. sparnacensis*.

***Melanopsis elongata* Gassies, 1874 [invalid]**

Original source. Gassies 1874: 384.

Type locality. “Bourail”, New Caledonia.

Remarks. Junior homonym of *Melanopsis buccinoidea elongata* Férussac, 1822. Pallary (1916: 82) introduced *M. goulvaini* as replacement name for the junior homonym.

† ***Melanopsis narzolina* var. *elongata* Locard, 1878 [invalid]**

Original source. Locard 1878: 58.

Type horizon. Tortonian, late Miocene.

Type locality. “À Tersannes près de Hauterives (Drôme)” [at Tersanne near Hauterives], France.

Remarks. Junior homonym of *Melanopsis buccinoidea elongata* Férussac, 1822. Wenz (1929: 2785) considered the taxon as a junior synonym of *M. narzolina* d’Archiac in Viquesnel, 1846.

† *Melanopsis bouei* var. *elongata* Handmann, 1882 [invalid]

Original source. Handmann 1882: 557.

Type horizon. Pannonian, zone D, late Miocene.

Type locality. “Kottingbrunn [...] Ziegelei a”, Austria.

Remarks. Junior homonym of *Melanopsis buccinoidea elongata* Férussac, 1822. Not included in the Fossilium Catalogus by Wenz (1929).

Melanella letourneuxi var. *elongata* Bourguignat, 1884

Original source. Bourguignat 1884: 26.

Type locality. “Ogulin”, Croatia.

† *Melanopsis (Martinia) vindobonensis* var. *elongata* Handmann, 1887 [invalid]

Original source. Handmann 1887: 28, pl. 6, figs 3–4.

Type horizon. Pannonian, zone B–D, late Miocene.

Type locality. “Leobersdorf”, Austria.

Remarks. Junior homonym of *Melanopsis buccinoidea elongata* Férussac, 1822.

† *Melanopsis citharella* var. *elongata* Locard, 1893 [invalid]

Original source. Locard 1893: 178, pl. 9, fig. 17.

Type horizon. “Helicidenmergel”, late Burdigalian, early Miocene.

Type locality. “Uetken, en Argovie”, Switzerland.

Remarks. Junior homonym of *Melanopsis buccinoidea elongata* Férussac, 1822. Neubauer et al. (2014c: 29) considered this taxon as a junior synonym of *Melanopsis citharella* Merian, 1849.

† *Melanopsis hazayi* var. *elongata* Brusina, 1903 [invalid]

Original source. Brusina 1903: 112.

Type horizon. Late Pleistocene–early Holocene.

Type locality. “Bischofsbad” [Püspökfürdő, Băile 1 Mai, Lake Peșea], Romania.

Remarks. Junior homonym of *Melanopsis buccinoidea elongata* Férussac, 1822. Neubauer et al. (2014d: 125) considered this taxon as a junior synonym of *Microcolpia parreyssii sikorai* (Brusina, 1903).

† *Melanopsis elongata* Doncieux, 1908 [invalid]

Original source. Doncieux 1908: 202, pl. 11, fig. 10.

Type horizon. Middle Lutetian, Eocene.

Type locality. “Au Nord d’Albas”, France.

Remarks. Junior homonym of *Melanopsis buccinoidea elongata* Férussac, 1822. Pallary (1916: 82) introduced *M. sublongata* [sic] as replacement name.

Melanopsis (Mesopotamia) khabourensis var. *elongata* Pallary, 1939 [invalid]

Original source. Pallary 1939: 103, pl. 5, fig. 16.

Type locality. “Ras el ‘Ain du Khabour” [Chabur river near Ra’s al ‘Ayn], Syria.

Remarks. Junior homonym of *Melanopsis buccinoidea elongata* Férussac, 1822. Heller et al. (2005: 239) tentatively considered the variety as a junior synonym of *Melanopsis khabourensis* Pallary, 1939.

“*Melanopsis praemorsa* f. *elongata*” mentioned in Pérès (1939) [unavailable]

Locality. Collection stations 80 [tributary rivulet to the river Tensift] and 144 [Oued Sidi Raho, approximately 6 km upstream of its confluence with the Bou Regreg] (Pérès 1939: 139), Morocco.

Remarks. First of all, the name as given by Pérès (1939) is a nomen nudum – Pérès apparently considered the expression “*elongata*” self-explanatory and did not describe it. Moreover, he obviously used the name not as separate taxon but rather as descriptive term to fit existing species into his morphological concept. He even indicated *M. olivieri* Bourguignat, 1884 as its “type”.

† *Melanopsis (Canthidomus) defensa elongata* Gillet & Marinescu, 1971 [invalid]

Original source. Gillet and Marinescu 1971: 55, pl. 23, figs 38–48.

Type horizon. Transdanubian, Pannonian, late Miocene.

Type locality. “Rădmănești”, Romania.

Remarks. Junior homonym of *Melanopsis buccinoidea elongata* Férussac, 1822. Neubauer et al. (2014c: 31) considered this variety a junior synonym of *Melanopsis defensa* Fuchs, 1870.

† *Melanopsis elophila* Pallary, 1925

Original source. Pallary 1925: 257.

Type horizon. Messinian, late Miocene.

Type locality. “Della valle della Sterza” (Capellini 1880: 420) [La Sterza near Lajatico], Italy.

Remarks. Introduced for *M. buccinoidea* sensu Capellini, 1880, non Férussac, 1823.

“*Melanopsis elysaea*” mentioned in Brot (1874–1879: 419) [unavailable]

Locality. Not indicated.

Remarks. Nomen nudum, “in schedis” name from Tarnier listed in the synonymy list of *M. buccinoidea* (Olivier, 1801) by Brot (1879).

Melanopsis foleyi var. *emaciata* Pallary, 1928 [unresolved]

Original source. Pallary 1928a: 271.

Type locality. “Aïn Mélias, près de Figuig” [Ain Melias near Figuig], Algeria.

Remarks. Homonym of the simultaneously published name *Melanopsis letourneuxi emaciata* Pallary, 1928 (see Note 1). This case requires the action of a First Reviser.

Melanopsis letourneuxi var. *emaciata* Pallary, 1928 [unresolved]

Original source. Pallary 1928a: 273, pl. 5, fig. 7.

Type locality. “Berguent” [Aïn Beni Mathar], Morocco.

Remarks. Homonym of the simultaneously published name *Melanopsis foleyi emaciata* Pallary, 1928 (see Note 1). This case requires the action of a First Reviser.

Melanopsis monteli var. *emaciata* Pallary, 1936 [invalid]

Original source. Pallary 1936: 61.

Type locality. Not explicitly stated but probably the same as for the species (“L’Oued Sous, au pont des Aït Melloul, sur la route d’Agadir à Tiznit, à 14 kil. S. O. d’Agadir” [in the Oued Sous, at the bridge of Ait Melloul, at the road from Agadir to Tiznit, 14 km southwest of Agadir], Morocco).

Remarks. Junior homonym of *Melanopsis letourneuxi emaciata* Pallary, 1928 and *M. foleyi emaciata* Pallary, 1928 (simultaneously published; no priority fixed yet; see Note 1).

***Melanopsis cerithiopsis* var. *emaciata* Pallary, 1939 [invalid]**

Original source. Pallary 1939: 96, pl. 6, figs 40–43.

Type locality. “À Jéricho, dans ‘Ain Solthan” [at Jericho, in the spring ‘Ayn as Sulṭān], Palestine.

Remarks. Junior homonym of *Melanopsis letourneuxi emaciata* Pallary, 1928 and *M. foleyi emaciata* Pallary, 1928 (simultaneously published; no priority fixed yet; see Note 1).

† ***Melanopsis emerici* d’Orbigny, 1850**

Original source. d’Orbigny 1850: 309.

Type horizon. Early Rupelian, Oligocene.

Type locality. “Levit, près de Castellanne” [Vit-de-Castellane near Castellane], France.

† ***Melanopsis geniculata* f. *enodata* Brusina, 1897**

Original source. Brusina 1897: 11.

Type horizon. Early Langhian, middle Miocene.

Type locality. “Sinj (Župića potok)”, Croatia.

Types. The syntypes are stored in the Croatian Natural History Museum, Zagreb; no number indicated (Milan et al. 1974: 91).

Remarks. Wenz (1929: 2734) considered the taxon as a junior synonym of *Melanopsis geniculata* Brusina, 1874, while Neubauer et al. (2016b: 23) treated it as a distinct species. The name “*enodota*” as mentioned in Wenz (1929: 2734) is an incorrect subsequent spelling.

“*Melania enodes* Ziegler” mentioned in Reeve (1860) [unavailable]

Locality. Not indicated.

Remarks. Perhaps a manuscript or “in schedis” name from Ziegler. Reeve (1860) considered this taxon as a junior synonym of *Melania hollandrii* [sic].

† ***Melanopsis entzi* Brusina, 1902**

Original source. Brusina 1902: pl. 6, figs 34–37.

Type horizon. Transdanubian, Pannonian, late Miocene.

Type locality. “Tihany; Kenese” [Tihany; Balatonkenese], Hungary.

Types. Milan et al. (1974: 90) defined a “lectosyntypus”, a term not accepted by the Code, based on the specimen from Tihany illustrated by Brusina (1902: pl. 6, figs

34–36). This fixation is insufficient for a valid lectotype designation (Art. 74). The specimen is stored in the Croatian Natural History Museum, Zagreb, coll. no. 2515-161/1-2.

Remarks. Bartha (1955: 294) considered this taxon as a junior synonym of *Melanopsis fuchsi* Handmann, 1882.

† *Melanopsis eocenica* Pallary, 1916

Original source. Pallary 1916: 79.

Type horizon. Sparnacian, early Ypresian, Eocene.

Type locality. “Mont Bernon bei Epernay” (only this locality is the type locality as Pallary clearly referred to the specimen illustrated by Sandberger 1870–1875: pl. 9, fig. 5), France.

Remarks. Introduced for *M. buccinoidea* sensu Sandberger, 1870. Wenz (1929: 2660) considered that record a junior synonym of *M. antidiluviana* (Poiret, 1801). The name “*eocaenica*” as mentioned in Wenz (1929: 2662) is an incorrect subsequent spelling.

Melanopsis episema Bourguignat, 1884

Original source. Bourguignat 1884: 88.

Type locality. “Près de Biskra, dans un ruisseau d’eau chaude à Ouargla, dans un puits artésien de Mazer au Ziban, ainsi qu’aux environs de Boghar et dans les rivières du djebel Takreda entre Mogador et Maroc [sic]” [near Biskra, in a stream of warm water at Ouargla, in an artesian well of Mazer, as well as around Boghar (Algeria) and in rivers of Jebel Takreda (?) between Essaouira and Morocco (?)].

Melanopsis eremita Tristram, 1865

Original source. Tristram 1865: 542.

Type locality. “Wady Um Bagkek, between Sebbeh and Jebel Usdum, at the south-west corner of the Dead Sea”, Israel.

Remarks. Heller et al. (2005: 236) considered the species as a junior synonym of *Melanopsis ammonis* Tristram, 1865.

Melanopsis laevigata var. *erosa* Roth, 1839

Original source. Roth 1839: 24.

Type locality. “In Peloponneso”, Greece.

***Melanopsis esperi* Férussac, 1823**

Original source. Férussac 1823: 160.

Type locality. “La Laybach” [Ljubljana river], Slovenia.

Remarks. Type species of *Esperiana* Bourguignat, 1877.

† ***Melanopsis esperioides* Stefanescu, 1896**

Original source. Stefanescu 1896: 128, pl. 11, figs 6–11.

Type horizon. Pliocene.

Type locality. “À Bucovatz, dans la vallée de Jiu, et à Milcov, près de Slatina, dans la vallée de l’Oltu” [at Bucovăț, in the valley of the river Jiu, and at Milcov, near Slatina, in the valley of the river Olt], Romania.

Remarks. The name “*esperiodea*” as mentioned in Huică (1977: 41) is an incorrect subsequent spelling.

***Melanopsis lorcani* var. *etrusca* Westerlund, 1886**

Original source. Westerlund 1886: 128.

Type locality. “Italien, Balearen [Westerlund]; aus Toscana als auch aus Jumilla (Spanien) und Merknies (Marocco) [Brot]” [Italy, Balearic Islands (Spain); Tuscany (Italy), Jumilla (prov. Murcia, Spain), Merknies (Morocco)].

Remarks. The name was first mentioned by Brot (1862: 63) as “*etrusca* Villa (v. min)” from the locality “Toscane”, without description or bibliographical reference. Brot listed it under the name “*Melanopsis Dufourii* Fér.”, for which a bibliographical reference (to “Rossm. Icon. 835–839”) and locality “Espagne” was given. Since *etrusca* was neither described by Brot nor by Rossmässler, the name *etrusca* is not available from this case (Art. 12.1).

Brot (1879: 434) listed the name (“*Melanopsis etrusca* Villa MSS.”) as synonym of *Melanopsis Dufourii* var. β , which he introduced there with a short Latin description. On p. 435 Brot even associated *etrusca* with an illustration of a specimen from Toscana (pl. 47, fig. 3). However, the name was explicitly referred to as synonym of the (unnamed) variety β , which is not a valid name, and thus the requirements of Art. 11.6 (and therefore 12.1) are not met (see Note 2).

Westerlund (1886: 128) used the name as “[*Melanopsis lorcani*] var. *etrusca* Villa ap. Brot (Cat. syst. Melan. 1862)” and provided a description and thus made it available as species-group taxon. Since he referred to Brot’s catalogue, the type locality comprises both his and Brot’s records. The authority of *Melanopsis etrusca* remains with Westerlund. Art. 11.6.1 does not apply here as Brot (1879) had listed the name as synonym of “Var. β ”, which is not a valid name (see Note 2).

† *Melanopsis eulimopsis* Brusina, 1902

Original source. Brusina 1902: pl. 5, figs 42–44.

Type horizon. Middle Pannonian, late Miocene.

Type locality. “Kúp”, Hungary.

Types. The illustrated syntypes are stored in the Croatian Natural History Museum, Zagreb, coll. no. 2494-140/1-3 (Milan et al. 1974: 90).

Remarks. Considered a junior synonym of *M. pygmaea mucronata* Handmann, 1887 by Papp (1953b: 150).

Melanopsis eumorphia Bourguignat, 1884

Original source. Bourguignat 1884: 146.

Type locality. “Dans le Jourdain, à 4 kilomètres au-dessus de la Mer Morte” [in the Jordan river, 4 km above the Dead Sea], Palestine/Jordan.

Remarks. Apparently by mistake, Heller et al. (2005: 245, 247) listed the species in synonymy of *Melanopsis lampra* Bourguignat, 1884 as well as of *Melanopsis obliqua* Bourguignat, 1884, both of which they ranked as subspecies of *M. costata* (Olivier, 1804).

† *Melanopsis (Lyrcea) euphrosinae* Cobălcescu, 1883

Original source. Cobălcescu 1883: 124, pl. 9, figs 11a–b.

Type horizon. Dacian, Pliocene.

Type locality. “Beceni”, Romania.

† *Melanopsis eurystoma* Neumayr in Neumayr & Paul, 1875

Original source. Neumayr and Paul 1875: 49, pl. 8, fig. 30.

Type horizon. Cernikian, Pliocene.

Type locality. “Čaplthal” [Čapljia trench near Slavonski Brod], Croatia.

Melanopsis excoriata Pallary, 1920

Original source. Pallary 1920a: 31.

Type locality. “Aït Brahim; Tazouta; El Menzel; [...] l’Oued Raha, à Agourai” [Aït Brahim, Tazouta, El Menzel, Oued Rha at Agourai], Morocco.

***Melanopsis chehirensis* var. *exilis* Pallary, 1939**

Original source. Pallary 1939: 95, pl. 6, figs 51, 52, 63.

Type locality. “Yeni Chehir” [Yenişehir], Turkey.

***Melanella eximia* Bourguignat, 1884**

Original source. Bourguignat 1884: 13.

Type locality. “Rivière d’Ostaria, entre Plaski et Ogulin, en Croatie; la Save, à Sissek, en Slavonie; la Narenta, en Dalmatie” [river at Oštarije, between Plaški and Ogulin; Sava river at Sisak in Slavonia; in the Narenta river in Dalmatia], Croatia.

Remarks. Note that Bourguignat denoted the authority as “Bourguignat, 1880”.

***Melanopsis eximia* Pallary, 1928**

Original source. Pallary 1928a: 260, pl. 5, figs 16–18.

Type locality. “Agourai, à 29 kil. sud de Fès” [Agourai, claimed to be 29 km south of Fès; judging from the map, Pallary obviously mixed up Fès with Meknès in this case], Morocco.

***Melanopsis magnifica* var. *expansa* Pallary, 1920**

Original source. Pallary 1920a: 33.

Type locality. “Agourai” [Agourai, south of Meknes], Morocco.

† *Melanopsis (Lyrcea) delessei* var. *expansa* Magrograssi, 1928 [invalid]

Original source. Magrograssi 1928: 259, pl. 6, fig. 15.

Type horizon. Plio-Pleistocene.

Type locality. “Coo: frequente nella zona centrale fossilifera” [Kos island: common in the central fossiliferous zone], Greece.

Remarks. Junior homonym of *Melanopsis magnifica expansa* Pallary, 1920 (see Note 1).

† *Melanopsis (Martinia) martiniana* var. *extensa* Handmann, 1887

Original source. Handmann 1887: 26, pl. 5, figs 3–4.

Type horizon. Pannonian, zone B–D, late Miocene.

Type locality. “Leobersdorf”, Austria.

Remarks. Wenz (1929: 2720) considered the taxon as a junior synonym of *Melanopsis fossilis* (Gmelin, 1791).

† *Melanopsis faberi* Brusina, 1884

Original source. Brusina 1884a: 167, pl. 29, fig. 1.

Type horizon. Portaferrian (Pannonian Basin), late Miocene–Pliocene.

Type locality. “Okrugljak” [near Zagreb], Croatia.

Types. Milan et al. (1974: 90) indicated a holotype, but it is uncertain whether the specimen was the only one Brusina had at hand (holotype by monotypy, Art. 73.1.2). The specimen is stored in the Croatian Natural History Museum, Zagreb, coll. no. 2966-612.

Melanella fagotiana Bourguignat, 1884

Original source. Bourguignat 1884: 18.

Type locality. “Dans la rivière de la Krapina, à Sused, en Croatie” [in the river Krapina at Podsused, in Zagreb], Croatia.

Remarks. Note that Bourguignat denoted the authority as “Bourguignat, 1879”.

† *Melanopsis fallax* Pantanelli, 1886

Original source. Pantanelli 1886a: 64 or 1886b: 67, pl. 3, figs 12–13 (precedence not established).

Type horizon. Messinian, late Miocene.

Type locality. “Sterza” (Capellini 1873: 550) [near Lajatico], Italy.

Remarks. Introduced for the record of *M. acicularis* sensu Capellini, 1873, non Férussac, 1823.

Melanopsis fasciata Gassies, 1874

Original source. Gassies 1874: 381.

Type locality. “Nékété” [Nakéty], New Caledonia.

Remarks. Brot (1879: 444) considered the taxon as a junior synonym of *Melanopsis frustulum* Morelet, 1857.

† *Melanopsis fasciata* Handmann, 1882 [invalid]

Original source. Handmann 1882: 559.

Type horizon. Pannonian, zone D, late Miocene.

Type locality. “Kottingbrunn [...] Ziegelei a”, Austria.

Remarks. Junior homonym of *Melanopsis fasciata* Gassies, 1874. Pallary (1916: 86) introduced *M. vittata* as replacement name. The taxon was considered as a junior synonym of *M. haueri* by Wenz (1929: 2741).

Melanopsis doriae var. *fasciata* Biggs, 1937 [invalid]

Original source. Biggs 1937: 248.

Type locality. “From a ‘qanat’ [= typical Iranian channel] at Jelalabad, near Kerman”, Iran.

Remarks. Junior homonym of *Melanopsis fasciata* Gassies, 1874.

“*Melania fasciata*” mentioned in Brusina (1866: 106) and Brusina (1867: 86) [unavailable]

Locality. “Cetina”, Croatia.

Remarks. Nomen nudum, “in schedis” name in collection of Kucik (also read as “Kutschig”), occasionally listed with authority “Stentz”. If available, the name would be junior homonym of *Melania fasciata* Sowerby, 1818 described from the Paleogene of the United Kingdom.

Melanopsis maroccana var. *fasciolata* Bourguignat, 1864

Original source. Bourguignat 1864: 260.

Type locality. “Ruisseau de Ngouca” [N’Goussa], Algeria.

Melanopsis fasensis Pallary, 1920

Original source. Pallary 1920a: 32.

Type locality. “Taforalt. Aïn Sfa. Fès (Ras el Mâ). [...] Dar Batha, jardin public de bou Djeloud, séguias de dar el Maghzen, séguias en dehors de bab sidi bou Jida” [Taforhalt; Aïn Sfa; Fes; Ras el Ma; Dar Batha, park of Bou Jeloud, irrigation channels of Dar El Makhzen, irrigation channels outside Bab Sidi Boujida (all latter localities are within Fes)], Morocco.

***Melanopsis faseolaria* Brot, 1868**

Original source. Brot 1868: 58, pl. 2, fig. 10.

Type locality. “Persepolis”, Iran.

Remarks. Brot attributed the authority to Parreyss, based on an “in schedis” determination by that author. However, from the following text it becomes clear that Brot is the author of the species.

† *Melanopsis fateljensis* Neubauer, Mandic & Harzhauser, 2014

Original source. Neubauer et al. 2014b: 52, figs 2–11.

Type horizon. Langhian, middle Miocene.

Type locality. “NW slope of Fatelj hill near Kupres”, Bosnia and Herzegovina.

Types. Geological-Paleontological Department, Natural History Museum Vienna, Austria, coll. no. 2011/0138/0107a.

Remarks. Introduced for formerly misidentified *Melanopsis mojsisovicsi* Neumayr, 1880 in Neubauer et al. (2013: 137, figs 5E–F, I–K).

***Melanopsis feliciani* Bourguignat, 1884**

Original source. Bourguignat 1884: 145.

Type locality. “Dans le Jourdain, non loin de son embouchure dans la Mer Morte” [in the Jordan river, near its mouth at the Dead Sea], Palestine/Jordan.

***Melanopsis ferussaci* Roth, 1839**

Original source. Roth 1839: 24, pl. 2, fig. 10.

Type locality. “Smyrna” [Izmir], Turkey.

Remarks. Heller et al. (2005: 232) considered the species as a junior synonym of *Melanopsis buccinoidea* (Olivier, 1801).

***Melanopsis excoriata* var. *festiva* Pallary, 1920**

Original source. Pallary 1920c: 144, pl. 4, figs 21–22.

Type locality. “Aït Brahim”, Morocco.

***Melanopsis foleyi* var. *festiva* Pallary, 1928 [invalid]**

Original source. Pallary 1928a: 271.

Type locality. “Aïn Mélias, près de Figuig” [Ain Melias near Figuig], Algeria.

Remarks. Junior homonym of *Melanopsis excoriata festiva* Pallary, 1920 (see Note 1).

† ***Melanopsis filifera* Neumayr, 1880**

Original source. Neumayr 1880c: 479, pl. 7, figs 6–7.

Type horizon. Early Langhian, middle Miocene.

Type locality. “Westlich von Drvar” [west of Drvar], Bosnia and Herzegovina.

† ***Melanopsis flammulata* De Stefani, 1877**

Original source. De Stefani 1877: 306, pl. 18, fig. 7.

Type horizon. Villafranchian, Plio-Pleistocene.

Type locality. “Monte Albuccio, [...] Valli della Pescaia, e del Riluogo [...] presso Siena; Castellacela presso Massa Marittima [...]; Colline di Piedimonte presso Terni nell’Umbria [...], tra S. Gemine e Carsoli sulla strada di Narni, e fra Otricoli e le Vigne sulla strada da Roma a Foligno?” [Monte Albuccio, Valli della Pescaia, Riluogo near Siena, Castellacela near Massa Marittima, Piedimonte hills near Terni in Umbria, between San Gemini and Carsoli on the road to Narni, and between Otricoli and Le Vigne], Italy.

Remarks. Wenz (1929: 2663) considered the taxon as a junior synonym of “*M. antiqua* Férussac”, which is not an available name.

“*Melania holandri* var. *flava* Zelebor” mentioned in Brot (1874–1879: 12) [unavailable]

Locality. Not indicated.

Remarks. Nomen nudum, “in schedis” name from Zelebor listed in synonymy of “*Melania holandri*” [sic] by Brot (1874). If available, it would be junior homonym of *Melania flava* Deshayes in Deshayes & Jullien, 1876 described from Cambodia.

***Melanopsis foleyi* Pallary, 1920**

Original source. Pallary 1920a: 29.

Type locality. “Aïn Mélias, près de Figuig” [Ain Melias near Figuig], Algeria.

† *Melanopsis foliacea* Gümbel, 1861

Original source. Gümbel 1861: 753.

Type horizon. Chattian, Oligocene.

Type locality. “Wester-Buchberg, Loherflötz, Aubach, Rohmbach, Sulzgrabenflötz, Schlierachthal, Neumühle, bei Rimselrain, Pensberg, im Höllgraben, im Buchbergflötz, bei Schmerold, am hohen Peissenberge” [all localities near Miesbach, southern Bavaria], Germany.

† *Melanopsis (Canthidomus) fontanensis* Villatte in Tambareau & Villatte, 1966

Original source. Tambareau and Villatte 1966: 512, pl. 1, figs 4–18.

Type horizon. Sparnacian, early Ypresian, Eocene.

Type locality. “Fontané (Ariège)”, France.

Types. Laboratoire de Géologie de la Faculté des Sciences de Toulouse; no number indicated.

† *Melanopsis (Martinia) capulus* var. *fornicata* Handmann, 1887

Original source. Handmann 1887: 21, pl. 2, fig. 20.

Type horizon. Pannonian, zone B–D, late Miocene.

Type locality. “Leobersdorf”, Austria.

Melanopsis costata var. *forticostata* Paetel, 1888

Original source. Paetel 1888: 400.

Type locality. “Lake Huleh [Palestine]” (Nevill 1884: 212), Palestine.

Remarks. Originally introduced as infrasubspecific taxon (“subvariety”) by Nevill (1884), but made available by Paetel (1888) who treated it as variety (Art. 45.5.1). Paetel clearly referred to the description of Nevill.

† *Melania fossariformis* Tournouër, 1879

Original source. Tournouër 1879: 261.

Type horizon. Cernikian, Pliocene.

Type locality. “Nisipulu, Rumaniae” [Nisip], Romania.

Remarks. Considered to belong in the genus *Amphimelania* by Cossmann (1909: 126) and Wenz (1929: 2873).

† *Buccinum fossile* Gmelin, 1791

Original source. Gmelin 1791: 3485.

Type horizon. Sarmatian (sensu stricto), middle Miocene.

Type locality. “Zwischen Oedinburg and Ruß” (Martini 1777: 204) [between Sopron (Hungary) and Rust (Austria)], Hungary/Austria.

Types. Papp (1953b: 133) designated the specimen illustrated by Martini (1777: pl. 94, figs 913) as lectotype (see also Art. 74.5). Since Gmelin explicitly referred to Martini’s work when describing the species, this designation is valid.

Remarks. The taxon in question was first mentioned in the Catalogus by Walch (1768: 121, pl. C. II.*, figs 1–5), yet without name. Martini (1777: 203, pl. 94, figs 912–914) described and illustrated the species as “*Pyrum fossile monstrosus* f. *amorphon*” and referred to Walch’s work. Species-group names introduced in volumes 1–11 of Martini and Chemnitz’ Neues Systematischer Conchylien Cabinet (1769–1795) are, however, unavailable according to Opinion 184 (ICZN 1944). Gmelin (1791: 3485) referred to Martini’s work, described the species and was the first to assign an available name to the taxon: *Buccinum fossile*.

Férussac (1823), who was the first to combine the species with *Melanopsis*, listed the records by Walch, Martini and Gmelin but apparently did not consider “*fossile*” a valid name and introduced “*Melanopsis martiniana*” as new name. That name is permanently invalid as it is an objective synonym (see also Pallary 1916: 81).

The last summary of the status of *M. fossilis* was provided by Fischer (1996a), but his conclusions were unfortunately based on misinterpretations and nomenclatural errors. He was unaware of the fact that “*Melanopsis martiniana*” is an objective synonym of *M. fossilis*. Moreover, he considered the outcrops near Siegendorf as “locus typicus restrictus”, a term not existing in nomenclature.

Melanopsis foucauldiana Pallary, 1920

Original source. Pallary 1920a: 30.

Type locality. “Ouggarta, à 60 kil. au sud de Beni Abbès, Sahara occidental” [Ouggarta, prov. Béchar], Algeria.

Melanopsis foureaui Pallary, 1920

Original source. Pallary 1920a: 29.

Type locality. “Tabebala, dans le Sahara occidental” [Tabebala (?), in the western Sahara], Algeria.

***Melanopsis fragilis* Gassies, 1874**

Original source. Gassies 1874: 382.

Type locality. “Ouagap” [Wagap], New Caledonia.

“*Melania fragilis* Schmidt” mentioned in Brusina (1867: 86) [unavailable]

Locality. Not indicated.

Remarks. Nomen nudum, “in schedis” name in collection of Kucik (also read as “Kutschig”). If available, the name would be junior homonym of *Melania fragilis* Lamarck, 1804 from the Paleogene (Eocene?) of the Paris Basin.

† *Melanopsis franciscae* Brusina, 1903

Original source. Brusina 1903: 113.

Type horizon. Late Pleistocene–Holocene.

Type locality. “Bischofsbad” [Püspökfürdő, Băile 1 Mai, Lake Peșea], Romania.

Remarks. Neubauer et al. (2014d: 125) considered this taxon as a junior synonym of *Microcolpia parreyssii sikorai* (Brusina, 1903).

† *Melanopsis (Canthidomus) freybergi* Kühn, 1951

Original source. Kühn 1951: 186, pl. 17, figs 4–5.

Type horizon. Mammal zone MN 10, late Miocene.

Type locality. “Nordwestlich Nea Liossia” [northwest of Ilion], Greece.

Types. Geological-Paleontological Department, Natural History Museum Vienna, Austria, coll. no. 1949/0004/0006.

† *Amphimelania frici* Brusina, 1897

Original source. Brusina 1897: 6.

Type horizon. Late Cernikian, late Pliocene–early Pleistocene.

Type locality. “Kutina (šuma Mišinka)” [Kutina, Mišinka forest], Croatia.

Types. Milan et al. (1974: 57) stated that Brusina (1897) had indicated one of the specimens illustrated by him (pl. 3, figs 27–28) as type. This can be accepted as valid lectotype designation in the sense of Art 74.5. The specimen is stored in the Croatian Natural History Museum, Zagreb, coll. no. 2964-610/1.

† *Melanopsis friedeli* Brusina, 1885

Original source. Brusina 1885: 160.

Type horizon. Cernikian, Pliocene.

Type locality. “Kravarsko und Podvornica”, Croatia.

Types. Milan et al. (1974: 91) defined a neotype based on the specimens illustrated by Brusina (1897: 8, pl. 6, figs 5–8) because the specimen from 1885 was apparently lost. The type fixation is insufficient with respect to Art. 75.3 of the Code. The specimen is stored in the Croatian Natural History Museum, Zagreb, coll. no. 2998-644.

† *Melanopsis fritzei* Thomä, 1845

Original source. Thomä 1845: 158, pl. 2, figs 7a–b.

Type horizon. Mammal zone MN 2, early Miocene.

Type locality. “Mühltal bei Wiesbaden”, Germany.

Remarks. The name “*Fritzii*” as mentioned in Chenu (1859: 297) is an incorrect subsequent spelling.

Melanopsis frustulum Morelet, 1857

Original source. Morelet 1857: 33.

Type locality. “[Ad Sanctam-Mariam de Balade]” [Balade], New Caledonia.

† *Melanopsis fuchsi* Handmann, 1882

Original source. Handmann 1882: 556.

Type horizon. Pannonian, zone D, late Miocene.

Type locality. “Kottingbrunn [...] Ziegelei a”, Austria.

Remarks. Wenz (1929: 2813) synonymized the species with *M. pygmaea* Hörnes, 1856. Jekelius (1944: 75) invalidly synonymized it with *M. handmanni*, which is the objective junior synonym of *M. fuchsi* Handmann, 1882.

† *Melanopsis fuchsi* Brusina, 1884 [invalid]

Original source. Brusina 1884a: 168.

Type horizon. Transdanubian, Pannonian, late Miocene.

Type locality. “Radmanest” (Fuchs 1870: 353) [Rădmănești], Romania.

Types. The syntypes are stored in the Croatian Natural History Museum, Zagreb; no number indicated (Milan et al. 1974: 91).

Remarks. Introduced for *M. costata* sensu Fuchs, 1870, non Olivier, 1804. Junior homonym of *Melanopsis fuchsi* Handmann, 1882 (see *M. confusa* Strausz, 1941, *M. handmanni* Brusina, 1892 and *M. hungarica* Pallary, 1916, as well as Neubauer et al. 2014a: 461 for details).

† *Amphimelania fuchsi* Wenz, 1928

Original source. Wenz 1928a: 119.

Type horizon. Late Pliocene to early Pleistocene.

Type locality. “Kalamaki” [near Corinth], Greece.

Remarks. Replacement name for *Paludina* (*Vivipara*) *ornata* Fuchs, 1877, non *Vivipara ornata* Neumayr, 1875, which was considered a Melanopsidae by Wenz (1928a).

Melanopsis fulgurans Gassies, 1858

Original source. Gassies 1858: 371.

Type locality. “Des petits cours d’eau à Balade; [...] à l’île des Pins” [in small streams near Balade; L’Île des Pins], New Caledonia.

Remarks. Brot (1879: 444) considered the taxon as a junior synonym of *Melanopsis frustulum* Morelet, 1857.

“*Melanopsis fulminans* Born.” mentioned in De Cristofori & Jan (1832: Conchyliia terrestria et fluviatilia, p. 7) [unavailable]

Locality. “Austr.” [Australia].

Remarks. Nomen nudum, found only in the species list of De Cristofori and Jan (1832).

Melanopsis fulminata Brot, 1879

Original source. Brot 1874–1879: 458, pl. 49, fig. 10.

Type locality. “Ouagap” [Wagap], New Caledonia.

† *Melanopsis costata* var. *funiculata* P. Fischer in Gaudry, 1867

Original source. Gaudry 1862–1867: 446, pl. 62, figs 13–15.

Type horizon. Pliocene.

Type locality. “Mégare” (p. 444), Greece.

Remarks. The taxon is not included in the Fossilium Catalogus of Wenz (1929).

***Melanopsis fusca* Gassies, 1870**

Original source. Gassies 1870: 147.

Type locality. “Prope Pouebo” [near Pouébo], New Caledonia.

† ***Melanopsis fusiformis* Sowerby, 1822**

Original source. Sowerby 1821–1823: 35, pl. 332, figs 1–7.

Type horizon. Eocene.

Type locality. “Isle of Wight; [...] New Charlton; [...] Woolwich; [...] New Cross near Deptford; [...] Hordwell”, United Kingdom.

† ***Melanopsis buccinoidea* var. *fusiformis* Grateloup, 1838 [invalid]**

Original source. Grateloup 1838: 146.

Type horizon. Burdigalian, early Miocene.

Type locality. “Dax. [...] Mandillot”, France.

Remarks. Based on some of the illustrated syntypes of “*M. antiqua* Férussac, 1823” (actually, Grateloup referred to the illustrations of *M. buccinoidea* in Férussac 1823; “*M. antiqua* Férussac” is not an available name). Junior homonym of *Melanopsis fusiformis* Sowerby, 1822.

***Melanopsis fusiformis* Gassies, 1870 [invalid]**

Original source. Gassies 1870: 148.

Type locality. “Prope Kanala; insula Ouen” [near Canala; Île Ouen], New Caledonia.

Remarks. Junior homonym of *Melanopsis fusiformis* Sowerby, 1822. Gassies (1880) introduced *M. rossiteri* as replacement name. Unaware of this, Pallary (1916) introduced *M. cookiana* as replacement name, which is its junior objective synonym.

† ***Melanopsis fusiformis* Handmann, 1882 [invalid]**

Original source. Handmann 1882: 560.

Type horizon. Pannonian, zone D, late Miocene.

Type locality. “Kottingbrunn [...] Ziegelei a”, Austria.

Remarks. Junior homonym of *Melanopsis fusiformis* Sowerby, 1822. Pallary (1916: 82) introduced *M. angusta* as replacement name. Wenz (1929: 2742) considered the taxon as a junior synonym of *M. haueri* Handmann, 1882.

† *Melanopsis (Canthidomus) proteus* var. *fusiformis* Magrograssi, 1928 [invalid]

Original source. Magrograssi 1928: 260, pl. 6, fig. 17.

Type horizon. Plio-Pleistocene.

Type locality. “Coo: V. Tuvachiu, fra Antimachia e Pili” [Kos island: Tuvachia valley (?), between Antimácheia and Pýli], Greece.

Remarks. Junior homonym of *Melanopsis fusiformis* Sowerby, 1822.

† *Melanopsis praemorsa* var. *fusulatina* Sacco, 1895

Original source. Sacco 1895: 9, pl. 1, fig. 13.

Type horizon. Messinian, late Miocene.

Type locality. “Priosa presso Narzole, Castelletto d’Orba, S. Marzano Oliveto”, Italy.

† *Melanopsis gaasensis* Vergnau-Saubade, 1968

Original source. Vergnau-Saubade 1968: 201, [unnumbered plate], fig. 10.

Type horizon. Oligocene.

Type locality. “Gaas - Lesbarritz”, France.

Remarks. First mentioned in an unpublished thesis (Vergnau-Saubade 1966).

† *Melania gaji* Brusina, 1878

Original source. Brusina 1878: 347.

Type horizon. Middle–late Cernikian, late Pliocene–early Pleistocene.

Type locality. “Sibinj”, Croatia.

Types. Milan et al. (1974: 57) indicated a holotype, but it is uncertain whether the specimen is part of the original type series and whether it was the only one Brusina had at hand (holotype by monotypy, Art. 73.1.2). The specimen is stored in the Croatian Natural History Museum, Zagreb, coll. no. 2963-609.

Remarks. Considered to belong in the genus *Amphimelania* by Brusina (1897: 6), Cossmann (1909: 126) and Wenz (1929: 2874). The names “*caji*” as mentioned in Cossmann (1909: 126) and “*gayi*” as given by Bandel (2000: 140) are incorrect subsequent spellings.

***Fagotia gallandi* Bourguignat, 1884**

Original source. Bourguignat 1884: 41.

Type locality. “Affluents du lac Sabandja, près d’Ismidt” [tributaries to the Lake Sapanca near İzmit], Turkey.

Remarks. Appeared first as a nomen nudum in Locard (1883a).

***Microcolpia gallandi* Bourguignat, 1884**

Original source. Bourguignat 1884: 60.

Type locality. “Constantinople [...], a été trouvée dans la rivière d’Ismidt (Anatolie)” [Istanbul, found in a river at İzmit], Turkey.

Remarks. Starobogatov et al. (1992: 68) considered the species as a junior synonym of *Microcolpia coutagniana* (Pfeiffer, 1828).

† *Melanopsis galloprovincialis* Matheron, 1842

Original source. Matheron 1842: 291, pl. 37, figs 1–6.

Type horizon. Campanian–Maastrichtian, Cretaceous.

Type locality. “Les Martigues, les Pennes, Simiane, Gardanne, Fuveau, Peynier, Trets”, France.

† *Melanopsis garumnica* Munier-Chalmas, 1884

Original source. Munier-Chalmas 1884: 330, pl. 7, figs 21–22.

Type horizon. Maastrichtian, Cretaceous.

Type locality. “D’Auzas”, France.

Remarks. Munier-Chalmas (1884) indicated the authority with “Mun. Ch., 1870”, which is probably due to the fact that issues 1–3 of vol. 1 of *Annales de Malacologie*, in which the original publication appeared, were published in 1870 (issue 4, however, in 1884).

***Melanopsis gassiesiana* Crosse, 1867**

Original source. Crosse 1867: 435, pl. 12, fig. 7.

Type locality. “Wagap”, New Caledonia.

† *Melanopsis gearyae* Neubauer, Harzhauser, Georgopoulou, Mandic & Kroh, 2014

Original source. Neubauer et al. 2014a: 456.

Type horizon. Early Pleistocene (?).

Type locality. “Weganrisse und Regenrisse 3 Km N Antirrion in Akarnanien” (Symeonidis et al. 1986: 342) [sections 3 km N of Antirrio, Aetolia-Acarnania], Greece.

Types. Senckenberg Forschungsinstitut und Naturmuseum Frankfurt; no number indicated.

Remarks. Replacement name for *M. posterior* Schütt, 1986, non Papp, 1953.

† *Melanopsis geniculata* Brusina, 1874

Original source. Brusina 1874: 40, pl. 1, figs 9–10.

Type horizon. Langhian, middle Miocene.

Type locality. “Miočič”, Croatia.

Types. Milan et al. (1974: 91) indicated a holotype, but it is uncertain whether the specimen was the only one Brusina had at hand (holotype by monotypy, Art. 73.1.2). The specimen is stored in the Croatian Natural History Museum, Zagreb, coll. no. 3207-853.

Melanopsis germaini Pallary, 1939

Original source. Pallary 1939: 93, pl. 6, figs 5–8, 14.

Type locality. “Sources du Nahr es Sine, au Sud de Lattaquié, sur la route de Beyrouth” [Nahr as Sinn, south of Latakia, at the road to Beirut], Syria.

Remarks. Heller et al. (1999: 61) considered the species as a junior synonym of *Melanopsis costata* (Olivier, 1804).

† *Melanopsis gersondei* Willmann, 1981

Original source. Willmann 1981: 143.

Type horizon. Palioskala Formation, middle Tortonian, late Miocene.

Type locality. “Bachriß beim Höhenzug von Palioskala, ca. 700 m westlich vom Kap Phoka” [brook section near the mountain range of Palioskala, ca. 700 m west of Ágios Fokás, Kos Island], Greece.

Types. Geological-Paleontological Institute, University of Kiel, Germany; no number indicated.

† *Melanopsis gibbosula* Grateloup, 1838

Original source. Grateloup 1838: 141, pl. 4, fig. 50.

Type horizon. Chattian–Burdigalian (?), late Oligocene–early Miocene.

Type locality. “Dax. [...] d’Abesse et de Quillac à Saint-Paul” [Dax; Abesse and Quillac at Saint-Paul-lès-Dax], France.

Remarks. Wenz (1929: 2774) considered the taxon as a junior synonym of “*Melanopsis major* Férussac, 1823”, which is not an available name.

Melanella gigantea Bourguignat, 1884

Original source. Bourguignat 1884: 19.

Type locality. “Rivière près d’une villa sur la route de Pregrada, aux environs de Krapina, en Croatie” [river near a villa on the road to Pregrada, near Krapina], Croatia.

Remarks. Note that Bourguignat denoted the authority as “Bourguignat, 1879”.

† *Melanopsis narzolina gigantea* Robles, 1975

Original source. Robles 1975: 362, pl. 2, figs 2–8.

Type horizon. Late Miocene.

Type locality. “Venta del Moro (Valencia)”, Spain.

Types. Museo Nacional de Ciencias Naturales, Madrid, coll. no. M-327.

† *Melanopsis costata* var. *glabra* Brusina, 1874

Original source. Brusina 1874: 41, pl. 7, fig. 9.

Type horizon. Cernikian, Pliocene.

Type locality. “Sibinj; Farkašić; Dubranjec”, Croatia.

Types. Milan et al. (1974: 89) stated that the single specimen of this taxon was lost.

† *Melanopsis (Melanoptychia) glabra* Pavlović, 1927 [invalid]

Original source. Pavlović 1927: 60, pl. 6, figs 18–19.

Type horizon. Middle Pannonian, late Miocene.

Type locality. “У Караџы” [from Karagača near Vrčin], Serbia.

Types. The illustrated syntype is stored in the Natural History Museum, Belgrade, coll. no. 209 (Milošević 1962: 23).

Remarks. Junior homonym of *Melanopsis glabra* Brusina, 1874. Neubauer et al. (2014a: 457) introduced *M. vrcinensis* as replacement name.

† “*Melanopsis glabra*” mentioned in Bogachev (1938: 13, 33, 53) [unavailable]

Horizon. Miocene.

Locality. “Tori bei Borshomi” [Tori near Borjomi], Georgia.

Remarks. The name was only mentioned in a species list by Bogachev without description or illustration. Moreover, he applied multiple original spellings: the name is given as “*glabra*” on p. 53, but “*glarba*” on p. 13, 33.

† *Melanoptychia lyrata glabra* Olujić, 1999 [invalid]

Original source. Olujić 1999: 20, 48, pl. 1, fig. 1.

Type horizon. Langhian, middle Miocene.

Type locality. It is unclear from the original work in which of the studied localities/sections along the valleys of the Sutina, Batarelov and Vojskava rivers (4 km W of Sinj) the taxon occurred and in which not, Croatia.

Remarks. Junior secondary homonym of *Melanopsis glabra* Brusina, 1874. Neubauer et al. (2011: 207) considered the taxon as a junior synonym of *Melanopsis lyrata* Neumayr, 1869.

Melanopsis parreyssii var. *glabrata* Clessin, 1890

Original source. Clessin 1887–1890: 690, fig. 475.

Type locality. “Ungarn ([...] ohne nähere Fundortangabe)” [Hungary, without indication of a locality], Hungary.

Remarks. Neubauer et al. (2014d: 125) considered this taxon as a junior synonym of *Microcolpia parreyssii* (Philippi, 1847).

† *Melanopsis glandicula* Sandberger, 1872

Original source. Sandberger 1870–1875: pl. 26, figs 3–3b.

Type horizon. Mammal zone MN 5, early–middle Miocene.

Type locality. “Pontlevoy bei Blois” [Pontlevoy near Blois], France.

Remarks. Plate 26 of Sandberger’s monograph was issued in 1872, while the description on p. 520 appeared in 1875 (Woodward 1906).

Melanopsis foleyi var. *glandinopsis* Pallary, 1928

Original source. Pallary 1928a: 271, pl. 6, figs 10, 15.

Type locality. “Aïn Mélias, près de Figuig” [Ain Melias near Figuig], Algeria.

† *Melanopsis glaubrechtii* Neubauer & Harzhauser in Neubauer et al., 2015

Original source. Neubauer et al. 2015b: 145, figs 3A–D.

Type horizon. Middle Pannonian, late Miocene.

Type locality. “Brickyard near the town of Martin, Turiec Basin”, Slovak Republic.

Types. Múzeum Andreja Kmeťa in Martin, Slovak Republic, coll. no. SNM 16/2011 (PZ-703g).

Microcolpia glinaica Bourguignat, 1884

Original source. Bourguignat 1884: 54.

Type locality. “La Glina en Hongrie” [Glina river], Croatia.

Remarks. Starobogatov et al. (1992: 65) considered the species as a junior synonym of *Microcolpia acicularis* (Férussac, 1823).

Melanopsis glinensis Brot, 1878

Original source. Brot 1874–1879: 370–371, pl. 38, fig. 4e.

Type locality. “Glina Fl.” [Glina river], Croatia.

Remarks. “In schedis” name by Parreyss, introduced by Brot (1878) in synonymy of “*Hemisinus acicularis* (Férussac, 1823)”. Clessin (1890) made it available by treating it as a valid name (see Note 2). Starobogatov et al. (1992: 65) again considered the species as a junior synonym of *Microcolpia acicularis*.

† *Melanopsis gorceixi* var. *globosa* Magrograssi, 1928

Original source. Magrograssi 1928: 259, pl. 6, fig. 14.

Type horizon. Plio-Pleistocene.

Type locality. “Coo: V. Bocasia, torrente Sefto, C. Foca, tra Antimachia e Pili” [Kos island: Vokasia valley, Sefto river, Ágios Fokás, between Antimácheia and Pýli], Greece.

Melanopsis pachya var. *gombaulti* Pallary, 1939

Original source. Pallary 1939: 87, pl. 5, figs 37–39.

Type locality. “Dans les sources de Mézérib, au N. O. de Derâa (Syrie méridionale)” [in the sources of the Muzayrīb, northwest of Dar’â], Syria.

† *Melanopsis gorceixi* Tournouër, 1875

Original source. Tournouër 1875: 76.

Type horizon. Tafi Formation, early Pleistocene.

Type locality. “Prope vicum Antimaki [...] et prope civitatem Cos” [near near the village Antimácheia and near the city of Kos], Greece.

Melanopsis goulvaini Pallary, 1916

Original source. Pallary 1916: 82.

Type locality. “Bourail” (Gassies 1874: 384), New Caledonia.

Remarks. Replacement name for *M. elongata* Gassies, 1874, non Férussac, 1822 (see Note 1).

Melanopsis gracilentia Pallary, 1911

Original source. Pallary 1911: 130, [unnumbered plate], fig. 19.

Type locality. “Tout près d’Oudjda, à 4 kilom. S.-E., sourdent les belles sources de Sidi-Yahia qui alimentent une véritable oasis, puis la ville d’Oudjda, et vont finalement se déverser dans l’oued Isly” [near Oujda, 4 km southeast, at the sources of Sidi Yahya that feed an oasis and the city of Oujda, and ultimately will flow into the Oued Isly], Morocco.

† *Melanopsis (Melanosteira) conemenosiana graciliformis* Papp, 1955

Original source. Papp 1955: 129, pl. 20, fig. 26.

Type horizon. Early Pleistocene.

Type locality. “Brochitza, Elis” [Vrokhítsa], Greece.

Types. Museum of Palaeontology and Geology of the University of Athens; no number indicated.

† *Melanopsis bouei* var. *gracilis* Brusina, 1874

Original source. Brusina 1874: 47.

Type horizon. Pannonian, zone C, late Miocene.

Type locality. “Sused (Sopot bei Goljak)” [Podsused, in Zagreb], Croatia.

Remarks. Wenz (1929: 2673) considered the variety as a junior synonym *M. bouei* Férussac, 1823.

† *Melanopsis neumayri* var. *gracilis* Fontannes, 1880 [invalid]

Original source. Fontannes 1879–1882: 174.

Type horizon. Mammal zone MN 11, late Miocene.

Type locality. “Les marnes à *Potamides Basteroti* de Visan (Vaucluse)” [in the marls with *P. basteroti* at Visan, Dép. Vaucluse], France.

Remarks. Junior homonym of *Melanopsis bouei gracilis* Brusina, 1874 (see Note 1). Wenz (1929: 2793) considered the variety as a junior synonym *M. neumayri* Tournouër, 1874.

Melanopsis costata var. *gracilis* Locard, 1883 [invalid]

Original source. Locard 1883a: 288.

Type locality. “Lac d’Homs” [Lake Homs], Syria.

Remarks. Junior homonym of *Melanopsis bouei gracilis* Brusina, 1874 (see Note 1).

† *Melanopsis (Hyphantria) gracilis* Handmann, 1887 [invalid]

Original source. Handmann 1887: 37, pl. 8, fig. 17.

Type horizon. Pannonian, zone B–D, late Miocene.

Type locality. “Leobersdorf”, Austria.

Remarks. Junior homonym of *Melanopsis bouei gracilis* Brusina, 1874 (see Note 1). Wenz (1929: 2673) considered the variety as a junior synonym *M. bouei* Férussac, 1823.

† *Melanopsis graciosa* Seninski, 1905

Original source. Seninski 1905: 59, pl. 1, fig. 16.

Type horizon. Duab Beds, middle to late Kimmerian, Pliocene.

Type locality. “Моквинские пласты, разрыв р. Дуабъ” [Mokvi layers at Duab river], Georgia.

† *Melanopsis gradata* Rolle, 1858

Original source. Rolle 1858: 28, pl. 2, fig. 13.

Type horizon. Paleocene.

Type locality. “Herrmanns-Stollen im Lubellina-Graben, Gemeinde Ober-Skallis, nordöstlich von Schönstein” [Hermanns adit in the Lubellina ditch in Škale, northeast of Šoštanj], Slovenia.

Remarks. After Wenz (1929: 2505) this species belongs in the genus *Pyrgulifera* Meek, 1877 (Pleuroceridae), but Yen (1958: 194) doubted this classification.

† *Melanopsis gradata* Fuchs, 1870 [invalid]

Original source. Fuchs 1870b: 539, pl. 20, figs 13–14.

Type horizon. Transdanubian, Pannonian, late Miocene.

Type locality. “Tihany” (p. 533), Hungary.

Remarks. Junior homonym of *Melanopsis gradata* Rolle, 1858. Wenz (1928b: 219) introduced *M. tihanyensis* as replacement name.

Melanopsis (Mesopotamia) khabourensis var. *gradata* Pallary, 1939 [invalid]

Original source. Pallary 1939: 103, pl. 5, fig. 20.

Type locality. “Ras el ‘Ain du Khabour” [Chabur river near Ra’s al ‘Ayn], Syria.

Remarks. Junior homonym of *Melanopsis gradata* Rolle, 1858.

Melanopsis graellsii Villa & Villa in Graells, 1846

Original source. Graells 1846: 17, [unnumbered plate], figs 16–19.

Type locality. Spain [no locality indicated].

Remarks. Graells (1846) indicated that the “brothers Villa” dedicated the species to him. The name “*graellsii*” as mentioned in Pallary (1924: 250) is an incorrect subsequent spelling. It is unclear why Pallary considered the species as a junior synonym of “*Buccinum annulatum* Chemnitz”. That species, which is not available from Chemnitz (1795: 91, pl. 188, figs 1812–1813) according to Opinion 184 (ICZN 1944), is not even a freshwater snail but dwells in the southern Pacific (Nassariidae?).

Melanopsis chehirensis var. *granifera* Pallary, 1939

Original source. Pallary 1939: 95.

Type locality. Not explicitly stated but probably the same as for the species (“Dans le source de Yeni Chehir, [...] entre Antioche et Alep, à l’intersection de la route d’Alexandrette” [at the source of Yenişehir, between Antakya and Aleppo, at the intersection of the road from İskenderun], Turkey).

† *Melanopsis granum* Calvert & Neumayr, 1880

Original source. Calvert and Neumayr 1880: 376, pl. 2, fig. 23.

Type horizon. Late Sarmatian, Khersonian, late Miocene.

Type locality. “Renkiöi” [north of İntepe], Turkey.

† “*Melanopsis grateloupii*” mentioned in Hoeninghaus (1831: 142) [unavailable]

Horizon. Burdigalian, early Miocene (?).

Locality. “Dax”, France.

Remarks. Nomen nudum. Grateloup (1838: 139) listed the name (as “*grateloupii*”) in synonymy of *Melanopsis aquensis* Grateloup, 1838. Wenz (1929: 2774) synonymized the name (as “*grateloupi*”) with “*M. major* Férussac, 1823”, which is not an available name.

Fagotia gravida Bourguignat, 1884

Original source. Bourguignat 1884: 40.

Type locality. “Rivières du bassin du lac Sabandja (Anatolie)” [rivers of the basin of Lake Sapanca], Turkey.

Remarks. Note that Bourguignat denoted the authority as “Bourguignat, 1880”. Starobogatov et al. (1992: 63) considered the species as a junior synonym of *Fagotia* [= *Esperiana*] *gallandi* Bourguignat, 1884.

† *Melanopsis groyei* Pallary, 1916

Original source. Pallary 1916: 78.

Type horizon. Sparnacian, early Ypresian, Eocene.

Type locality. “Entre St.-Germini et Carsoli” (Férussac 1823: 164) [between San Gemini and Carsoli], Italy.

Remarks. Introduced for a part of the material of *Melanopsis buccinoidea* sensu Férussac, 1823 (pl. 8, fig. 3).

† *Melanopsis guernei* Brusina, 1902

Original source. Brusina 1902: pl. 5, fig. 60.

Type horizon. Middle Pannonian, late Miocene.

Type locality. “Markuševac”, Croatia.

Types. Milan et al. (1974: 91) indicated a holotype, but it is uncertain whether the specimen is part of the original type series and whether it was the only one Brusina had at hand (holotype by monotypy, Art. 73.1.2). The specimen is stored in the Croatian Natural History Museum, Zagreb, coll. no. 2500-146.

Remarks. The name “*gueneri*” as mentioned in Wenz (1929: 2736) is an incorrect subsequent spelling.

***Melanopsis etrusca* var. *guineensis* Azpeitia Moros, 1929**

Original source. Azpeitia Moros 1929: 247, pl. 9, fig. 223.

Type locality. “De las cercanías del Cabo San Juan en la Guinea española” [in the vicinity of Cabo San Juan], Equatorial Guinea.

Remarks. Certainly not a Melanopsidae.

***Melanopsis guiraoi* Bourguignat, 1884**

Original source. Bourguignat 1884: 108.

Type locality. “Prope Cehejin Prov. Murcica” (Brot 1879: 438) [near Cehegín, prov. Murcia], Spain.

Remarks. Replacement name for *M. obesa* Brot, 1879, non Gassies, 1856.

***Melanopsis hadiensis* Pallary, 1928**

Original source. Pallary 1928a: 269, pl. 6, figs 11–14.

Type locality. “Bel Hadi-Kenadsa (Sud Oranais)” [Bel Hadi, near Kenadsa], Algeria.

***Microcolpia hagenmuelleriana* Bourguignat, 1884**

Original source. Bourguignat 1884: 57.

Type locality. “Le Danube près de Buda-Pesth, en Hongrie” [in the Danube river near Budapest], Hungary.

Remarks. Bourguignat (1884) listed the name as “*Hagenmulleria*” on p. 51 but as “*Hagenmülleriana*” on p. 57. To my knowledge no later author has acted as First Reviser sensu Art. 24.2.3 and has mentioned both names and selected one as correct. Starobogatov et al. (1992: 65) considered the species as a junior synonym of *Microcolpia cornea* (Pfeiffer, 1828).

† *Melanopsis entzi* var. *balavatsi* Pavlovic, 1927

Original source. Pavlović 1927: 68.

Type horizon. Middle Pannonian, late Miocene.

Type locality. “Kapagača” [Karagača near Vrčin], Serbia.

Remarks. From the writing in the original work it seems that Pavlović had apparently doubts about the validity of the taxon.

Melanopsis hammamensis Gassies, 1856

Original source. Gassies 1856: 11, figs 9–10.

Type locality. “L’Oued-el-Hammam [entre le Sig et Mascara]” [Oued el Hammam between Sig and Mascara], Algeria.

† *Melanopsis handmanni* Brusina, 1892 [invalid]

Original source. Brusina 1892: 140.

Type horizon. Early–middle Pannonian, late Miocene.

Type locality. “Kotingbrunn; [...] Leobersdorf”, Austria.

Types. The illustrated syntypes are stored in the Croatian Natural History Museum, Zagreb, coll. no. 2513-159/1-4 (Milan et al. 1974: 92).

Remarks. Introduced as replacement name for the senior (!) homonym *M. fuchsi* Handmann, 1882, non Brusina, 1884. Wenz (1929: 2813–2814) synonymized *M. handmanni* as well as *M. fuchsi* Handmann, 1882 with *M. pygmaea*.

† *Melanopsis handmanniana* W. Fischer, 1996

Original source. Fischer 1996a: 27, figs 3, 5–10.

Type horizon. Pannonian, zone B–D, late Miocene.

Type locality. “Leobersdorf”, Austria.

Types. Fischer (1996a: 27) designated a lectotype (for *M. constricta* Handmann), which is stored in the collection of the Geological Survey Austria, Vienna; no number indicated.

Remarks. Replacement name for *M. martiniana constricta* Handmann, 1887, non Brusina, 1878 (see Note 1).

† *Melanopsis hantkeni* Hofmann, 1870

Original source. Hofmann 1870: 26, pl. 3, figs 5a–c.

Type horizon. Egerian, late Oligocene–early Miocene.

Type locality. “A zsily-völgyi [...], továbbá Valia-Aninossaban és Paren lui Marin-ban, Lupény mellett. [...] gyakori Buda és Esztergom között [...], jelesen a Miklóshegyen, Annavölgyen, Mogyoróson és Pomázon” [in the Zsily valley; also in Valea Ani-noasa and Paren lui Marin (?) near Lupeni (Romania); between Buda and Esztergom, at Miklóshegy, Annavölgy, Mogyorós and Pomáz (Hungary)], Hungary.

† *Melanopsis haranti* De Laubrière & Carez, 1881

Original source. De Laubrière and Carez 1881: 403, pl. 16, figs 1–2.

Type horizon. Lutetian, Eocene.

Type locality. “Brasles”, France.

Remarks. Cossmann (1888: 279) and Wenz (1929: 2640) classified the species in the genus *Faunus* Montfort, 1810 (Pachychilidae).

Melanopsis costata var. *harpa* Westerlund, 1892

Original source. Westerlund 1892: 199.

Type locality. “Bei Sevilla im Guadalquivir” [near Sevilla, in the Guadalquivir river], Spain.

† *Melanopsis harpula* Neumayr in Neumayr & Paul, 1875

Original source. Neumayr and Paul 1875: 38, pl. 7, fig. 1.

Type horizon. Cernikian, Pliocene.

Type locality. “Čaplathal bei Podwin” [Čaplja graben near Slavonski Brod], Croatia.

† *Melanopsis hassiaca* Sandberger, 1873

Original source. Sandberger 1870–1875: 315.

Type horizon. Early Rupelian, Oligocene.

Type locality. “Grossalmerode [...]; Nordshausen” (Speyer 1870: 95–96), Germany.

Remarks. Introduced for *M. subulata* sensu Speyer, 1870, non Sowerby, 1822 and *M. praerosa* [= *M. praemorsa*] sensu Speyer, 1870, non Linnaeus, 1758.

† *Melanopsis hastata* Neumayr in Neumayr & Paul, 1875

Original source. Neumayr and Paul 1875: 40, pl. 7, figs 7–8.

Type horizon. Cernikian, Pliocene.

Type locality. “Gromačnik; [...] Slobodnica; [...] Sabin [Sibinj]; [...] Malino”, Croatia.

† *Melanopsis haueri* Handmann, 1882

Original source. Handmann 1882: 558.

Type horizon. Pannonian, zone D, late Miocene.

Type locality. “Kottingbrunn [...] Ziegelei a”, Austria.

† *Melanopsis haugi* Popescu-Voitești, 1910

Original source. Popescu-Voitești 1910: 360, pl. 21 (4), figs 4–4l.

Type horizon. Lutetian, Eocene.

Type locality. “Gropile Vulpilor près Titești” [Gropile Vulpilor (?) near Titești], Romania.

Remarks. Wenz (1929: 2570) classified the species within the genus *Coptostylus* Sandberger, 1872 (Thiaridae).

† *Melanopsis hazayi* Brusina, 1903

Original source. Brusina 1903: 112.

Type horizon. Late Pleistocene–Holocene.

Type locality. “Bischofsbad” [Püspökfürdő, Băile 1 Mai, Lake Peșea], Romania.

Remarks. Neubauer et al. (2014d: 125) considered this taxon as a junior synonym of *Microcolpia parreyssii sikorai* (Brusina, 1903).

† *Melania heberti* Hantken, 1878

Original source. Hantken 1878: 179–180, fig. 23.

Type horizon. Late Santonian–early Campanian, late Cretaceous.

Type locality. “[Ajka]” (p. 175, 178), Hungary.

Remarks. Currently attributed to the melanopsid genus *Campylostylus* (Bandel and Riedel 1994: 21).

Melanopsis hebraica Bourguignat, 1884

Original source. Bourguignat 1884: 131.

Type locality. “Aïn-Saadi, près de Kaifa, en Syrie” [Aïn Saadi, in Haifa], Israel.

Remarks. Heller et al. (2002: 601) considered the species as a junior synonym of *Melanopsis saulcyi* Bourguignat, 1853.

† *Amphimelania heckneri* Koch, 1917

Original source. Koch 1917: 9, figs 1–6.

Type horizon. Cernikian, Pliocene.

Type locality. “U okolici Dubranca, [...] u Zagreb šumi, [...] kod Prvonožina” [near Dubranec, in the Zagreb woods, and at Prvonožina], Croatia.

† *Melanopsis heeri* Mayer-Eymar, 1887

Original source. Mayer-Eymar 1887: 68, pl. 6, fig. 10.

Type horizon. Chattian, Oligocene.

Type locality. “Schloss Ralligen” [Ralligen Castle near Sigriswil], Switzerland.

Remarks. Appeared first as a nomen nudum (as “*heerii*”) in Sandberger (1873: 341).

† *Melanopsis heldreichi* Neumayr, 1880

Original source. Neumayr 1880b: 295, pl. 2, figs 2–3.

Type horizon. Plio-Pleistocene.

Type locality. “Zwischen Pyllé und Antimachia; [...] beim Castell von Antimachia und in der näheren Umgebung des Dorfes Antimachia” [between Pýli and Antimácheia; near the citadel of and in the vicinity of the village Antimácheia, Kos Island], Greece.

Melania helena Von dem Busch in Philippi, 1847

Original source. Philippi 1847: 170, pl. 4, fig. 4.

Type locality. “Insula Java”, Indonesia.

Remarks. Although described as *Melania* and not considered a Melanopsidae today, this species is included and briefly discussed here because it has been repeatedly considered a *Melanopsis*. The specimens had been apparently sent to Von dem Busch or Philippi by Meder, who had named the species “*Melanopsis Helena*”. The description, however, stems clearly from Von dem Busch. The species epithet is a noun in apposition and needs not to agree in gender with the generic name (Art. 31.2.1). Currently considered to belong in the marine genus *Clea* H. Adams & A. Adams, 1855 (Buccinidae) (e.g., Coelho et al. 2013).

***Melanopsis heliophila* Bourguignat, 1884**

Original source. Bourguignat 1884: 161.

Type locality. “Dans l’oasis d’Aïn-Chair, à l’extrême sud saharien du Maroc” [in the oasis of Ain Chair (= Oued Mellah), at the far southern Sahara in Morocco], Morocco.

Remarks. Note that Bourguignat denoted the authority as “Bourguignat, 1872”.

† ***Melania bellespontica* Calvert & Neumayr, 1880**

Original source. Calvert and Neumayr 1880: 374, pl. 2, fig. 14.

Type horizon. Late Sarmatian, Khersonian, late Miocene.

Type locality. “Renkiöi” [north of İntepe], Turkey.

Remarks. Considered to belong in the genus *Amphimelania* by Wenz (1929: 2874).

† ***Melanopsis hemimorpha* Blanckenhorn, 1897**

Original source. Blanckenhorn 1897: 138, pl. 10, fig. 26.

Type horizon. Plio-Pleistocene.

Type locality. “Dschisr esch-Schurr, Dreissensiaschicht” [Jisr Ash-Shughur, *Dreissena* layer], Syria.

† ***Melanopsis hennersdorfensis* W. Fischer, 1993**

Original source. Fischer 1993: 180, figs 1, 3.

Type horizon. Pannonian, zone D, late Miocene.

Type locality. “Tegelgrube der Fa. Wienerberger, Hennersdorf, NÖ” [claypit of the Wienerberger company at Hennersdorf, Lower Austria], Austria.

Types. Geological Survey Austria, Vienna; no number indicated.

† ***Melanopsis hericarti* Fontannes, 1881**

Original source. Fontannes 1881: 981, pl. 1, figs 5–6.

Type horizon. Rupelian, Oligocene.

Type locality. “Baume-Cornillane”, France.

***Melanopsis hiera* Bourguignat, 1884**

Original source. Bourguignat 1884: 121.

Type locality. “Sources du Jourdain; [...] d’Aïn-el-Mellaha, dans la plaine du Bahr-el-Houlé; [...] lacs d’Homs et d’Antioche” [Sources of the Jordan; [...] Aïn Mallahah, in the plains of the Hula valley; [...] lakes Homs and Anuk (also as Amik)], Syria.

Remarks. Introduced for *M. costata* sensu Kobelt, 1880, non Olivier, 1804. Bourguignat attributed the authority to Letourneux, but there is no evidence that the description really derived from that author. Heller et al. (2005: 243) listed the species as a junior synonym of *Melanopsis costata* (Olivier, 1804) as well as of *Melanopsis saulcyi* Bourguignat, 1884.

† *Fusus (Anura) decipiens* var. *hilarionis* De Gregorio, 1880

Original source. De Gregorio 1880: 89, pl. 7, fig. 61.

Type horizon. Eocene.

Type locality. “San Giovanni Ilarione”, Italy.

Remarks. Considered a *Melanopsis* and a separate species by Quaggiotto and Mellini (2008: 44).

Melania (Amphimelania) hispanica Westerlund, 1898

Original source. Westerlund 1898: 178.

Type locality. “Albarracin”, Spain.

† *Melanopsis hispidula* Pallary, 1916

Original source. Pallary 1916: 84.

Type horizon. Pannonian, zone B–D, late Miocene.

Type locality. “Leobersdorf” (Handmann 1887), Austria.

Remarks. Replacement name for the junior homonym *M. turrita* Handmann, 1887, non Rossmässler, 1854.

† *Melanopsis hoernesii* Sandberger, 1872

Original source. Sandberger 1870–1875: pl. 25, figs 32–32a.

Type horizon. Burdigalian, early Miocene.

Type locality. “St. Paul, Mandillot (Landes)”, France.

Remarks. Plate 25 of Sandberger’s monograph was issued in 1872, while the description on p. 512 appeared in 1875 (Woodward 1906). Introduced for *M. buccinoides* [sic] sensu Grateloup, 1840, non Olivier, 1801, for which d’Orbigny (1852) had already introduced *M. subbuccinoides* as replacement name. Hence, *M. hoernesii* is a junior objective synonym of *M. subbuccinoides*. Sandberger attributed the authority to Mayer, apparently based on an “in schedis” determination.

***Melanopsis costata* var. *hoernesi* Blanckenhorn, 1897 [invalid]**

Original source. Blanckenhorn 1897: 130, pl. 10, figs 9–10.

Type horizon. Pleistocene (?)–Recent.

Type locality. “Fossil [...] bei Antakije. Lebend am unteren Orontes [...], desgleichen im Kara Su, einem nördlichen Zufluss des Sees von Antiochia und im Sadjür Su bei Aleppo” [as fossil near Antakya (Turkey). Alive in the lower Orontes (Turkey/Syria), as well as in the Karasu, a northern tributary of the Lake Anuk (Turkey), and in the Säjür river near Aleppo (Syria)].

Remarks. Junior homonym of *M. hoernesi* Sandberger, 1872 (note that latter is a junior objective synonym of *M. subbuccinoides* and thus invalid). Heller et al. (2005: 244) considered the species as a junior synonym of *Melanopsis costata* (Olivier, 1804).

***Melania holandrii* Pfeiffer, 1828**

Original source. Pfeiffer 1828: 47.

Type locality. “Bei Kroatisch Feistritz, am Fusse des Berges Terglou in Illyrien” [near Bohinjska Bistrica, at the foot of Mt. Triglav], Slovenia.

Remarks. Type species of *Holandriana* Bourguignat, 1884 and *Amphimelania* P. Fischer, 1885. The names “*holandri*” and “*hollandri*”, each occurring multiple time in the literature (e.g., Walderdorff 1864: 512; Brusina 1867: 85; Tschapeck 1881: 101), are incorrect subsequent spellings.

† *Melanopsis branilovici* Brusina, 1897

Original source. Brusina 1897: 12, pl. 5, fig. 12.

Type horizon. Pannonian, late Miocene.

Type locality. “Ad viam inter Sarajevo et Lukavica” [at the road between Sarajevo and Lukavica], Bosnia and Herzegovina.

Types. Milan et al. (1974: 92) indicated a holotype, but it is uncertain whether the specimen was the only one Brusina had at hand (holotype by monotypy, Art. 73.1.2). The specimen is stored in the Croatian Natural History Museum, Zagreb, coll. no. 2985-631.

***Melanopsis huidobroi* Azpeitia Moros, 1929**

Original source. Azpeitia Moros 1929: 326, pl. 12, fig. 280.

Type locality. “Guadalquivir en Lora del Río, Sevilla” [in the Guadalquivir river at Lora del Río, prov. Sevilla], Spain.

***Melanopsis humilis* Pallary, 1928**

Original source. Pallary 1928a: 255, pl. 4, figs 14–15.

Type locality. “Fès et Meknès”, Morocco.

† *Melanopsis hungarica* Kormos, 1904

Original source. Kormos 1904: 107, fig. 2.

Type horizon. Late Pleistocene–early Holocene.

Type locality. “Püspökfürdő” [Băile 1 Mai, Lake Pețea], Romania.

Remarks. Neubauer et al. (2014d: 125) considered this taxon as a junior synonym of *Microcolpia parreyssii* (Philippi, 1847).

† *Melanopsis hungarica* Pallary, 1916 [invalid]

Original source. Pallary 1916: 83.

Type horizon. Transdanubian, Pannonian, late Miocene.

Type locality. “Radmanest” (Fuchs 1870: 353) [Rădmănești], Romania.

Remarks. Replacement name for *M. fuchsi* Brusina, 1884 non Handmann, 1882, which itself was introduced for *M. costata* sensu Fuchs, 1870, non Olivier, 1804. It is a junior homonym of *M. hungarica* Kormos, 1904 (see *M. confusa* Strausz, 1941).

† *Melanopsis hybostoma* Brusina, 1874

Original source. Brusina 1874: 31.

Type horizon. Cernikian, Pliocene.

Type locality. “Podvinje (Čaplja)” [Čaplja trench near Slavonski Brod], Croatia.

Remarks. The names “*hibostoma*” and “*hypostoma*” as mentioned in Pană (2003: 324) and Macarovici (1940: 344) are incorrect subsequent spellings.

“*Melanopsis tingitana* var. *hybrida*” mentioned in Brot (1874–1879: 442) [unavailable]

Locality. “Morocco, Mogador; [...] Spanien, Andalousien” [Morocco, Essaouira; Spain, Andalusia].

Remarks. Nomen nudum, listed in synonymy of “*M. tingitana* var. β ” by Brot (1879), which is described briefly in Latin. He attributed the name to Morelet, but it remains unclear why - the only author mentioning a (nameless) variety of *M. tingitana* that matches the description by Brot is Mousson (1874).

† *Melanopsis impressa* Krauss, 1852

Original source. Krauss 1852: 143, pl. 3, fig. 3.

Type horizon. Ottnangian, middle Burdigalian, early Miocene.

Type locality. “Kirchberg an der Iller” [Illerkirchberg], Germany.

† *Melanopsis incerta* Férussac, 1822

Original source. Férussac 1819–1851: *Mélanopsides fossiles*, pl. 1 (1822), fig. 12.

Type horizon. Sarmatian (sensu lato), middle–late Miocene.

Type locality. “De Sestos” [Sestos, prov. Çanakkale], Turkey.

Remarks. The species was also illustrated on pl. 2 of the “*Mélanopsides fossiles*” in the “*Histoire naturelle*” (Férussac 1819–1851) and on pl. 8 in Férussac (1823). However, Férussac obviously mixed up *M. incerta* with *M. daudebartii* and erroneously illustrated the latter of both species, which was already recognized by Deshayes when preparing the final explanations for the plates of the “*Histoire naturelle*”.

† *Melanopsis incerta* Fuchs, 1877 [invalid]

Original source. Fuchs 1877: 14, pl. 2, figs 13–21.

Type horizon. Pliocene.

Type locality. “Megara”, Greece.

Remarks. Junior homonym of *Melanopsis incerta* Férussac, 1822. Pallary (1920b: 110) introduced *M. revelata* as replacement name.

Melanopsis inconspicua Pallary, 1928

Original source. Pallary 1928a: 261, pl. 4, figs 22–25.

Type locality. “Sources tièdes un peu avant Kerrando, au S.-E. de Rich (grand Atlas oriental, versant sud)” [hot Springs just before Kerrando, southeast of Rich (High Atlas range, southern side)], Morocco.

Microcolpia inconspicua Starobogatov in Starobogatov et al., 1992

Original source. Starobogatov et al. 1992: 66, fig. 3 (17).

Type locality. “Фёслау, близ Вены” [Vöslau near Vienna], Austria.

Types. Zoological Institute of Russian Academy of Sciences, St.-Petersburg; no number indicated.

† *Melanopsis inconstans* Neumayr, 1869

Original source. Neumayr 1869: 356, pl. 11, figs 9–18.

Type horizon. Langhian, middle Miocene.

Type locality. “Miocic” [Miočić], Croatia.

Types. Illustrated syntypes are stored at the Geological Survey Austria, Vienna, coll. no. 1869/01/3/1-10.

Melania (Amphimelania) induta Westerlund, 1898

Original source. Westerlund 1898: 179.

Type locality. “Muchalatka” [not found; stated to be on the Crimean Peninsula], Ukraine.

† *Melanopsis inermis* Handmann, 1882

Original source. Handmann 1882: 554.

Type horizon. Pannonian, zone D, late Miocene.

Type locality. “Kottingbrunn [...] Ziegelei a”, Austria.

† *Melanopsis klerici inermis* Brusina, 1897 [invalid]

Original source. Brusina 1897: 8.

Type horizon. Pannonian, zone D–E, late Miocene.

Type locality. “Begaljica”, Serbia.

Types. Milan et al. (1974: 93) indicated a holotype, but it is uncertain whether the specimen was the only one Brusina had at hand (holotype by monotypy, Art. 73.1.2). The specimen is stored in the Croatian Natural History Museum, Zagreb, coll. no. 3020-666.

Remarks. Junior homonym of *Melanopsis inermis* Handmann, 1882. Neubauer et al. (2014a) introduced *M. magyari* as replacement name.

† *Boistelia inermis* Jekelius, 1944 [invalid]

Original source. Jekelius 1944: 137, pl. 57, figs 22–27.

Type horizon. Early Pannonian, late Miocene.

Type locality. “Turislav-Tal bei Soceni” [Turislav valley near Soceni], Romania.

Remarks. Junior secondary homonym and junior synonym of *M. inermis* Handmann, 1882 (see discussion in Neubauer et al. 2014a: 463).

† *Melanopsis inexpectata* Willmann, 1981

Original source. Willmann 1981: 193.

Type horizon. Gurniati Formation, late Pliocene to early Pleistocene.

Type locality. “Küstenaufschluß ca. 4 km südwestl. von Kardhamena/Kos (Profil K 3)” [section K 3, ca. 4 km southwest of Kardámaina, Kos Island], Greece.

Types. Geological-Paleontological Institute, University of Kiel, Germany; no number indicated.

Remarks. The name “*inexpectata*” as given by Kapan Yeşilyurt and Taner (2002: 107) is an incorrect subsequent spelling.

† *Melanopsis buccinoidea* var. γ) *antiquua inflata* Férussac, 1822

Original source. Férussac 1819–1851: *Mélanopsides fossiles*, pl. 1 (1822), figs 1–3, 9.

Type horizon. Eocene; late Villafranchian, early Pleistocene.

Type locality. “Des environs d’Épernay. [...] Du dépôt situé entre St.-Germini et Carsoli” [surroundings of Épernay (France); from deposits between San Gemini and Carsoli (Italy)].

Remarks. Introduced within the var. γ , to which Férussac attributed the term “*antiquua*”, which was probably not intended as species-group name (see introduction for a detailed discussion of the names introduced by Férussac 1823).

† *Melanopsis pygmaea* var. *inflata* Handmann, 1882 [invalid]

Original source. Handmann 1882: 553.

Type horizon. Pannonian, zone D, late Miocene.

Type locality. “Kottingbrunn [...] Ziegelei a, Ziegelei c”, Austria.

Remarks. Junior homonym of *Melanopsis buccinoidea inflata* Férussac, 1822. Wenz (1929: 2813) considered the taxon as a junior synonym of *Melanopsis pygmaea* Hörnes, 1856.

† *Melanopsis pygmaea inflata* Sauerzopf, 1952 [invalid]

Original source. Sauerzopf 1952: 13, pl. 2, fig. 4.

Type horizon. Pannonian, zone D–F, late Miocene.

Type locality. “Stegersbach, Litzelsdorf, Olbendorf, Oberdorf”, Austria.

Remarks. Junior homonym of *Melanopsis buccinoidea inflata* Férussac, 1822. Neubauer et al. (2014c: 28) considered the taxon as a junior synonym of *Melanopsis fuchsi* Handmann, 1882.

***Melanopsis costata* var. *inflexa* Pallary, 1939**

Original source. Pallary 1939: 90.

Type locality. Not explicitly stated but probably the same as for the species (“Disjre ech Chogour” [Jisr Ash-Shughur], Syria).

***Melanopsis costata* var. *infracincta* Martens, 1874**

Original source. Martens 1874: 32, pl. 5, fig. 38.

Type locality. “Quellen des Chabur bei Ras-el-Ain” [source of Chabur river near Ra’s al ‘Ayn], Syria.

***Melanopsis magnifica* var. *ingens* Pallary, 1928**

Original source. Pallary 1928a: 273, pl. 5, fig. 15.

Type locality. “Aïn Chekef, près Fès” [Aïn Chkef, near Fes], Morocco.

***Melanopsis parreyssi* f. *innodata* Westerlund, 1886**

Original source. Westerlund 1886: 123.

Type locality. “Bei Gross-Wardein in der schnellen Körös” [near Oradea in the Crişul Repede], Romania.

Remarks. Based on a specimen of *M. parreyssii* figured in Kobelt (1880: 19, pl. 189, fig. 1909d). Neubauer et al. (2014d: 125) considered this taxon as a junior synonym of *Microcolpia parreyssii* (Philippi, 1847).

***Melanopsis monteli* var. *innodata* Pallary, 1936 [invalid]**

Original source. Pallary 1936: 61.

Type locality. Not explicitly stated but probably the same as for the species (“L’Oued Sous, au pont des Aït Melloul, sur la route d’Agadir à Tiznit, à 14 kil. S. O. d’Agadir” [in the Oued Sous, at the bridge of Ait Melloul, at the road from Agadir to Tiznit, 14 km southwest of Agadir], Morocco).

Remarks. Junior homonym of *M. parreyssii innodata* Westerlund, 1886 (see Note 1).

***Melanopsis costata* var. *inodata* Pallary, 1939**

Original source. Pallary 1939: 91.

Type locality. Not explicitly stated but probably the same as for the species (“Djishr ech Chegour” [Jisr Ash-Shughur], Syria).

***Melanopsis insignis* Locard, 1883**

Original source. Locard 1883a: 202.

Type locality. “Samava” [As Samawah], Iraq.

Remarks. Appeared first as a nomen nudum in Martens (1874: 43, 67). Locard (1883a) introduced the species by referring to “*Melanopsis Turcica* de Mousson [...] 1874, p. 44” [act. p. 49], which he obviously considered different from the variety described in Mousson (1874: 33).

† ***Melanopsis intermedia* Rzehak, 1883**

Original source. Rzehak 1883: 43, pl. 2, figs 7a–c.

Type horizon. *Oncophora* Beds, middle Burdigalian, early Miocene.

Type locality. “Eibenschitz, [...] Oslawan” [Ivančice; Oslavany], Czech Republic.

Remarks. Čtyroký (1972: 74) considered this taxon as a junior synonym of *Melanopsis impressa* Krauss, 1852.

† ***Melanopsis (Lyrcea) senatoria* var. *intermedia* Handmann, 1887 [invalid]**

Original source. Handmann 1887: 19, pl. 2, fig. 14.

Type horizon. Pannonian, zone B–D, late Miocene.

Type locality. “Leobersdorf”, Austria.

Remarks. Junior homonym of *Melanopsis intermedia* Rzehak, 1883.

***Melanopsis mellalensis* var. *intermedia* Pallary, 1928 [invalid]**

Original source. Pallary 1928b: 19, pl. 2, fig. 20.

Type locality. “À Béni Mellal” [at Beni Mellal], Morocco.

Remarks. Junior homonym of *Melanopsis intermedia* Rzehak, 1883.

† ***Melanopsis involuta* Handmann, 1882**

Original source. Handmann 1882: 554.

Type horizon. Pannonian, zone D, late Miocene.

Type locality. “Kottingbrunn [...] Ziegelei a, Ziegelei c”, Austria.

***Melanopsis iraqensis* Pallary, 1939**

Original source. Pallary 1939: 89, pl. 4, figs 1, 2, 4, 5.

Type locality. “Tappah, à 3 km Est de Belad Sindjar et ‘Ain Haglan” [Tappah, 3 km east of Sinjar and Ain Haglan (?)], Iraq.

***Melanopsis irellii* Pallary, 1920**

Original source. Pallary 1920b: 111.

Type locality. “M. Mario” (Cerulli-Irelli 1914: 184), Italy.

Remarks. Introduced for *M. nodosa* sensu Cerulli-Irelli, 1914, non Férussac, 1822. Girotti (1972: 232) considered *M. nodosa* sensu Cerulli-Irelli, 1914 a junior synonym of “*Melanopsis affinis* Férussac”, which is not an available name.

***Melanopsis jordanica* var. *irregularis* Mousson, 1861**

Original source. Mousson 1861: 149.

Type locality. “Lac de Tiberias” [Sea of Galilee], Israel.

† *Melanopsis irregularis* Handmann, 1882 [invalid]

Original source. Handmann 1882: 557.

Type horizon. Pannonian, zone D, late Miocene.

Type locality. “Kottingbrunn [...] Ziegelei a”, Austria.

Remarks. Junior homonym of *Melanopsis jordanica irregularis* Mousson, 1861 (see Note 1).

***Melanopsis isabelae* Ahuir Galindo, 2016**

Original source. Ahuir Galindo 2016: 27.

Type locality. “In a Za river affluent, between Guefait and Hassi Blal”, Morocco.

Types. Museo Malacologico di Cupra Marittima, Italy; no number indicated.

***Melanopsis isseli* Bourguignat, 1884**

Original source. Bourguignat 1884: 115.

Type locality. “Dans le lac d’Accesa, près de Massa, en Toscane (Italie); aux environs d’Oran; [...] dans la vallée du Nahr-el-Kelb, près de Beyrouth” [in the Lago

dell'Accesa near Massa Marittima, in Tuscany (Italy); in the surroundings of Oran (Algeria); in the cave of Nahr el-Kalb near Beirut (Lebanon)].

Remarks. Introduced for *M. maroccana* sensu Bourguignat, 1864, non “Chemnitz” (which is not an available name; see there for details).

† *Melanopsis martiniana* var. *italica* Sandberger, 1872

Original source. Sandberger 1870–1875: 666, 687, pl. 26, fig. 25.

Type horizon. Tortonian–early Messinian, late Miocene.

Type locality. “Stazzano”, Italy.

Remarks. Plate 26 of Sandberger’s monograph appeared already in 1872, while the description on p. 666 (see also p. 687) was issued in 1875 (Woodward 1906). Harzhauser et al. (2015: 10) considered the taxon as a junior synonym of *Melanopsis narzolina* d’Archiac in Viquesnel, 1846.

† *Melanopsis (Stilospirula) jansseni* Anderson, 1964

Original source. Anderson 1964: 208, pl. 13, fig. 118.

Type horizon. Middle Miocene.

Type locality. “Bachbett der Küningsbeke an der Königsmühle bei Dingden” [Küningsbeke brook at the Königsmühle near Dingden], Germany.

Remarks. Probably not a Melanopsidae.

† *Melanopsis jasonis* Stache in Sandberger, 1871

Original source. Sandberger 1870–1875: 133, pl. 19, fig. 12.

Type horizon. Liburnian, Danian, Paleocene.

Type locality. “Zablachie bei Sebenico in Dalmatien” [Zablaće near Šibenik], Croatia.

Melanopsis jebusitica Bourguignat, 1884

Original source. Bourguignat 1884: 126.

Type locality. “Dans quelques sources de la plaine de Jéricho (Syrie)” [in a few springs in the plains of Jericho], Palestine.

Remarks. Bourguignat denoted the authority as “Letourneux, 1882”, but there is no evidence that the description really derived from that author.

† *Melanopsis (Canthidomus) jekeliusi* Gillet & Marinescu, 1971 [invalid]

Original source. Gillet and Marinescu 1971: 55, pl. 22, figs 20–30.

Type horizon. Transdanubian, Pannonian, late Miocene.

Type locality. “Radmanest” (Fuchs 1870: 353) [Rădmănești], Romania.

Remarks. Replacement name for the junior homonyms *M. fuchsi* Brusina, 1902, non Handmann, 1882 and *M. hungarica* Pallary, 1916, non Kormos, 1904, which were in turn introduced for *M. costata* sensu Fuchs, 1870, non Olivier, 1804. However, already Strausz (1941) had introduced *M. confusa* as replacement name, which makes *M. jekeliusi* its junior objective synonym.

Melanopsis costata var. *jordanica* Roth, 1839

Original source. Roth 1839: 25, pl. 2, figs 12–13.

Type locality. “In flumine Jordano; in mari Galilaeo” [in Jordan river; in Sea of Galilee], Israel.

Remarks. The name “*judaica*” as mentioned in Morelet (1880: 75) is an incorrect subsequent spelling.

Melanopsis jordanicensis Germain, 1921 [invalid]

Original source. Germain 1921: 466, 496.

Remarks. Unjustified emendation and therefore junior objective synonym of *M. jordanica* Roth, 1839.

† *Melanopsis kacici* Brusina, 1902

Original source. Brusina 1902: pl. 6, figs 56–57.

Type horizon. Middle Pannonian, late Miocene.

Type locality. “Markuševac”, Croatia.

Types. The syntypes are stored in the Croatian Natural History Museum, Zagreb; no number indicated (Milan et al. 1974: 92).

Remarks. Wenz (1929: 2678) considered this taxon as a junior synonym of *Melanopsis bouei* Férussac, 1823.

† *Melanopsis (Lyrcea) vindobonensis* var. *karagacensis* Pavlović, 1927

Original source. Pavlović 1927: 84, pl. 12, figs 1–2.

Type horizon. Middle Pannonian, late Miocene.

Type locality. “Капагач” [from Karagača near Vrčin], Serbia.

Types. The illustrated syntype is stored in the Natural History Museum, Belgrade, coll. no. 226 (Milošević 1962: 23).

† “*Melanopsis kaltenbachii*” mentioned in Mertin (1939: 254) [unavailable]

Horizon. Heidelberg Formation, late Santonian, late Cretaceous.

Locality. “Flugplatz Quedlinburg” [airfield at Quedlinburg], Germany.

Remarks. Nomen nudum. Obviously a lapsus calami of *Odostomia kaltenbachii* Mertin, 1939.

† *Melanopsis karici* Pavlović, 1903

Original source. Pavlović 1903: 156, pl. 3, figs 3–6.

Type horizon. Middle Miocene.

Type locality. “Из Бабуноз Дола близу Скопља” (p. 155) [Babin Dol near Skopje], Macedonia.

Types. The illustrated syntype is stored in the Natural History Museum, Belgrade, coll. no. 1448 (Milošević 1962: 24).

† *Melanopsis (Lyrcaea) nobilis karpovi* Gozhik in Gozhik & Datsenko, 2007

Original source. Gozhik and Datsenko 2007: 94, pl. 88, fig. 5.

Type horizon. Pliocene.

Type locality. “Оз. Сасык” [Lake Sasyk], Ukraine.

Types. Institute of Geological Sciences of the National Academy of Sciences of Ukraine, coll. no. 4602.

† *Melanopsis katzeri* Brusina, 1904

Original source. Brusina 1904: 495, pl. 2, fig. 4, pl. 3, fig. 4.

Type horizon. Langhian, middle Miocene.

Type locality. “Varcar-Vakufa” [Mrkonjić Grad], Bosnia and Herzegovina.

† *Melanopsis kernerii* Kühn, 1946

Original source. Kühn 1946: 75, textfig. 6.

Type horizon. Late middle Eocene–early Oligocene.

Type locality. “Monte Promina” [Promina Mountains], Croatia.

Types. Geological-Paleontological Department, Natural History Museum Vienna, Austria, coll. no. 1869/0009/0013.

***Melanopsis (Mesopotamia) khabourensis* Pallary, 1939**

Original source. Pallary 1939: 102, pl. 5, figs 13–21.

Type locality. “Sources de Khabour, dites Ras el ‘Ain” [source of Chabur river near Ra’s al ‘Ayn], Syria.

“*Melanopsis kindermanni* Zelebor” mentioned in Brot (1874–1879: 429) [unavailable]

Locality. Not indicated.

Remarks. Nomen nudum, apparently based on an unpublished manuscript name from Zelebor listed in the synonymy list of *M. saulcyi* by Brot (1879).

† *Melanopsis kispatici* Brusina, 1897

Original source. Brusina 1897: 10.

Type horizon. Langhian, middle Miocene.

Type locality. “Miočić”, Croatia.

Types. Milan et al. (1974: 93) indicated a holotype, but it is uncertain whether the specimen is part of the original type series and whether it was the only one Brusina had at hand (holotype by monotypy, Art. 73.1.2). The specimen is stored in the Croatian Natural History Museum, Zagreb, coll. no. 2970-616.

Remarks. Neubauer et al. (2013: 135) considered this taxon as a junior synonym of *Melanopsis lyrata* Neumayr, 1869.

† *Melanopsis (Canthidomus) kittli* Handmann, 1887

Original source. Handmann 1887: 32, pl. 7, fig. 14.

Type horizon. Pannonian, zone B–D, late Miocene.

Type locality. “Leobersdorf”, Austria.

Remarks. Wenz (1929: 2675) considered the taxon as a junior synonym of *Melanopsis bouei* Férussac, 1823.

† *Melanoptychia klecakiana* Bourguignat, 1880

Original source. Bourguignat 1880: 37.

Type horizon. Langhian, middle Miocene.

Type locality. “Vallée de la Cettina” [Cetina river valley], Croatia.

Remarks. The taxon is not included in the Fossilium Catalogus of Wenz (1929).

† *Melanopsis kleinii* Kurr, 1856

Original source. Kurr 1856: 42.

Type horizon. Middle–late Burdigalian, early Miocene.

Type locality. “Zwiefalten und Andelfingen” (Dunker 1848: 159), Germany.

Remarks. Introduced for *M. praerosa* [= *M. praemorsa*] sensu Dunker, 1848 and Klein, 1852, non Linnaeus, 1758. The name “*kleini*” as mentioned in Wenz (1929: 2762) is an incorrect subsequent spelling.

† *Melanopsis klerici* Brusina, 1893

Original source. Brusina 1893: 203 (Serbian part), 53 (Italian part).

Type horizon. Pannonian, zone D–E, late Miocene.

Type locality. “Begaljica”, Serbia.

Types. The syntypes are stored in the Croatian Natural History Museum, Zagreb; no number indicated (Milan et al. 1974: 93).

Melania (Melanopsis) kotschy Von dem Busch in Philippi, 1847

Original source. Philippi 1847: 175, pl. 4, fig. 11.

Type locality. “Persepolis Persiae”, Iran.

† *Melanopsis kottingbrunnensis* Handmann, 1882

Original source. Handmann 1882: 559.

Type horizon. Pannonian, zone D, late Miocene.

Type locality. “Kottingbrunn [...] Ziegelei a”, Austria.

Remarks. Wenz (1929: 2741) considered the taxon as a junior synonym of *Melanopsis haueri* Handmann, 1882.

† “*Melanopsis kozlovskii*” mentioned in Bogachev (1938: 13, 33, 53) [unavailable]

Horizon. Miocene.

Locality. “Tori bei Borshomi” [Tori near Borjomi], Georgia.

Remarks. The name was only mentioned in a species list by Bogachev without description or illustration.

† *Melanopsis krambergeri* Brusina, 1892

Original source. Brusina 1892: 142.

Type horizon. Middle Pannonian, late Miocene.

Type locality. “Markuševac”, Croatia.

Types. Milan et al. (1974: 93) indicated collection numbers for “syntypes” illustrated in Brusina (1902), but it is uncertain whether the specimens were part of the original type series. They are stored in the Croatian Natural History Museum, Zagreb, coll. no. 2499-145/1-2.

Remarks. Appeared first as a nomen nudum in Brusina (1884a: 138).

† *Amphimelania krambergeri* Brusina, 1902

Original source. Brusina 1902: pl. 5, figs 10–11.

Type horizon. Cernikian, Pliocene.

Type locality. “Hrastina”, Croatia.

Types. Milan et al. (1974: 58) indicated a holotype, but it is uncertain whether the specimen was the only one Brusina had at hand (holotype by monotypy, Art. 73.1.2). The specimen is stored in the Croatian Natural History Museum, Zagreb, coll. no. 2481-127.

Remarks. Appeared first as a nomen nudum in Gorjanović-Kramberger (1892: 109).

Melanella krapinensis Bourguignat, 1884

Original source. Bourguignat 1884: 18.

Type locality. “Dans le canal de sortie des eaux thermales de Krapina-Toeplitz, en Croatie” [in the outlet channel of the thermal waters at Krapinske toplice], Croatia.

Remarks. Bourguignat denoted the authority as “Letourneux, 1879”, but there is no evidence that the description really derived from that author.

† *Melanopsis kupensis* Fuchs, 1870

Original source. Fuchs 1870b: 544, pl. 22, figs 3–4.

Type horizon. Middle Pannonian, late Miocene.

Type locality. “Kúp”, Hungary.

† *Melanopsis kurdica* Brusina, 1902

Original source. Brusina 1902: pl. 7, figs 8–11.

Type horizon. Portaferrian (Pannonian Basin), late Miocene–Pliocene.

Type locality. “Kurd”, Hungary.

Types. The illustrated syntypes are stored in the Croatian Natural History Museum, Zagreb, coll. no. 2535-181/1-4 (Milan et al. 1974: 93).

† *Melanopsis kuzmici* Brusina, 1907

Original source. Brusina 1907: 197.

Type horizon. Late Burdigalian–early Langhian, early–middle Miocene.

Type locality. Not indicated by Brusina (1905, 1907), apart from “probably Sinj Basin”, Croatia.

Remarks. Appeared first as a nomen nudum in Brusina (1905: 32).

† *Melanopsis lactacea* Cossmann, 1888

Original source. Cossmann 1888: 283, pl. 11, figs 12–13.

Type horizon. Thanetian, Paleocene.

Type locality. “Chenay [...]; Jonchery [...]; Châlons-sur-Vesle”, France.

Melanopsis laevigata Lamarck, 1822

Original source. Lamarck 1822: 168.

Type locality. “Dans les rivières des îles de l’Archipel” [it is unknown which Archipel Lamarck referred to].

Remarks. The species first appeared without name on a plate in the “Tableau encyclopédique et méthodique, vol. 3” by Lamarck (1797). The captions for most plates of the three volumes work were prepared by Bory de Saint-Vincent, but apparently not before 1824, and added to the “Tableau encyclopédique et méthodique, vol. 1” of Bruguière (1791; see note on pp. 83–84). In the captions, Bory de Saint-Vincent also referred to the “Histoire naturelle” (vol. 6, part 2) by Lamarck (1822), where the name first occurred.

***Melania holandri* var. *laevigata* Rossmässler, 1839**

Original source. Rossmässler 1838–1844: Heft 4 (10), 37, pl. 50, figs 664–667.

Type locality. “Gradaschza und Ringelsza [bei Laibach]; aus einem Mühlbache bei Nassenfuss in Unterkrain” [in Gradaščica and Ringelsza (?) brooks near Ljubljana; from a mill creek near Mokronog] (the remaining localities mentioned cannot be assigned to the present variety without doubt), Slovenia.

Remarks. Note that Rossmässler (1839) seems to have listed *M. holandri* in synonymy of this variety, but from the text it is clear that the list is intended to refer to the species as a whole.

† *Melanopsis laevigata* Łomnicki, 1886 [invalid]

Original source. Łomnicki 1886: 76.

Type horizon. Late Miocene.

Type locality. “Wyczółki (wrzynka kol. trans. na zachodnio-południowym końcu wsi)” [Goncharivka, at southwestern end of village], Ukraine.

Remarks. Junior homonym of *Melanopsis laevigata* Lamarck, 1822. Wenz (1928a: 119) introduced *M. lomnickii* as replacement name.

***Melanopsis tunetana* var. *laevigata* Pallary, 1912 [invalid]**

Original source. Pallary 1912: 20, [unnumbered plate], figs 31–33.

Type locality. “Nefta”, Tunisia.

Remarks. Junior homonym of *Melanopsis laevigata* Lamarck, 1822.

***Melanopsis penchinati* var. *laevigata* Pallary, 1924 [invalid]**

Original source. Pallary 1924: 248, pl. 15, fig. 12.

Type locality. “Eaux de Mathen, Alhama de Aragon” [waters of Mathen (?), Alhama de Aragón], Spain.

Remarks. Junior homonym of *Melanopsis laevigata* Lamarck, 1822. The taxon appears at infrasubspecific rank in the plate captions of the original publication.

† *Melanopsis laevis* Stoliczka, 1860

Original source. Stoliczka 1860: 484, pl. 1, figs 4a–b.

Type horizon. Late Turonian, late Cretaceous.

Type locality. “Neualpe im Russbachthal” [Neualm near Russbach am Pass Gschütt], Austria.

***Melanopsis vespertina* var. *laevis* Bourguignat, 1884 [invalid]**

Original source. Bourguignat 1884: 125.

Type locality. “Ruisseau de l’île d’Ivice, aux Baléares” [stream on the island of Ibiza], Spain.

Remarks. Junior homonym of *Melanopsis laevis* Stoliczka, 1860.

***Melanopsis douttei* var. *laevis* Pallary, 1920 [invalid]**

Original source. Pallary 1920c: 150.

Type locality. “Oued Masmouda; Ruisseau de Bab Hadid” [Oued Masmouda (?); stream Bab Hadid (?)], Morocco.

Remarks. Junior homonym of *Melanopsis laevis* Stoliczka, 1860.

***Melanopsis foleyi* var. *laevis* Pallary, 1928 [invalid]**

Original source. Pallary 1928a: 270.

Type locality. “Aïn Mélias, près de Figuig” [Ain Melias near Figuig], Algeria.

Remarks. Junior homonym of *Melanopsis laevis* Stoliczka, 1860.

***Melanopsis lamarckii* Potiez & Michaud, 1838**

Original source. Potiez and Michaud 1838: 351, pl. 31, figs 5–6.

Type locality. “Les rivières de Madagascar?” [rivers of Madagascar?].

Remarks. Köhler and Glaubrecht (2010: 873) considered the taxon as a junior synonym of *Madagasikara spinosa* (Lamarck, 1822) (Pachychilidae).

† *Melanopsis lamarckii* Deshayes, 1862 [invalid]

Original source. Deshayes 1861–1864: 472, pl. 31, figs 25–26.

Type horizon. Calcaire de Montabuzard, middle–late Burdigalian.

Type locality. “Damery”, France.

Remarks. Junior homonym of *Melanopsis lamarckii* Potiez & Michaud, 1838. Cossmann (1888: 279) and Wenz (1929: 2640) classified the species in the genus *Faunus* Montfort, 1810 (Pachychilidae). The name “*lamarcki*” as mentioned in Wenz is an incorrect subsequent spelling.

***Melanopsis lamberti* Souverbie, 1872**

Original source. Souverbie 1872: 148.

Type locality. “Baye du Sud” [Baie Sud], New Caledonia.

Remarks. Brot (1879: 463) considered the taxon as a junior synonym of *Melanopsis mariei* Crosse, 1869.

***Melanopsis olivieri* var. *lamellata* Bourguignat, 1884**

Original source. Bourguignat 1884: 98.

Type locality. “Environs de Constantinople” [surroundings of Istanbul], Turkey.

***Melanopsis lampra* Bourguignat, 1884**

Original source. Bourguignat 1884: 132.

Type locality. “Belus, près de Saint-Jean-d’Acre, en Syrie” [in the Na’aman river, near Acre], Israel.

Remarks. Heller et al. (2002: 596) considered the species as a junior synonym of *Melanopsis costata* (Olivier, 1804). Heller et al. (2005: 244) in turn treated it as an accepted name.

† *Melanopsis lanceolata* Neumayr in Neumayr & Paul, 1875

Original source. Neumayr and Paul 1875: 39, pl. 7, figs 5, 15.

Type horizon. Cernikian, Pliocene.

Type locality. “Slobodnica; [...] Sibir [Sibirj]; [...] Malino; [...] Gromačnik; [...] Čapla [Čaplja trench near Slavonski Brod]; [...] Cigelnik”, Croatia.

† *Melanopsis (Melanoptychia) langhofferi* Pavlović, 1927

Original source. Pavlović 1927: 61, pl. 7, figs 7–8.

Type horizon. Middle Pannonian, late Miocene.

Type locality. “Из каратачких neckoџа” [from the sands of Karagača near Vrčin], Serbia.

Types. The illustrated syntype is stored in the Natural History Museum, Belgrade, coll. no. 211 (Milošević 1962: 23).

† *Melanopsis lanzaeana* Brusina, 1874

Original source. Brusina 1874: 34.

Type horizon. Langhian, middle Miocene.

Type locality. “Ribarić; Turjake [Turjaci]”, Croatia.

Types. Milan et al. (1974: 94) indicated a holotype, but it is uncertain whether the specimen is part of the original type series and whether it was the only one Brusina had at hand (holotype by monotypy, Art. 73.1.2). The specimen is stored in the Croatian Natural History Museum, Zagreb, coll. no. 2982-628.

Remarks. The name “*lanzae*” as mentioned in Olujić (1999: 20, 49) is an incorrect subsequent spelling.

Melanopsis latastei Letourneux & Bourguignat, 1887

Original source. Letourneux and Bourguignat 1887: 158.

Type locality. “Des environs de Nefta; [...] dans l’Oued Gabès et aux alentours de Tozer” [surroundings of Nefta; in the Oued Gabès and around Tozeur], Tunisia.

† *Melanopsis laubrierei* Carez, 1879

Original source. Carez 1879: 637, pl. 12, figs 1–5.

Type horizon. Early–middle Eocene.

Type locality. “La Maladrerie (Brasles); Gland (Aisne)”, France.

† *Melanopsis lebedai* Lueger, 1980

Original source. Lueger 1980: 104.

Type horizon. Pannonian, zone D, late Miocene.

Type locality. “Föllig d1 [...] und d2” [Föllig hill near Großhöflein], Austria.

Types. Geological-Paleontological Department, Natural History Museum Vienna, Austria. The number Lueger provided is not the collection number but the acquisition number, which could be used to trace back the object in our database.

Melanopsis lecointrei Pallary, 1918

Original source. Pallary 1918: 150.

Type locality. “Taza, Ain en nsa, source est, chaude” [Taza, in the hot spring Ain en nsa], Morocco.

***Melania holandri* var. *legitima* Rossmässler, 1839**

Original source. Rossmässler 1838–1844: Heft 4 (10), 37, pl. 50, figs 662–663.

Type locality. “In der Laibach; in der Save” [Ljubljanska and Sava rivers] (the remaining localities mentioned cannot be assigned to the present variety without doubt), Slovenia.

“*Melanopsis leMBERGENSIS*” mentioned in Brot (1870: 310) and Brot (1874: 370) [unavailable]

Locality. “Lemberg, Galizien” [Lviv], Ukraine.

Remarks. Nomen nudum, apparently based on an unused manuscript name. It appears only in the synonymy list of *Hemisinus* [now *Esperiana*] *acicularis* in Brot (1870) with the authority “Schröter” as well in Brot (1874) with the authority “Parreyss”.

***Melanopsis lentiginosa* Reeve, 1860**

Original source. Reeve 1860: Section *Melanopsis*, pl. 3, figs 9a–b.

Type locality. “New Caledonia” (France) [no locality indicated].

Remarks. Brot (1879: 444) considered the taxon as a junior synonym of *Melanopsis frustulum* Morelet, 1857.

† *Melanopsis (Martinia) leobersdorfensis* Handmann, 1887

Original source. Handmann 1887: 23, pl. 3, fig. 10.

Type horizon. Pannonian, zone B–D, late Miocene.

Type locality. “Leobersdorf”, Austria.

Remarks. Wenz (1929: 2719) considered the taxon as a junior synonym of *Melanopsis fossilis* (Gmelin, 1791).

† *Melanopsis lepavinensis* Brusina, 1897

Original source. Brusina 1897: 9, pl. 6, figs 17–20.

Type horizon. *Congerina rhomboidea* Zone, Portaferrian, late Miocene.

Type locality. “Lepavina”, Croatia.

Types. Milan et al. (1974: 94) indicated a holotype, but it is uncertain whether the specimen is part of the original type series and whether it was the only one Brusina had at hand (holotype by monotypy, Art. 73.1.2). The specimen is stored in the Croatian Natural History Museum, Zagreb, coll. no. 3003-649.

***Fagotia letourneuxi* Bourguignat, 1884**

Original source. Bourguignat 1884: 45.

Type locality. “Rivière entre Plaski et Ostaria, et dans la Save entre Agram et Sissek” [river between Plaški and Oštarije, and in the Sava river between Zagreb and Sisak], Croatia.

Remarks. Note that Bourguignat denoted the authority as “Bourguignat, 1879”. Starobogatov et al. (1992: 60) considered the species as a junior synonym of *Fagotia* [= *Esperiana*] *acroxia* Bourguignat, 1884

***Melanella letourneuxi* Bourguignat, 1884**

Original source. Bourguignat 1884: 25.

Type locality. “Dans la rivière d’Ogulin, en Croatie; [...] dans la Migliaska, près de Séràjewo” [in the river at Ogulin (Croatia); in the Miljacka river near Sarajevo (Bosnia and Herzegovina)].

Remarks. Appeared first as a nomen nudum in Servain (1884: 379) [January]. Bourguignat (1884) [May] denoted the authority as “Bourguignat, 1879”.

***Melanopsis letourneuxi* Bourguignat, 1884**

Original source. Bourguignat 1884: 116.

Type locality. “Dans la source et la rivière de la Moulouiah, à l’ouest de Lalla-Maghnia” [in the source and the river Moulouya, west of Maghnia], Morocco or Algeria.

Remarks. Bourguignat denoted the authority as “Bourguignat, 1872, et Letourneux”, apparently referring to Letourneux (1876b). That work is not available to me, but after Crosse (1877: 33) that record is a nomen nudum. It appears also as nomen nudum in Letourneux (1876a: 48). In addition, Bourguignat listed “Bourguignat. Spec. nov. Moll., no. 198, 1878”, which displays the second part of “Species novissimae Molluscorum” (Bourguignat 1876), but which was never published.

***Microcolpia letourneuxi* Bourguignat, 1884**

Original source. Bourguignat 1884: 64.

Type locality. “La Save, au-dessous d’Agram” [Sava river below Zagreb], Croatia.

***Melanopsis crenocarinata* var. *leucostoma* Moricand, 1841**

Original source. Moricand 1841: 62.

Type locality. “Rio de Pedra Branca, procince de Bahia” [Pedra Branca river, province Bahia], Brazil.

Remarks. Although not explicitly stated, this variety was apparently considered to belong to the new genus *Verena* by Adams and Adams (1854) (Thiaridae), of which *Melanopsis crenocarinata* is the type species (see Nuttall 1990: 253).

† *Melanopsis libanensis* Delpey, 1940

Original source. Delpey 1940: 93, fig. 66, pl. 5, fig. 5.

Type horizon. Aptian, early Cretaceous.

Type locality. “Bekfaya; [...] Mar Abda” [Bikfaiya; Mâr Abdâ], Lebanon.

Remarks. Probably not a Melanopsidae.

† *Melanopsis liburnica* Stache, 1889

Original source. Stache 1889: 145, pl. 3, figs 25–27, pl. 5, fig. 12.

Type horizon. Danian, Paleocene.

Type locality. “Corgnale, Divacca und unteres Gaberg-Gehänge” [Lokev, Divača, and lower slopes of Mt. Vremščica], Italy.

Remarks. Appeared first as a nomen nudum in Stache (1880: 199).

† *Hemisinus lignitarius* Tausch, 1886

Original source. Tausch 1886: 8, pl. 1, figs 24–27.

Type horizon. Late Santonian–early Campanian, late Cretaceous.

Type locality. “Ajka”, Hungary.

Remarks. Oppenheim (1892: 757) and Bandel and Riedel (1994: 20) considered the species as a junior synonym of *Esperiana obeloides* (Tausch, 1886).

† *Melanopsis limbata* Pallary, 1916

Original source. Pallary 1916: 83.

Type horizon. Pannonian, zone D, late Miocene.

Type locality. “Kottingbrunn [...] Ziegelei a” (Handmann 1882: 559), Austria.

Remarks. Replacement name for *M. scalaris* Handmann, 1882, non Gassies, 1856. Wenz (1929: 2741) considered the taxon as a junior synonym of *Melanopsis haueri* Handmann, 1882.

***Melanopsis costata* var. *lineata* Pallary, 1939**

Original source. Pallary 1939: 91, pl. 5, fig. 33.

Type locality. “Djishr ech Chegour” [Jisr Ash-Shughur], Syria.

Remarks. Pallary erroneously gave the name as “*unicincta*” in the plate captions. Heller and Sivan (2002a: 49) considered the taxon as a junior synonym of *Melanopsis multiformis* Blanckenhorn, 1897.

† ***Pseudofagotia lineata* Anistratenko, 1993**

Original source. Anistratenko 1993: 72, textfig. 2.

Type horizon. Duab Beds, middle to late Kimmerian, Pliocene.

Type locality. “Окр. с. Мокбу, Очамчирский р-н” [near the village Mok’vi, Ochamchirskiy rayon], Georgia.

Types. Schmalhausen Institute of Zoology of National Academy of Sciences of Ukraine, Kiev; no number indicated.

Remarks. Type species of the genus *Pseudofagotia* Anistratenko, 1993.

***Melanopsis lineolata* Gassies, 1857**

Original source. Gassies 1857: 276, pl. 9, figs 9–10.

Type locality. “La rivière Balade” [in the river Balade], New Caledonia.

Remarks. Gassies (1861) apparently considered *Melania lineolata* Gray in Griffith & Pidgeon, 1833 as a *Melanopsis*, which resulted in secondary homonymy for his species *Melanopsis lineolata* Gassies, 1857. Thus, he introduced *Melanopsis livida* as a replacement name. Secondary homonymy, however, is not given anymore, since *Melania lineolata* Gray was recently shown to belong to the genus *Cerithidea* Swainson, 1840 (Potamididae) by Reid (2014). (Note, moreover, that *Melania lineolata* Gray is a junior homonym of *Melania lineolata* Wood, 1828). *M. livida* is a junior objective synonym of *Melanopsis lineolata* Gassies (see also Art. 59.3). Brot (1879: 444) considered the taxon as a junior synonym of *Melanopsis frustulum* Morelet, 1857.

***Melanopsis liocephala* Pallary, 1936**

Original source. Pallary 1936: 55, pl. 3, fig. 5.

Type locality. “Guefäit (Maroc oriental)”, Morocco.

***Melanopsis lirata* Gassies, 1869**

Original source. Gassies 1869: 77.

Type locality. “Prope Noumea” [near Nouméa], New Caledonia.

Remarks. Brot (1879: 444) considered the taxon as a junior synonym of *Melanopsis frustulum* Morelet, 1857.

***Melanopsis livida* Gassies, 1861 [invalid]**

Original source. Gassies 1861: 290, pl. 7, fig. 9.

Type locality. “Le Diahot à Balade” [in the Diahot river at Balade], New Caledonia.

Remarks. Replacement name for the presumed secondary homonym *Melanopsis lineolata* Gassies, 1856, non *Melania lineolata* Gray in Griffith & Pidgeon, 1833. Secondary homonymy, however, is not given anymore, since *Melania lineolata* Gray was recently shown to belong to the genus *Cerithidea* Swainson, 1840 (Potamididae) by Reid (2014) (see also Art. 59.3). (Note, moreover, that *Melania lineolata* Gray is a junior homonym of *Melania lineolata* Wood, 1828). *M. livida* is a junior objective synonym of *Melanopsis lineolata* Gassies. Brot (1879: 444) considered the taxon as a junior synonym of *Melanopsis frustulum* Morelet, 1857.

***Melanopsis locardi* Blanckenhorn, 1897**

Original source. Blanckenhorn 1897: 132, pl. 10, fig. 15.

Type horizon. Pleistocene (?)–Recent.

Type locality. “Lebend im See von Antiochia. Halbfossil bei Selemije” [living in Lake Anuk (also as Amik) (Turkey); subfossil near As Salamīyah (Syria)].

***Fagotia locardiana* Bourguignat, 1884**

Original source. Bourguignat 1884: 39.

Type locality. “Dans le lac Sabandja, près d’Ismidt (Anatolie)” [Lake Sapanca near İzmit], Turkey.

Remarks. Note that Bourguignat denoted the authority as “Bourguignat, 1882”.

† *Melanopsis lomnickii* Wenz, 1928

Original source. Wenz 1928a: 119.

Type horizon. Badenian, middle Miocene.

Type locality. “Wyczółki (wrzynka kol. trans. na zachodnio-południowym końcu wsi)” (Łomnicki 1886: 76) [Goncharivka, at southwestern end of village], Ukraine.

Remarks. Replacement name for *M. laevigata* Łomnicki, 1886, non Lamarck, 1792.

† *Melanopsis nodosa* var. *longa* Deshayes in Férussac, 1839

Original source. Férussac 1819–1851: *Mélanopsides fossiles*, pl. 2, fig. 8 (altered caption by Deshayes delivered with Livraison 29; see introduction for details).

Type horizon. Late Miocene.

Type locality. “D’Athènes” [Athens], Greece.

† *Microcolpia longa* Gozhik in Gozhik & Datsenko, 2007

Original source. Gozhik and Datsenko 2007: 95, pl. 95, figs 1–5, pl. 97, fig. 3.

Type horizon. Late Pleistocene.

Type locality. “Аллювія V террасы р. Дунай у с. Назорное” [Alluvial terrace V of the Danube river near Nagornoye], Ukraine.

Types. Institute of Geological Sciences of the National Academy of Sciences of Ukraine, coll. no. 6461.

Remarks. Originally the gender was indicated as masculine (“*longus*”), but *Microcolpia* is feminine, which is why the name must be “*longa*”.

† *Melanopsis (Lyrcea) senatoria* var. *longata* Handmann, 1887

Original source. Handmann 1887: 19, pl. 2, figs 12–13.

Type horizon. Pannonian, zone B–D, late Miocene.

Type locality. “Leobersdorf”, Austria.

† *Melanopsis longirostris* Pallary, 1920

Original source. Pallary 1920b: 112.

Type horizon. Pontian (sensu stricto), late Miocene.

Type locality. “Babadjan” (Andrusov 1909: captions of pl. 5) [Babadzhan], Azerbaijan.

Remarks. Based on the record of “*Melanopsis* sp.” in Andrusov (1909: 88, pl. 5, figs 1–2).

***Melanopsis cesari* var. *longirostris* Pallary, 1928 [invalid]**

Original source. Pallary 1928a: 255.

Type locality. “Dans les séguias de Beni Abbès” [in the irrigation channel of Beni Abbès], Algeria.

Remarks. Junior homonym of *Melanopsis longirostris* Pallary, 1920.

† *Melanopsis (Canthidomus) macrosculpturata longitesta* Papp, 1953

Original source. Papp 1953a: 110, pl. 23, figs 16–18.

Type horizon. Gelasian, early Pleistocene.

Type locality. “Pyrgos (Elis)”, Greece.

Types. Museum of Palaeontology and Geology of the University of Athens; no number indicated.

† *Melanopsis praemorsa* var. *longopyrulata* Sacco, 1895

Original source. Sacco 1895: 9, pl. 1, fig. 15.

Type horizon. Early Messinian, late Miocene.

Type locality. “S. Marzano Oliveto”, Italy.

Remarks. Wenz (1929: 2733) considered the taxon as a junior synonym of *Melanopsis fusulatina* Sacco, 1895.

***Melanopsis lorcana* Guirao, 1854**

Original source. Guirao 1854: 32.

Type locality. “In rivulo Rambla de Viznaga et in Pantano de Puentes non procul Lorca in Regno Murcico” [in the river Rambla de Viznaga and in Pantano de Puentes, not far from Lorca, prov. Murcia], Spain.

† *Melanopsis (Canthidomus) lorentheyi* Andrusov, 1909

Original source. Andrusov 1909: 82, 158, pl. 4, figs 11–20.

Type horizon. Pontian (sensu stricto), late Miocene.

Type locality. “Babadjan” [Babadzhan], Azerbaijan.

† *Melanopsis lorioli* Locard, 1893

Original source. Locard 1893: 187, pl. 9, fig. 22.

Type horizon. Middle–late Burdigalian, early Miocene.

Type locality. “Vernier, près Genève”, Switzerland.

Melanopsis lortetiana Locard, 1883

Original source. Locard 1883a: 271, pl. 23, figs 50–51.

Type locality. “Lac d’Antioche” [Lake Anuk (also as Amik)], Turkey.

Remarks. Appears as “*Lorteti*” in Locard’s remarks on p. 272, which is apparently based on a typesetting error.

† *Melanopsis lozanici* Brusina, 1893

Original source. Brusina 1893: 201 (Serbian part), 34 (Italian part), pl. 2, fig. 6.

Type horizon. Pannonian, zone D–E, late Miocene.

Type locality. “Ripanj”, Serbia.

Types. Milan et al. (1974: 94) indicated a holotype, but it is uncertain whether the specimen is part of the original type series and whether it was the only one Brusina had at hand (holotype by monotypy, Art. 73.1.2). The specimen is stored in the Croatian Natural History Museum, Zagreb, coll. no. 3639-1279/1.

† *Melanopsis lucanensis* Neubauer in Neubauer et al., 2011

Original source. Neubauer et al. 2011: 206, pl. 1, figs 10–12.

Type horizon. Early Langhian, middle Miocene.

Type locality. “Sinj, Lučane [= Sutina] section”, Croatia.

Types. Geological-Paleontological Department, Natural History Museum Vienna, Austria, coll. no. 2010/0042/0001.

† ?*Amphimelania lucida* Cossmann, 1886

Original source. Cossmann 1886: 228, pl. 10, fig. 7.

Type horizon. Eocene.

Type locality. “Houdan”, France.

Remarks. This taxon was considered to belong to the marine genus *Nozeba* Iredale, 1915 (Iravadiidae) by Ponder (1984: 56).

“*Melanopsis lucio*” mentioned in Erber (1868: 904) [unavailable]

Locality. “Rhodus” [Rhodes island, no locality indicated], Greece.

Remarks. Nomen nudum. Erber attributed the authority to Mousson. Given as “*Luciae*” in Paetel (1888).

***Melanopsis dircaeana* var. *luctuosa* Pallary, 1939**

Original source. Pallary 1939: 88, pl. 4, fig. 19, pl. 6, fig. 35.

Type locality. “Lac de Homs”, Syria.

† *Melanopsis ludwigi* Speyer, 1870

Original source. Speyer 1870: 97, pl. 15, figs 3–4.

Type horizon. Early Rupelian, Oligocene.

Type locality. “Grossalmerode” (Ludwig 1865: 71), Germany.

Remarks. Replacement name for *M. costata* Ludwig, 1865, non Olivier, 1804.

† *Melanopsis lushani* d’Archiac in Viquesnel, 1846

Original source. Viquesnel 1846: 265, pl. 16, fig. 1.

Type horizon. Pannonian, late Miocene.

Type locality. “Entre Koulana et Lus-han” [between Koulana and Lus-han, S of Drin river mouth], Albania.

Remarks. Originally the name was introduced as “*Lus-Hani*”; the name “*Lush-ami*” as mentioned in Chenu (1859: 297) is an incorrect subsequent spelling. Wenz (1929: 2715) considered the taxon as a junior synonym of *M. fossilis* (Gmelin, 1791).

***Melanopsis iraqensis* var. *lutea* Pallary, 1939**

Original source. Pallary 1939: 89.

Type locality. “Ain Haglan” [not found], Iraq.

***Melanopsis (Sistaniana) lutensis* Starobogatov & Izzatullaev, 1985**

Original source. Starobogatov and Izzatullaev 1985: 35, fig. 5.

Type locality. “Пустыне Дешме-Лым” [Dasht-e Loot desert], Iran.

Types. Zoological Institute of Russian Academy of Sciences, St.-Petersburg; no number indicated.

***Melanopsis costata* var. *luteopsis* Germain, 1921**

Original source. Germain 1921: 494, pl. 20, figs 9–10.

Type locality. “Lac d’Homs”, Syria.

Remarks. Heller et al. (2005: 248) considered the variety as a junior synonym of *Melanopsis saulcyi* Bourguignat, 1853.

† ***Melanopsis lyra* Matheron, 1842**

Original source. Matheron 1842: 293, pl. 37, figs 8–10.

Type horizon. Early Campanian, Cretaceous.

Type locality. “Les Martigues”, France.

Remarks. Sandberger (1871: 88) attributed the species to the genus *Paludomus* Swainson, 1840 (Paludomidae).

† ***Melanopsis (Canthidomus) lyrata* Neumayr, 1869 [invalid]**

Original source. Neumayr 1869: 358, pl. 11, fig. 8.

Type horizon. Langhian, middle Miocene.

Type locality. “Ribaric” [Ribarić], Croatia.

Types. Illustrated syntype is stored at the Geological Survey Austria, Vienna, coll. no. 1869/01/6.

Remarks. Unlike stated by Neubauer et al. (2016b: 25), the name *M. lyrata* Neumayr, 1869 [June] is deemed to be a junior homonym of *M. lirata* Gassies, 1869 [January] after Art. 58.2. Pallary (1916: 85) introduced *M. dissimilis* as replacement name. Wenz (1929: 2698) was aware of the homonymy issue and synonymized this species with *M. cylindracea* Brusina, 1874.

***Microcolpia mabilliana* Bourguignat, 1884**

Original source. Bourguignat 1884: 59.

Type locality. “La rivière entre Plaski et Ostaria (Croatie)” [river between Plaški and Oštarije], Croatia.

Remarks. Starobogatov et al. (1992: 65) considered the species as a junior synonym of *Microcolpia acicularis* (Férussac, 1823).

† ***Melania macedonica* Burgerstein, 1877**

Original source. Burgerstein 1877: 248, pl. 3, figs 13–16.

Type horizon. Late Miocene.

Type locality. “Ueskueb” [Skopje], Macedonia (not Zvezdan in Serbia as given by Wenz 1929).

Remarks. Considered to belong in the genus *Amphimelania* by Brusina (1893: 63) and Wenz (1929: 2875).

Melanopsis macilenta Pallary, 1928

Original source. Pallary 1928b: 17, pl. 2, figs 15–17.

Type locality. “Beni Mellal, dans l’oued Taguenout” [Beni Mellal, in the Oued Taguenout (?)], Morocco.

“*Melania macilenta* Ziegler” mentioned in Brot (1874–1879: 13) [unavailable]

Locality. “In der Muhr” [in the river Mur], Austria.

Remarks. Nomen nudum, apparently based on an unused manuscript name from Parreyss listed in synonymy of “*Melania Holandri*” [sic] by Brot (1874).

† *Melanopsis (Canthidomus) macrosculpturata* Papp, 1953

Original source. Papp 1953a: 109, pl. 23, figs 13–18.

Type horizon. Gelasian, early Pleistocene.

Type locality. “Pyrgos”, Greece.

Types. Museum of Palaeontology and Geology of the University of Athens; no number indicated.

Melanopsis macrostoma Bourguignat, 1884

Original source. Bourguignat 1884: 157.

Type locality. “Le Guadalquivir aux environs de Séville” [in the Guadalquivir river, around Sevilla], Spain.

Melanopsis maculata Lea, 1837

Original source. Lea 1837: 82, pl. 19, fig. 75.

Type locality. “Peru” [no locality indicated].

Remarks. Junior synonym of *Hemisinus osculati* (Villa in Villa & Villa, 1854) (Thiaridae) after Ramírez et al. (2003: 272).

† *Melanopsis magna* Pallary, 1916 [invalid]

Original source. Pallary 1916: 78.

Type horizon. Burdigalian, early Miocene (?).

Type locality. “Dax”, France.

Remarks. Pallary (1916) erroneously treated “*magna*” as available name attributed to Férussac (1823), who solely used the word as descriptive term (Latin “big”), which he applied to many of his varieties. The name became nevertheless available from Pallary (1916) who associated the name with an illustration in Férussac (1823: pl. 7, fig. 16). The illustrated specimen is, however, the holotype (by monotypy) of *M. dufourii* Férussac, 1822 (Férussac used the same plates in both of his works and thus Pallary actually referred to the specimen illustrated as *M. dufourii* in the “Histoire naturelle”; see also introduction and Table 1 for details of Férussac’s publications). Therefore, *Melanopsis magna* Pallary, 1916 is a junior objective synonym of *M. dufourii*.

Melanopsis letourneuxi var. *magna* Pallary, 1928 [invalid]

Original source. Pallary 1928a: 273, pl. 5, fig. 6.

Type locality. “Berguent” [Aïn Beni Mathar], Morocco.

Remarks. Junior homonym of *Melanopsis magna* Pallary, 1916.

† *Melanopsis (Lyrcaea)* [sic] *magna* Lubenescu, 1985 [invalid]

Original source. Lubenescu 1985: 78, pl. 2, fig. 8.

Type horizon. Middle Pannonian, late Miocene.

Type locality. “Village Cut, district Alba”, Romania.

Types. Institute of Geology and Geophysics, University of Bucharest, coll. no. 14655.

Remarks. Originally the gender was indicated as masculine (“*magnus*”), but *Melanopsis* is feminine, which is why the name must be “*magna*”. Junior homonym of *Melanopsis magna* Pallary, 1916.

Melanopsis magnifica Bourguignat, 1884

Original source. Bourguignat 1884: 152.

Type locality. “Environs de Fez” [surroundings of Fes], Morocco.

† *Melanopsis magyari* Neubauer, Harzhauser, Georgopoulou, Mandic & Kroh, 2014

Original source. Neubauer et al. 2014a: 457.

Type horizon. Middle Pannonian, late Miocene.

Type locality. “Begaljica” (Brusina 1897: 8), Serbia.

Types. Milan et al. (1974: 93) indicated a holotype, but it is uncertain whether the specimen was the only one Brusina had at hand when describing *M. klerici inermis* (holotype by monotypy, Art. 73.1.2). The specimen is stored in the Croatian Natural History Museum, Zagreb, coll. no. 3020-666.

Remarks. Replacement name for *M. klerici inermis* Brusina, 1897, non Handmann, 1882 (see Note 1).

† “*Melanopsis major* Férussac, 1823” mentioned in Wenz (1929: 2773) [unavailable]

Horizon. Burdigalian, early Miocene.

Locality. “De Mandillot, près de Dax” [Mandillot, near Dax], France.

Remarks. This name has appeared several times in the literature (e.g., Bronn 1848: 717; Wenz 1929: 2773), but it is based on an error. Férussac (1823: 154) actually described *Melanopsis dufourii* var. ϵ and listed “*Fossilis, major*. Férussac” in synonymy referring to plate 1 of the “*Mélanopsides fossiles*” of his “*Histoire naturelle*” (Férussac 1819–1851; see also introduction for details about this work). However, in the captions of plate 1, it is given as “*Melanops. Dufourii*, var. ϵ). *Fossilis, maxima*”. Obviously neither *major* nor *maxima* was intended as species-group name by Férussac. Later, the name would have been a junior homonym of *M. buccinoidea major* Grateloup, 1838.

† *Melanopsis buccinoidea* var. *major* Grateloup, 1838

Original source. Grateloup 1838: 146, pl. 4, figs 52–53.

Type horizon. Burdigalian, early Miocene.

Type locality. “Dax. [...] Mandillot; à Saint-Paul”, France.

Melanopsis acicularis var. *major* Rossmässler, 1839 [invalid]

Original source. Rossmässler 1838–1844: 41, pl. 50, figs 673–675.

Type locality. “In der Lachina bei Tschernembl, [...] in dem Bug” [in the Lahinja river near Črnomelj (Slovenia) and in the Bug river (Ukraine)].

Remarks. Junior homonym of *Melanopsis buccinoidea major* Grateloup, 1838 (see Note 1).

Melanopsis maroccana var. *major* Bourguignat, 1864 [invalid]

Original source. Bourguignat 1864: 260.

Type locality. “Mostaghanem” [Mostaganem], Algeria.

Remarks. Junior homonym of *Melanopsis buccinoidea major* Grateloup, 1838 (see Note 1).

***Melanopsis cariosa* var. *major* Bourguignat, 1884 [invalid]**

Original source. Bourguignat 1884: 151.

Type locality. “Dans les aqueducs de Séville et dans le Guadalquivir” [in the aqueducts of Sevilla and in the Guadalquivir river], Spain.

Remarks. Junior homonym of *Melanopsis buccinoidea major* Grateloup, 1838 (see Note 1).

***Melanopsis macrostoma* var. *major* Bourguignat, 1884 [invalid]**

Original source. Bourguignat 1884: 157.

Type locality. “Le Guadalquivir aux environs de Séville” [in the Guadalquivir river, around Sevilla], Spain.

Remarks. Junior homonym of *Melanopsis buccinoidea major* Grateloup, 1838 (see Note 1).

***Melanopsis subscalaris* var. *major* Bourguignat, 1884 [invalid]**

Original source. Bourguignat 1884: 108.

Type locality. “Environs de Fez (Maroc)” [surroundings of Fes], Morocco.

Remarks. Junior homonym of *Melanopsis buccinoidea major* Grateloup, 1838 (see Note 1).

***Melanopsis pseudoferussaci* var. *major* Pallary, 1899 [invalid]**

Original source. Pallary 1899: 140.

Type locality. “Environs de Tétouan” [surroundings of Tétouan], Morocco.

Remarks. Junior homonym of *Melanopsis buccinoidea major* Grateloup, 1838 (see Note 1).

***Melanopsis tingitana* var. *major* Pallary, 1899 [invalid]**

Original source. Pallary 1899: 147.

Type locality. “L’O.[ued] Ida ou Guert, près de Mogador (p.); O. Aït Ouadel (p. 163)” [river (?) Ida Ou Gourdh at Essaouira;], Morocco.

Remarks. Junior homonym of *Melanopsis buccinoidea major* Grateloup, 1838 (see Note 1).

† *Melanopsis hammamensis* var. *major* Pallary, 1901 [invalid]

Original source. Pallary 1901a: 178, pl. 2, fig. 21.

Type horizon. Pleistocene.

Type locality. “De l’oued El Biod (Géryville)” [Oudeï el Biod near El Bayadh], Algeria.

Remarks. Junior homonym of *Melanopsis buccinoidea major* Grateloup, 1838 (see Note 1).

Melanopsis vespertina var. *major* Pallary, 1911 [invalid]

Original source. Pallary 1911: 130, [unnumbered plate], fig. 3.

Type locality. “Tout près d’Oudjda, à 4 kilom. S.-E., sourdent les belles sources de Sidi-Yahia qui alimentent une véritable oasis, puis la ville d’Oudjda, et vont finalement se déverser dans l’oued Isly” [near Oujda, 4 km southeast, at the sources of Sidi Yahya that feed an oasis and the city of Oujda, and ultimately will flow into the Oued Isly], Morocco.

Remarks. Junior homonym of *Melanopsis buccinoidea major* Grateloup, 1838 (see Note 1).

Melanopsis tunetana var. *major* Pallary, 1912 [invalid]

Original source. Pallary 1912: 20, fig. 30.

Type locality. “A Tozeur, sur les bords du Chott [Djerid?]” [banks of the Chott el Djérid at Tozeur], Tunisia.

Remarks. Junior homonym of *Melanopsis buccinoidea major* Grateloup, 1838 (see Note 1).

Melanopsis douttei var. *major* Pallary, 1920 [invalid]

Original source. Pallary 1920c: 150.

Type locality. “La Makina”, Morocco.

Remarks. Junior homonym of *Melanopsis buccinoidea major* Grateloup, 1838 (see Note 1).

***Melanopsis ricardi* var. *major* Pallary, 1920 [invalid]**

Original source. Pallary 1920c: 141, pl. 3, fig. 7.

Type locality. “Aïn Allou”, Morocco.

Remarks. Junior homonym of *Melanopsis buccinoidea major* Grateloup, 1838 (see Note 1).

***Melanopsis penchinati* var. *major* Pallary, 1924 [invalid]**

Original source. Pallary 1924: 248, pl. 15, fig. 10.

Type locality. “Eaux de Mathen, Alhama de Aragon” [waters of Mathen (?), Alhama de Aragón], Spain.

Remarks. Junior homonym of *Melanopsis buccinoidea major* Grateloup, 1838 (see Note 1).

***Melanopsis turrita* var. *major* Pallary, 1924 [invalid]**

Original source. Pallary 1924: 247, pl. 15, fig. 5.

Type locality. “Guadalquivir”, Spain.

Remarks. Junior homonym of *Melanopsis buccinoidea major* Grateloup, 1838 (see Note 1).

† *Melanopsis gorceixi* var. *major* Magrograssi, 1928 [invalid]

Original source. Magrograssi 1928: 259, pl. 6, fig. 12.

Type horizon. Plio-Pleistocene.

Type locality. “Coo: molto frequente in tutte e due le zone fossilifere” [Kos island: very common in both areas rich in fossils, i.e., between Antimáchei and Pýli and in the northeast of the island, near Ágios Fokás], Greece.

Remarks. Junior homonym of *Melanopsis buccinoidea major* Grateloup, 1838 (see Note 1).

***Melanopsis seurati* var. *major* Pallary, 1928 [invalid]**

Original source. Pallary 1928a: 267.

Type locality. Not explicitly stated but probably the same as for the species (“La Zousfana, à la hauteur de Figuig et à Beni Ounif” [Oued Zousfana, at the height of Figuig, and at Beni Ounif], Algeria).

Remarks. Junior homonym of *Melanopsis buccinoidea major* Grateloup, 1838 (see Note 1).

***Melanopsis subimpressa* var. *major* Pallary, 1928 [invalid]**

Original source. Pallary 1928a: 263, pl. 6, fig. 1.

Type locality. Not explicitly stated but probably the same as for the species (“Guefaït (Maroc oriental)”, Morocco).

Remarks. Junior homonym of *Melanopsis buccinoidea major* Grateloup, 1838 (see Note 1).

***Melanopsis tutulata* var. *major* Pallary, 1928 [invalid]**

Original source. Pallary 1928a: 272.

Type locality. Not stated but probably the same or partly as for the species (“Berguent; Aoûllout; Ras el Mâ de Fès; O. Chkef près Fès” [Aïn Beni Mathar, Aïn Aoûllout, Ras El Ma, Oued Aïn Chkef in Fes], Morocco).

Remarks. Junior homonym of *Melanopsis buccinoidea major* Grateloup, 1838 (see Note 1).

***Melanopsis liocephala* var. *major* Pallary, 1936 [invalid]**

Original source. Pallary 1936: 57.

Type locality. Not indicated, but probably in Morocco.

Remarks. Junior homonym of *Melanopsis buccinoidea major* Grateloup, 1838 (see Note 1).

***Melanopsis saulcyi* var. *major* Pallary, 1939 [invalid]**

Original source. Pallary 1939: 97, pl. 6, fig. 15.

Type locality. “Lac d’Antioche, lac de Homs, Nahr el Kébir, Markieh, Yeni Chehir” [Lake Anuk (also as Amik), Yenişehir (Turkey), Lake Homs, Nahr el Kebir, river Marqiyah (Syria)].

Remarks. Junior homonym of *Melanopsis buccinoidea major* Grateloup, 1838 (see Note 1).

† *Melanopsis kleini* mut. *major* Jodot, 1958 [unavailable]

Original source. Jodot 1958: 65, pl. 11, figs 2–5.

Type horizon. Middle Miocene.

Type locality. “Route de Caravaca (Murcie)” [road to Caravaca, prov. Murcia], Spain.

Remarks. Introduced as “mut. nov.” which is not ruled by the provisions of the Code.

† “*Melanopsis majoricensis*” mentioned in Hermite (1879: 184, 328) [unavailable]

Horizon. Early Eocene.

Locality. “De Binisalem et de Selva” [from Binissalem and Selva, Mallorca], Spain.

Remarks. Nomen nudum, listed by Hermite in a section called “Espèces nouvelles citées et non décrites” [= “new species identified and not described”], where he listed 18 new names that he intended to describe in the second volume of his “Études géologiques sur les îles Baléares”. That part, however, has never been published, probably because Hermite died in 1880.

† *Melanopsis malladae* Cossmann, 1906

Original source. Cossmann 1906: 149, pl. C, figs 5–7.

Type horizon. Ypresian, Eocene.

Type locality. “Perauba” [section Peralba near Àger], Spain.

† *Melanopsis mansiana* Noulet, 1854

Original source. Noulet 1854: 50.

Type horizon. Ludian?–Sannoisian, Priabonian–early Rupelian, late Eocene–early Oligocene.

Type locality. “Au Mas-Saintes-Puelles (Aude); [...] de la Massale, près de Castres; [...] à Labruguière; [...] à Saint-Genest-de-Contest, à Lautrec (Tarn)” [at Mas-Saintes-Puelles (Dép. Aude); from Massale near Castres; at Labruguière; at Saint-Genest-de-Contest, at Lautrec (Dép. Tarn)], France.

Remarks. The name “*masensis*” as mentioned in Dollfus (1906: 292) is an incorrect subsequent spelling.

† *Melanopsis maresi* Bourguignat, 1862

Original source. Bourguignat 1862: 106, pl. 6, figs 1–4.

Type horizon. Pleistocene.

Type locality. “Dans la daya de Habessa, ancien lac desséché, situé à plus de 200 lieues environ au sud d’Oran” [in the Daïa Habessa, an old, desiccated lake, located more than 800 km south of Oran], Algeria.

† *Melanopsis margili* Robles, 1975

Original source. Robles 1975: 360, pl. 2, fig. 1.

Type horizon. Mammal zone MN 13–15, late Miocene–Pliocene.

Type locality. “Fuente del Viso (Albacete)” [near Villatoya], Spain.

Types. Museo Nacional de Ciencias Naturales, Madrid, coll. no. M-435.

† *Melanopsis marginata* Braun in Walchner, 1851

Original source. Braun in Walchner 1851: 1127.

Type horizon. Early Miocene.

Type locality. “Mainzer Becken” [Mainz Basin, no locality indicated], Germany.

Remarks. Wenz (1929: 2726) considered the taxon as a junior synonym of *Melanopsis fritzei* Thomä, 1845.

Melanopsis mariei Crosse, 1869

Original source. Crosse 1869a: 69 (Latin) [January].

Type locality. “In loco ‘Baie du Sud’” [Baie Sud], New Caledonia.

Remarks. Re-described in French by Crosse (1869b: 280, pl. 8, fig. 5) [October].

† *Melanopsis haueri markusevecensis* Wenz, 1930

Original source. Wenz 1930: 65.

Type horizon. Middle Pannonian, late Miocene.

Type locality. “Markuševac” (Brusina 1902: pl. 6, figs 71–72), Croatia.

Remarks. Replacement name for *M. austriaca croatica* Brusina, 1902, non Brusina, 1884 (see Note 1).

Melanopsis maroccana Morelet, 1853

Original source. Morelet 1853: 297.

Type locality. “Prov. Oranensem” [province of Oran; no locality indicated], Morocco.

Remarks. The name was first mentioned by Chemnitz (1795) as “*Buccina Maroccana*”. However, according to Opinion 184 (ICZN 1944), species names introduced in volumes 1–11 of Martini and Chemnitz’ “Neues Systematischer Conchylien Cabinet” (1769–1795) have no status in nomenclature and the name is therefore not available from this work. The name was again listed by Férussac (1823) in synonymy of two different species, i.e., *M. dufourii* Férussac, 1822 (referring to Chemnitz 1795: pl. 210, figs 2078–2081) and *M. costellata* (referring to Chemnitz 1795: pl. 210, figs 2082–2083). A mention in synonymy, however, does not make the name available (see Note 2).

The first to adopt *M. maroccana* as a valid name was Morelet (1853). (Note that in the same year Bourguignat mentioned the name in his “Catalogue raisonné des

mollusques terrestres et fluviatiles [...]”, which, according to the preface was published not before 1 December 1853, while Morelet’s paper was issued on 1 August 1853.) Morelet (as well as Bourguignat) referred to Chemnitz’ work – but only to figs 2080–2081 – and listed *M. dufourei* [sic] as a synonym. Morelet did not provide a description or illustrations on his own; the brief Latin description he gave below the synonymy list refers to the unnamed variety β . The reference to the illustrations in Chemnitz nonetheless suffices as indication of a new taxon; the correct name is therefore *Melanopsis maroccana* Morelet, 1853. Although Morelet (1853) considered the older *M. dufourii* as a junior synonym of *M. maroccana*, the two names are no objective synonyms because Morelet referred to more specimens than just the holotype of *M. dufourii* (fig. 16). Note, moreover, that Morelet’s synonymization is likely based on a wrong concept of *M. dufourii*, which is a fossil species described from the Miocene of France.

The name “*marocana*” as mentioned in Pallary (1920c: 145) is an incorrect subsequent spelling.

***Melanopsis marteli* Pallary, 1920**

Original source. Pallary 1920a: 32.

Type locality. “Près de Taforalt; Oued Cherâa à Berkane” [near Taforhalt; Oued Cherraa at Berkane], Morocco.

† *Melanopsis marticensis* Matheron, 1842

Original source. Matheron 1842: 292, pl. 37, fig. 7.

Type horizon. Early Campanian, Cretaceous.

Type locality. “Les Martigues”, France.

† *Melanopsis martiniana* Férussac, 1823

Original source. Férussac 1819–1851: *Mélanopsides fossiles*, pl. 2 (1823), figs 11–13 or Férussac 1823: 155, pl. 8, figs 11–13 (precedence not established; see introduction for discussion).

Type horizon. Pannonian, late Miocene.

Type locality. “Dans les environs de Bisentz et de Scharditz, en Moravie, dans la vallée de la Marsch, affluent du Danube” [in the surroundings of Bzenec and Šardice in Moravia, in the valley of the March river, a tributary of the Danube], Czech Republic.

Remarks. Junior objective synonym of *M. fossilis* (for details see there). The name “*martinii*” as mentioned by numerous authors (e.g., Chenu 1859: 297) is an incorrect subsequent spelling.

† *Melanopsis matheroni* Mayer, 1871

Original source. Mayer 1871: 201.

Type horizon. Messinian, late Miocene.

Type locality. “Narzole?” (p. 189; apparently the locality was uncertain), Italy.

Remarks. Wenz (1929: 2783) considered this taxon as a junior synonym of *Melanopsis narzolina* d’Archiac in Viquesnel, 1846.

Melanopsis letourneuxi var. *mattarica* Pallary, 1911

Original source. Pallary 1911: 132, [unnumbered plate], figs 13–14.

Type locality. “Berguent, le Ras al Aïqun des Beni-Mattar” [Ras el Aïn at Aïn Beni Mathar], Morocco.

Remarks. Introduced for *M. letourneuxi* sensu Pallary, 1899 (pl. 8, fig. 6), non Bourguignat, 1884.

Melanopsis mauretanica Pallary, 1922 [invalid]

Original source. Pallary 1922: 207.

Remarks. Unjustified emendation and therefore junior objective synonym of *M. mauritanica* Bourguignat, 1884.

Melanopsis mauritanica Bourguignat, 1884

Original source. Bourguignat 1884: 102.

Type locality. “Çà et là de Maroc” [here and there in Morocco; no locality indicated].

† *Melanopsis mausseneti* Cossmann, 1888

Original source. Cossmann 1888: 284, pl. 11, fig. 10.

Type horizon. Sparnacian, early Ypresian, Eocene.

Type locality. “Mont Bernon” [near Épernay], France.

Melanopsis saulcyi var. *maxima* Dautzenberg, 1894

Original source. Dautzenberg 1894: 345.

Type locality. “Bir Jaloûd” [not found, in the Middle East].

***Melanopsis praemorsa maximalis* Schütt & Bilgin, 1974**

Original source. Schütt and Bilgin 1974: 63, fig. 6.

Type locality. “Sakarya başı, main spring of Sakarya river near village Çifteler, 60 km SE Eskişehir, 160 km WSW of Ankara”, Turkey.

Types. Senckenberg Forschungsinstitut und Naturmuseum Frankfurt, coll. no. SMF 232011.

***Melanopsis maroccana* var. *media* Blanckenhorn, 1897**

Original source. Blanckenhorn 1897: 124, pl. 10, fig. 4.

Type locality. “Dans les eaux le l’ancien Léonthes” (Mousson 1854: 51) [in the Litani river], Lebanon.

Remarks. Replacement name for *M. brevis* Parreyss in Mousson, 1854, non Sowerby, 1826 (see also Pallary 1916). Blanckenhorn (1897) denoted the authority as “Bourg.”, but from the discussion it is clear that Blanckenhorn must be the author.

***Melanopsis marocana* [sic] var. *media* Pallary, 1920 [invalid]**

Original source. Pallary 1920c: 146, pl. 4, fig. 14.

Type locality. “Taza. Fes (dar el Maghzen)”, Morocco.

Remarks. Junior homonym of *Melanopsis maroccana media* Blanckenhorn, 1897 (see Note 1).

***Melanopsis sevillensis* var. *media* Pallary, 1924 [invalid]**

Original source. Pallary 1924: 252.

Type locality. “De la petite rivière de Guadaira qui se jette dans le Guadalquivir” [in the little river Guadaira which flows into the Guadalquivir], Spain.

Remarks. Junior homonym of *Melanopsis maroccana media* Blanckenhorn, 1897 (see Note 1).

† *Melanopsis medinae* Neubauer, Mandić, Harzhauser & Hrvatović, 2013

Original source. Neubauer et al. 2013: 135, figs 5A–D.

Type horizon. Langhian, middle Miocene.

Type locality. “Žepj” (Neumayr 1880c: 480) [Džepi], Bosnia and Herzegovina.

Remarks. Replacement name for the junior secondary homonym *Melanoptychia bittneri* Neumayr, 1880, non *Melanopsis bittneri* Fuchs, 1877. Neubauer et al. (2016c: 275–276) considered the species as a junior synonym of *Melanopsis carusi* (Brusina, 1902).

† *Melanopsis (Canthidomus) bouei* var. *megacantha* Handmann, 1887

Original source. Handmann 1887: 36, pl. 8, figs 13–15.

Type horizon. Pannonian, zone B–D, late Miocene.

Type locality. “Leobersdorf”, Austria.

Remarks. Originally the gender was indicated as masculine (“*megacanthus*”), but *Melanopsis* is feminine, which is why the name must be corrected to “*megacantha*”.

† *Melanopsis themaki* var. *megalostoma* Brusina, 1903

Original source. Brusina 1903: 111.

Type horizon. Late Pleistocene–early Holocene.

Type locality. “Bischofsbad” [Püspökfürdő, Băile 1 Mai, Lake Peța], Romania.

Remarks. Neubauer et al. (2014d: 125) considered this taxon as a junior synonym of *Microcolpia parreyssii sikorai* (Brusina, 1903).

† “*Melanopsis sikorai* var. *megaotyla*” mentioned in Brusina (1903: 112) [unavailable]

Horizon. Late Pleistocene–early Holocene.

Locality. “Bischofsbad” [Püspökfürdő, Băile 1 Mai, Lake Peța], Romania.

Remarks. Nomen nudum. If available, it would be a junior objective synonym of *M. sikorai*: Brusina (1903) indicated it as the typical form of the species. Neubauer et al. (2014d: 125) considered this taxon as a junior synonym of *Microcolpia parreyssii sikorai* (Brusina, 1903).

† *Melanopsis hazayi* var. *megatyla* Brusina, 1903

Original source. Brusina 1903: 112.

Type horizon. Late Pleistocene–early Holocene.

Type locality. “Bischofsbad” [Püspökfürdő, Băile 1 Mai, Lake Peța], Romania.

Remarks. Neubauer et al. (2014d: 125) considered this taxon as a junior synonym of *Microcolpia parreyssii sikorai* (Brusina, 1903).

Melanopsis meiotoma Heller & Sivan, 2000

Original source. Heller and Sivan 2000: 1, fig. 1.

Type locality. “En Haruv [...], a small spring on the Golan Heights that pours into a small cement pool” [near Kefar Haruv], Syria.

Types. National mollusc collection of the Hebrew University, Jerusalem, coll. no. HUJ 7966.

***Melanopsis crenocarinata* var. *melanostoma* Moricand, 1841**

Original source. Moricand 1841: 62.

Type locality. “Rio de Pedra Branca, procince de Bahia” [Pedra Branca river, province Bahia], Brazil.

Remarks. Although not explicitly stated, this variety was apparently considered to belong to the new genus *Verena* by Adams and Adams (1854) (Thiaridae), of which *Melanopsis crenocarinata* is the type species (see Nuttall 1990: 253).

***Melanopsis mellalensis* Pallary, 1928**

Original source. Pallary 1928b: 18, pl. 2, figs 18–23.

Type locality. “L’Oued Taguenout à Beni Mellal” [Beni Mellal, in the Oued Taguenout (?)], Morocco.

***Melanopsis (Mesopotamia) mesopotamica* Pallary, 1939**

Original source. Pallary 1939: 99, pl. 5, figs 1–6.

Type locality. “‘Ain Arouss (la source de la fiancée), près de Tell Abiad, d’où naît le Nahr Bâhlik, qui se jette dans Euphrate, un peu au-dessous de Rakka, la métropole de Haroun el Rachid” [‘Ayn al ‘Arūs (source of the bride), near Tall Abyaḍ, from which arises the Nahr al Balikh, which flows into the Euphrates, a little below Ar Raqqa, the metropolis of Harun al-Rashid], Syria.

Remarks. Heller et al. (2005: 254) considered the species as a junior synonym of *Melanopsis infracincta* Martens, 1874.

† *Melanopsis metochiana* Pavlović, 1932

Original source. Pavlović 1932: 239, 247, pl. 1, figs 3–5.

Type horizon. Pontian (Dacian Basin), late Miocene–Pliocene.

Type locality. “Села Дрсчука” [village Drsnik], Kosovo.

Types. The illustrated syntype is stored in the Natural History Museum, Belgrade, coll. no. 1196 (Milošević 1962: 24).

***Melanopsis cerithiopsis* var. *mezeribensis* Pallary, 1939**

Original source. Pallary 1939: 95, pl. 6, figs 37–39.

Type locality. “Mézérîb” [Muzayrib], Syria.

† *Pseudofagotia michailowskii* Anistratenko, 1993

Original source. Anistratenko 1993: 73, textfig. 2.

Type horizon. Duab Beds, middle to late Kimmerian, Pliocene.

Type locality. “Окр. с. Мокѳу, Очамчирский р-н” [near the village Mok’vi, Ochamchirskiy rayon], Georgia.

Types. Schmalhausen Institute of Zoology of National Academy of Sciences of Ukraine, Kiev; no number indicated.

† *Melanopsis michelottiana* Pallary, 1916

Original source. Pallary 1916: 81.

Type horizon. Late Miocene.

Type locality. “À St. Agata près de Tortone” (Michelotti 1847: 191; the other localities Michelotti apparently adopted from other works) [at Sant’Agata Fossili near Tortona], Italy.

Remarks. Introduced for *M. carinata* sensu Michelotti, 1847, non Sowerby, 1826. Wenz (1929: 2671) considered the taxon as a junior synonym of *Melanopsis bonellii* Manzoni, 1870.

Melanopsis microcolpia Bourguignat, 1884

Original source. Bourguignat 1884: 81.

Type locality. “Près de Jéricho, dans la fontaine de Jérémie (Palestine)” [near Jericho, in the spring of Jeremiah (?)], Palestine.

Melanopsis microstoma Bourguignat, 1884

Original source. Bourguignat 1884: 159.

Type locality. “Ruisseau de la source de la Moulouiah, près de Lalla-Maghnia sur la frontière marocaine (prov. d’Oran)” [in the source of the river Moulouya, near Maghnia at the border to Morocco], Algeria.

† “*Melanopsis microstoma*” mentioned in Newton (1891: 203) [unavailable]

Horizon. Woolwich Beds, early Eocene.

Locality. Woolwich, United Kingdom.

Remarks. Nomen nudum. If available, it would be a junior homonym of *M. microstoma* Bourguignat, 1884.

***Melanopsis mingrelica* Mousson, 1863**

Original source. Mousson 1863: 411.

Type locality. “De l’intérieur de la Mingrélie [...], puis de Réduktaleh” [from the interior of Samegrelo; from Q’ulevi (also read as Kulevi, former Redut-Kale)], Georgia.

Remarks. Mousson attributed the authority to Bayer, but there is no evidence that the description really derived from that author. Bayer only seems to have collected a part of the material.

† ***Melanopsis minima* Blanckenhorn, 1897**

Original source. Blanckenhorn 1897: 114, pl. 9, figs 2–5.

Type horizon. Plio-Pleistocene.

Type locality. “In der tiefsten Thonbank des linken Orontesufers bei Dschir esch-Schurr” [in the lowest clay bank at the left riverside of the Orontes near Jisr Ash-Shughur], Syria.

Remarks. Introduced as “n. mut.” but clearly as a binomen and hence not infrasubspecific in the sense of ICZN Art. 45.6.

† ***Melanopsis buccinoidea* var. *minor* Grateloup, 1838**

Original source. Grateloup 1838: 146.

Type horizon. Burdigalian, early Miocene.

Type locality. “Dax. [...] Mandillot; à Saint-Paul”, France.

***Melanopsis acicularis* var. *minor* Rossmässler, 1839 [invalid]**

Original source. Rossmässler 1838–1844: 41, pl. 50, fig. 672.

Type locality. “In schwach schwefeligen Quellen [...] bei Vöslau unweit Baden” [in weakly sulfurous springs stones and sands at Vöslau near Baden], Austria.

Remarks. Junior objective synonym of *M. daudebartii* [Prevost], 1821, which Rossmässler listed in synonymy. Moreover, the name is a junior homonym of *Melanopsis buccinoidea minor* Grateloup, 1838 (see Note 1).

***Melanopsis costata* var. *minor* Martens, 1874 [invalid]**

Original source. Martens 1874: 33, p. 5, fig. 40.

Type locality. “Quellen des Chabur bei Ras-el-Ain” [source of Chabur river near Ra’s al ‘Ayn], Syria.

Remarks. Junior homonym of *Melanopsis buccinoidea minor* Grateloup, 1838 (see Note 1).

† *Melanopsis bonellii* var. *minor* Locard, 1878 [invalid]

Original source. Locard 1878: 56.

Type horizon. Tortonian, late Miocene.

Type locality. “À Tersannes près de Hauterives (Drôme)” [at Tersanne near Hauterives], France.

Remarks. Junior homonym of *Melanopsis buccinoidea minor* Grateloup, 1838 (see Note 1). Wenz (1929: 2669) considered the taxon as a junior synonym of *M. bonellii* Manzoni, 1870.

Melanopsis belonidaea var. *minor* Bourguignat, 1884 [invalid]

Original source. Bourguignat 1884: 111.

Type locality. “Ruisseau d’eau chaude à Ouargla (prov. de Constantine) et eaux thermales du Djérid, au nord du chott Tiraoun (sud de la Tunisie)” [in warm waters at Ouargla (Algeria) and thermal water of Djérid, north of chott Tiraoun (Tunisia)].

Remarks. Junior homonym of *Melanopsis buccinoidea minor* Grateloup, 1838 (see Note 1).

Melanopsis prophetarum var. *minor* Bourguignat, 1884 [invalid]

Original source. Bourguignat 1884: 82.

Type locality. “Lac d’Antioche” [Lake Anuk (also as Amik)], Turkey.

Remarks. Junior homonym of *Melanopsis buccinoidea minor* Grateloup, 1838 (see Note 1). Blanckenhorn (1897) indicated himself as author of the taxon, but he clearly cited the record by Bourguignat (1884) in the synonymy list. Wenz (1929) introduced *M. blanckenhorni* as replacement name.

Melanopsis salomonis var. *minor* Bourguignat, 1884 [invalid]

Original source. Bourguignat 1884: 96.

Type locality. “Environs d’Alep, à Sadjour-Sou, à quatre kilom. en aval d’Aïn-Taïb; ruisseaux à Doumar, sur l’oued Baradah, près Aïn-Fidji, et à Banias, en Syrie” [Surroundings of Aleppo (Syria), at Sadjour-Sou, 4 km downstream of Gaziantep (Turkey); streams at Dummar, at the river Barada, near Aïn al-Fiji and at Bāniyās (Syria)].

Remarks. Junior homonym of *Melanopsis buccinoidea minor* Grateloup, 1838 (see Note 1).

***Microcolpia letourneuxi* var. *minor* Bourguignat, 1884**

Original source. Bourguignat 1884: 65.

Type locality. Not indicated, but probably as for the species (“Dans la Save, au-dessous d’Agram” [in the Sava river below Zagreb], Croatia).

Remarks. Starobogatov et al. (1992: 65) considered the taxon as a junior synonym of *Microcolpia cornea* (Pfeiffer, 1828).

Melania (*Amphimelania*) *holandri* f. *minor* Westerlund, 1886 [invalid]

Original source. Westerlund 1886: 104.

Type locality. Not indicated.

Remarks. Junior homonym of *Melania tuberculata minor* Brot, 1877 from Sri Lanka (see Note 1).

***Melanopsis gracilentia* var. *minor* Pallary, 1911 [invalid]**

Original source. Pallary 1911: 130, [unnumbered plate], fig. 20.

Type locality. “Tout près d’Oudjda, à 4 kilom. S.-E., sourdent les belles sources de Sidi-Yahia qui alimentent une véritable oasis, puis la ville d’Oudjda, et vont finalement se déverser dans l’oued Isly” [near Oujda, 4 km southeast, at the sources of Sidi Yahya that feed an oasis and the city of Oujda, and ultimately will flow into the Oued Isly], Morocco.

Remarks. Junior homonym of *Melanopsis buccinoidea minor* Grateloup, 1838 (see Note 1).

***Melanopsis letourneuxi* var. *minor* Pallary, 1911 [invalid]**

Original source. Pallary 1911: 132, [unnumbered plate], fig. 16.

Type locality. “Berguent, le Ras al Aïqun des Beni-Mattar” [Ras el Aïn at Aïn Beni Mathar], Morocco.

Remarks. Junior homonym of *Melanopsis buccinoidea minor* Grateloup, 1838 (see Note 1).

***Melanopsis nobilis* var. *minor* Pallary, 1912 [invalid]**

Original source. Pallary 1912: 21, fig. 42.

Type locality. “Des bords du Chott Djerid, à Tozeur” [banks of the Chott el Djérid at Tozeur], Tunisia.

Remarks. Junior homonym of *Melanopsis buccinoidea minor* Grateloup, 1838 (see Note 1).

***Melanopsis attenuata* var. *minor* Pallary, 1920 [invalid]**

Original source. Pallary 1920a: 29.

Type locality. “Tétouan”, Morocco.

Remarks. Junior homonym of *Melanopsis buccinoidea minor* Grateloup, 1838 (see Note 1).

***Melanopsis douttei* var. *decorata minor* Pallary, 1920 [unavailable]**

Original source. Pallary 1920c: 150.

Type locality. Not indicated, but probably as for the variety (“La Makina”, Morocco).

Remarks. Introduced as infrasubspecific taxon, which is not ruled by the provisions of the Code.

***Melanopsis fasesnsis* var. *minor* Pallary, 1920 [invalid]**

Original source. Pallary 1920c: 148.

Type locality. “Dar Batha” [near Fes], Morocco.

Remarks. Junior homonym of *Melanopsis buccinoidea minor* Grateloup, 1838 (see Note 1).

***Melanopsis pseudoferussaci* var. *minor* Pallary, 1920 [invalid]**

Original source. Pallary 1920c: 147, pl. 4, figs 15–16.

Type locality. “Fès: source près de la ferme proche du pont neuf, à dar Mahrès; Sidi Harazen (source chaude); Moulay Idriss du Zehroun, source sulfureuse chaude; El Menzel; Tazouta” [Fes: source near the farm at the new bridge, at Dar Mahres; Sidi Harazem (hot spring); Moulay Driss Zerhoun, hot sulfur springs; El Menzel; Tazouta], Morocco.

Remarks. Junior homonym of *Melanopsis buccinoidea minor* Grateloup, 1838 (see Note 1).

***Melanopsis dufourii* var. *minor* Pallary, 1924 [invalid]**

Original source. Pallary 1924: 249.

Type locality. “Fortuna et Caravala [Caravaca?] (Murcia). Lorca, Valencia et Alicante”, Spain.

Remarks. Junior homonym of *Melanopsis buccinoidea minor* Grateloup, 1838 (see Note 1).

***Melanopsis lorcana* f. *minor* Pallary, 1924 [invalid]**

Original source. Pallary 1924: 247, pl. 15, fig. 8.

Type locality. Spain [no locality indicated].

Remarks. Junior homonym of *Melanopsis buccinoidea minor* Grateloup, 1838 (see Note 1).

***Melanopsis hadiensis* var. *minor* Pallary, 1928 [invalid]**

Original source. Pallary 1928a: 269.

Type locality. “Bel Hadi-Kenadsa (Sud Oranais)” [Bel Hadi, near Kenadsa], Algeria.

Remarks. Junior homonym of *Melanopsis buccinoidea minor* Grateloup, 1838 (see Note 1).

***Melanopsis macilenta* var. *minor* Pallary, 1928 [invalid]**

Original source. Pallary 1928b: 18, pl. 2, fig. 17.

Type locality. “O. Taguenout” [Oued Taguenout (?), said to be near Beni Mellal], Morocco.

Remarks. Junior homonym of *Melanopsis buccinoidea minor* Grateloup, 1838 (see Note 1).

***Melanopsis mellalensis* var. *minor* Pallary, 1928 [invalid]**

Original source. Pallary 1928b: 19, pl. 2, fig. 21.

Type locality. “Dans l’oued Daï” [not found], Morocco.

Remarks. Junior homonym of *Melanopsis buccinoidea minor* Grateloup, 1838 (see Note 1).

***Melanopsis seurati* var. *minor* Pallary, 1928 [invalid]**

Original source. Pallary 1928a: 267.

Type locality. Not explicitly stated but probably the same as for the species (“La Zousfana, à la hauteur de Figuig et à Beni Ounif” [Oued Zousfana, at the height of Figuig, and at Beni Ounif], Algeria).

Remarks. Junior homonym of *Melanopsis buccinoidea minor* Grateloup, 1838 (see Note 1).

***Melanopsis turgida* var. *minor* Pallary, 1928 [invalid]**

Original source. Pallary 1928a: 259, pl. 5, fig. 12.

Type locality. “Moulāï Taïeb” [Moulay Taïeb], Morocco.

Remarks. Name appears only in the plate captions. Junior homonym of *Melanopsis buccinoidea minor* Grateloup, 1838 (see Note 1).

***Melanopsis tutulata* var. *minor* Pallary, 1928 [invalid]**

Original source. Pallary 1928a: 272.

Type locality. Not stated but probably the same or partly as for the species (“Berguent; Aoûllout; Ras el Mâ de Fès; O. Chkef près Fès” [Aïn Beni Mathar, Aïn Aoûllout, Ras El Ma, Oued Aïn Chkef in Fes], Morocco).

Remarks. Junior homonym of *Melanopsis buccinoidea minor* Grateloup, 1838 (see Note 1).

***Melanopsis vondeli* var. *minor* Pallary, 1928 [invalid]**

Original source. Pallary 1928b: 16.

Type locality. “Dans l’oued Daï” [not found], Morocco.

Remarks. Junior homonym of *Melanopsis buccinoidea minor* Grateloup, 1838 (see Note 1).

***Melanopsis wagneri* var. *minor* Pallary, 1930 [invalid]**

Original source. Pallary 1930: 290.

Type locality. “Dans l’eau tiède du bassin de Diane, à Smyrne” [in warm water basin of Halkapınar at Izmir], Turkey.

Remarks. Junior homonym of *Melanopsis buccinoidea minor* Grateloup, 1838 (see Note 1).

***Melanopsis (Mesopotamia) khabourensis* var. *minor* Pallary, 1939 [invalid]**

Original source. Pallary 1939: 103, pl. 5, fig. 21.

Type locality. “Ras el ‘Ain du Khabour” [Chabur river near Ra’s al ‘Ayn], Syria.

Remarks. Junior homonym of *Melanopsis buccinoidea minor* Grateloup, 1838 (see Note 1).

***Melanopsis (Mesopotamia) mesopotamica* var. *minor* Pallary, 1939 [invalid]**

Original source. Pallary 1939: 100, pl. 4, fig. 6.

Type locality. “Dans le canal de la Butte de tir, à 7 km Sud de Baghdad” [in a trench of a firing hill (?), 7 km south of Baghdad], Iraq.

Remarks. Ranked as a variety of *M. nodosa* Férussac in the plate captions. Junior homonym of *Melanopsis buccinoidea minor* Grateloup, 1838 (see Note 1).

***Melanopsis pachya* var. *minor* Pallary, 1939 [invalid]**

Original source. Pallary 1939: 87.

Type locality. “Dans les sources de Mézériß, au N. O. de Derâa (Syrie méridionale)” [in the sources of the Muzayrîb, northwest of Dar’â], Syria.

Remarks. Junior homonym of *Melanopsis buccinoidea minor* Grateloup, 1838 (see Note 1).

† *Melanopsis (Canthidomus) oroposi minoriformis* Papp, 1979

Original source. Papp 1979: 672, pl. 3, figs 11–15.

Type horizon. Mammal zone MN 10, late Miocene.

Type locality. “Milessi” [Milesi], Greece.

Types. Institute of Paleontology, University of Vienna; no number indicated.

***Melanopsis monteli* var. *minorstricta* Pallary, 1936**

Original source. Pallary 1936: 61.

Type locality. Not explicitly stated but probably the same as for the species (“L’Oued Sous, au pont des Aït Melloul, sur la route d’Agadir à Tiznit, à 14 kil. S. O. d’Agadir” [in the Oued Sous, at the bridge of Ait Melloul, at the road from Agadir to Tiznit, 14 km southwest of Agadir], Morocco).

Remarks. Originally spelt as “*minor-stricta*”.

† *Melanopsis minotauri* Willmann, 1980

Original source. Willmann 1980: 285.

Type horizon. Viannos Formation, early Tortonian, late Miocene.

Type locality. “Tongrube 1 km westlich von Limin Chersonisou/Kreta” [claypit 1 km west of Chersonisos, near Agia Anna chapel], Greece.

† *Melanopsis minuta* Pallary, 1916

Original source. Pallary 1916: 77.

Type horizon. Mammal zone MN 10–12, late Miocene.

Type locality. “De Cuiseaux”, France.

Remarks. Used as valid name by Wenz (1929: 2780), but obviously not intended as species-group name by Férussac (1823: 150, pl. 7, fig. 4). The name became nevertheless available from Pallary (1916) who treated it as valid species-group name and associated it with an illustration in Férussac (1823: pl. 7, fig. 4). See introduction for a detailed discussion about the names used by Férussac (1823).

Melanopsis minutula Bourguignat, 1884

Original source. Bourguignat 1884: 92.

Type locality. “Fontaine froide du Hammam à Brousse (Anatolie); Nahr-Antalies dans le Liban (Syrie); puits artésien de Tamerna-Kedima, dans le Ziban (Algérie)” [cold springs in a Hamam at Bursa (Turkey); Nahr-Antalies (?) in the Lebanon; artesian well of Tamerna Kedima (Algeria)].

† *Melanopsis lyrata* var. *misera* Brusina, 1874

Original source. Brusina 1874: 45.

Type horizon. Langhian, middle Miocene.

Type locality. “Vrba; Sinj (Stuparuša); Turiake [Turjaci]”, Croatia.

Types. Milan et al. (1974: 94) indicated a holotype, but it is uncertain whether the specimen is part of the original type series and whether it was the only one Brusina had at hand (holotype by monotypy, Art. 73.1.2). The specimen is stored in the Croatian Natural History Museum, Zagreb, coll. no. 2972-618.

Remarks. Neubauer et al. (2011: 207) considered this taxon as a junior synonym of *Melanopsis lyrata* Neumayr, 1869.

† *Melanopsis (Lyrcaea)* [sic] *mitraeformis* Andrusov, 1909

Original source. Andrusov 1909: 87, 160, pl. 4, figs 34–35.

Type horizon. Pontian (sensu stricto), late Miocene.

Type locality. “Babadjan” [Babadzhan], Azerbaijan.

† *Melanopsis (Melanosteira) mitzopoulosi* Papp, 1955

Original source. Papp 1955: 126, pl. 20, figs 15–19.

Type horizon. Late Pliocene to early Pleistocene.

Type locality. “Nord-Nordwestlich von der Solfatare bei Susaki” [north-north-west of Solfatara Sousáki], Greece.

Types. Museum of Palaeontology and Geology of the University of Athens; no number indicated.

Remarks. First mentioned as nomen nudum in Papp and Thenius (1952: 4). Junior synonym of *Amphimelania gayi* [sic] after Bandel (2000: 140).

† *Melanopsis (Canthidomus) mitzopoulosi* Kühn, 1963 [invalid]

Original source. Kühn 1963: 386.

Type horizon. Mammal zone MN 10, late Miocene.

Type locality. “Peristeri”, Greece.

Types. Museum of Palaeontology and Geology of the University of Athens, coll. no. 1963/83.

Remarks. The status of this species is unclear. First of all, Kühn did not at all refer to the senior homonym *Melanopsis (Melanosteira) mitzopoulosi* Papp, 1955 but clearly described the species as new. Neither did Papp (1979: 668), who must have been aware of the issue and who ranked it as a subspecies of *M. longa* Deshayes in Férussac, 1839. Moreover, Papp stated that Kühn’s species was based on the material collected by Papp (1947) and considered the type locality mentioned by Kühn a mistake. Kühn (1963), however, did not refer to Papp’s material at all.

Melanopsis nodosa var. *moderata* Mousson, 1874

Original source. Mousson 1874: 48.

Type locality. Not explicitly stated but probably the same as for the species (“Au cours de l’Euphrate et du Tigre” [from the Euphrates and Tigris rivers], Iraq).

Remarks. Brot (1874–1879: 432) considered the taxon as a junior synonym of *Melanopsis nodosa* Férussac, 1822, which is actually a fossil species described from the Miocene of France.

† “*Melanopsis moderata*” mentioned in Bogachev (1938: 13, 33, 53) [unavailable]

Horizon. Miocene.

Locality. “Tori bei Borshomi” [Tori near Borjomi], Georgia.

Remarks. The name was only mentioned in a species list by Bogachev without description or illustration. If available, it would be a junior homonym of *Melanopsis nodosa moderata* Mousson, 1874 (see Note 1).

***Melanopsis neolithica* var. *moderna* Pallary, 1911**

Original source. Pallary 1911: 133.

Type locality. “À Colomb-Béchar” [Bechar], Algeria.

† “*Melanopsis modesta*” mentioned in Bogachev (1938: 13, 33, 53) [unavailable]

Horizon. Miocene.

Locality. “Tori bei Borshomi” [Tori near Borjomi], Georgia.

Remarks. The name was only mentioned in a species list by Bogachev without description or illustration.

***Melanopsis modica* Pallary, 1928**

Original source. Pallary 1928b: 19, pl. 2, figs 24–27.

Type locality. “Aïn Foum el Ançeur et Tirboula, près de Ksiba (Moyen Atlas méridional)” [Foum El Anceur and Tirboula, near Ksiba (southern Middle Atlas)], Morocco.

† ***Melanopsis moesiensis* Jekelius, 1944**

Original source. Jekelius 1944: 132, pl. 50, figs 1–2.

Type horizon. Early Pannonian, late Miocene.

Type locality. “Turislav-Tal bei Soceni” [Turislav valley near Soceni], Romania.

† ***Melanoptychia moesiensis* Jekelius, 1944 [invalid]**

Original source. Jekelius 1944: 139, pl. 58, figs 1–10.

Type horizon. Early Pannonian, late Miocene.

Type locality. “Turislav-Tal bei Soceni” [Turislav valley near Soceni], Romania.

Remarks. Junior secondary homonym of *Melanopsis moesiensis* Jekelius, 1944 (same work). Neubauer et al. (2014a: 463) considered both synonymous and gave priority to *Melanopsis moesiensis* (since *Melanoptychia* is considered a synonym of *Melanopsis*).

***Melanopsis mogadorensis* Pallary, 1911**

Original source. Pallary 1911: 134, [unnumbered plate], figs 27–30.

Type locality. “Dans les sources et la rivière du jardin du Sultan, à Mogador” [in the sources and the river of the garden of the Sultan, at Essaouira], Morocco.

***Melanopsis mohammedi* Bourguignat, 1884**

Original source. Bourguignat 1884: 110.

Type locality. “D’Agadir” (Morelet 1880: 71), Morocco.

Remarks. Introduced for *M. praerosa* [= *M. praemorsa*] sensu Morelet, 1880, non Linnaeus, 1758. Note that Bourguignat denoted the authority as “Bourguignat, 1881”.

† ***Melanoptychia mojsisovicsi* Neumayr, 1880**

Original source. Neumayr 1880c: 481, pl. 7, figs 9–10.

Type horizon. Langhian, middle Miocene.

Type locality. “Žepj” [Džepi], Bosnia and Herzegovina.

Types. The type material, with all specimens studied by Neumayr (1880), is lost. Neubauer et al. (2016c: 278) defined a neotype based on material from the type locality. The specimen is stored in the Geological-Paleontological Department, Natural History Museum Vienna, coll. no. 2014/0364/0013.

† ***Melanopsis molinai* Vidal, 1917**

Original source. Vidal 1917: 353, pl. 2, figs 9–10.

Type horizon. Cretaceous.

Type locality. “Lignitos de Selva y Binisalém” [lignites of Selva and Binissalem], Spain.

† ***Melanopsis bouei* var. *monacantha* Handmann, 1882**

Original source. Handmann 1887: 35, pl. 8, figs 8–9.

Type horizon. Pannonian, zone D, late Miocene.

Type locality. “Kottingbrunn [...] Ziegelei a”, Austria.

Remarks. Wenz (1929: 2675) considered this taxon as a junior synonym of *Melanopsis bouei* Férussac, 1823.

† “*Melanopsis monolithos*” mentioned in Dollfus (1922: 118) [unavailable]

Horizon. Apolakkia/Monolithos Formation, Pliocene.

Locality (Type locality of *M. biliottii*). “Rhodos” (locality specified as “Monolithos” in Bukowski 1893), Greece.

Remarks. Dollfus mentioned the name as nomen nudum without explanation alongside the discussion of [*M. costata*] “Var. *Biliottii* Buk.”. Apparently, he considered it a potential new name should *biliottii* be elevated to the species level. If not already a nomen nudum, *M. monolithos* would therefore be a junior objective synonym of *M. biliottii* Bukowski, 1892, which actually was introduced as a distinct species (see also Willmann 1981: 105).

† *Melanopsis impressa* var. *monregalensis* Sacco, 1889

Original source. Sacco 1889: 66, pl. 2, figs 10–12.

Type horizon. Messinian, late Miocene.

Type locality. “Santuario di Mondovì” [Santuario di Vicoforte], Italy.

Melanopsis monteli Pallary, 1936

Original source. Pallary 1936: 60, pl. 2, fig. 10.

Type locality. “L’Oued Sous, au pont des Aït Melloul, sur la route d’Agadir à Tiznit, à 14 kil. S. O. d’Agadir” [in the Oued Sous, at the bridge of Ait Melloul, at the road from Agadir to Tiznit, 14 km southwest of Agadir], Morocco.

Melania hollandri [sic] var. *montenegrina* Walderdorff, 1864

Original source. Walderdorff 1864: 512.

Type locality. “In den Bächen Krains” [in rivers of Carniola, a historical region that comprised parts of present-day Slovenia].

Melanopsis moreleti Pallary, 1916

Original source. Pallary 1916: 81.

Type locality. “[Sancta-Maria de Balade]” (Morelet 1857: 32) [Balade], New Caledonia.

Remarks. Replacement name for *Melanopsis brevis* Morelet, 1857, non Sowerby, 1826.

† *Melanopsis morrisi* Wenz, 1928

Original source. Wenz 1928b: 220.

Type horizon. Headon Beds, Priabonian, Eocene.

Type locality. “Headon Hill” (Forbes 1856: 156), United Kingdom.

Remarks. Introduced for *Melanopsis subcarinata* Morris in Forbes, 1856, non Deshayes, 1851.

Melanopsis mourebeyensis Pallary, 1901

Original source. Pallary 1901b: 228.

Type locality. “Oued Mourebēi sur la route de Merrakech” [Oued Mourebēi (?) at the road to Marrakech], Morocco.

Melanopsis moussoni Pallary, 1916 [invalid]

Original source. Pallary 1916: 81.

Type locality. “Dans les eaux de l’ancien Léonthes” (Mousson 1854: 51) [in the Litani river], Lebanon.

Remarks. Replacement name for *Melanopsis brevis* Parreyss in Mousson, 1854, non Sowerby, 1826, for which already Blanckenhorn (1897) had introduced *M. media* as replacement name. Thus, *M. moussoni* is a junior objective synonym of *M. media*.

† *Melanopsis pygmaea* var. *mucronata* Handmann, 1887

Original source. Handmann 1887: 13, pl. 1, fig. 1.

Type horizon. Pannonian, zone B–D, late Miocene.

Type locality. “Leobersdorf”, Austria.

† *Melanopsis mucronifera* Kormos, 1905

Original source. Kormos 1905: 395, 442, pl. 2, fig. 1.

Type horizon. Late Pleistocene–early Holocene.

Type locality. “Püspökfürdő” [Băile 1 Mai, Lake Pețea], Romania.

Remarks. Obviously unaware of the fact that variety names are available in nomenclature as species-group names, Kormos (1903) stated in the description of his new variety *M. sikorai carinata* that, if raised to species, he suggests “*M. mucronifera*” as name for the taxon. Both names were simultaneously published and are objective

synonyms. Since *M. carinata* Kormos is a junior homonym of *M. carinata* Sowerby, 1826, *M. mucronifera* is the valid name of the taxon. Neubauer et al. (2014d) considered the taxon as a junior synonym of *Microcolpia parreyssii sikorai* (Brusina, 1903).

† *Melanopsis bouei* var. *multicostata* Handmann, 1882

Original source. Handmann 1882: 557.

Type horizon. Pannonian, zone D, late Miocene.

Type locality. “Kottingbrunn [...] Ziegelei a”, Austria.

Remarks. The name “*multicostellata*” as mentioned by Handmann (1887: 33) is an incorrect subsequent spelling. Wenz (1929: 2674) considered this variety as a junior synonym of *M. bouei* Férussac, 1823.

† *Melanopsis tothi* var. *multiflora* Brusina, 1903

Original source. Brusina 1903: 114.

Type horizon. Late Pleistocene–early Holocene.

Type locality. “Bischofsbad” [Püspöckfürdő, Băile 1 Mai, Lake Peșea], Romania.

Remarks. Considered as a junior synonym of *Microcolpia parreyssii sikorai* (Brusina, 1903) by Neubauer et al. (2014d: 125).

† *Melanopsis doboi multiflora* Schréter, 1975 [invalid]

Original source. Schréter 1975: 10, pl. 2, figs 11–12, pl. 3, fig. 13.

Type horizon. Riss/Würm end to early Würm Ice Age, Pleistocene.

Type locality. “Eger, az egri vár Zárkándy bástyájának vasúti átmetszése” [Eger, section at the railway at the Zarkandy bastion of the fortress Eger], Hungary.

Types. Magyar Állami Földtani Intézet (Hungarian Geological Museum), Budapest; no number indicated.

Remarks. Junior homonym of *Melanopsis tothi* var. *multiflora* Brusina, 1903. Neubauer et al. (2016a) attributed the species *M. doboi* to the genus *Microcolpia*.

† *Melanopsis multiformis* Blanckenhorn, 1897

Original source. Blanckenhorn 1897: 116, pl. 9, figs 14–17.

Type horizon. Plio-Pleistocene.

Type locality. “In der ersten Thonbank des linken Orontesufers bei Dschir esch-Schurr” [in the first clay bank at the left riverside of the Orontes near Jisr Ash-Shughur], Syria.

Remarks. Introduced as “n. mut.” (see also discussion on p. 117) but clearly as a binomen and hence not infrasubspecific in the sense of ICZN Art. 45.6.

† *Melanopsis munieri* Roule, 1884

Original source. Roule 1884: 317, pl. 5, fig. 5.

Type horizon. Calcaire de Rognac, Maastrichtian, Cretaceous.

Type locality. “À Fuveau, Puyloubier, Ollières”, France.

Remarks. Probably not a Melanopsidae.

“*Melanopsis muraldi*” mentioned in Brot (1874–1879: 427) [unavailable]

Locality. Not indicated.

Remarks. Nomen nudum; apparently based on an unpublished manuscript name from Ziegler and listed in synonymy of *Melanopsis costata* (Olivier, 1804) by Brot (1879).

Melanopsis myosotidaea Bourguignat, 1884

Original source. Bourguignat 1884: 93.

Type locality. “Environs d’Agora et d’Alhama (Aragon), en Espagne; Relizane, dans la plaine du Cheliff, en Algérie” [surroundings of Agora (?) and Alhama de Aragón (Spain); Relizane, in the Plaine du Chelif (Algeria)].

Melanopsis mzabica Bourguignat, 1884

Original source. Bourguignat 1884: 89.

Type locality. “Ruisseau du puits artésien de Ngouça; Chetma, près de Biskra, et environs d’Ouargla, dans le sud de la province de Constantine” [creek from the artesian well of N’Goussa; Chetma near Biskra and in the surroundings of Ouargla], Algeria.

Melanopsis naini Pallary, 1928

Original source. Pallary 1928a: 274, pl. 5, figs 13–14.

Type locality. “Tiout, dans l’Anti Atlas”, Morocco.

Melania (Melanella) holandri var. *nana* Kobelt, 1881 [invalid]

Original source. Kobelt 1881: 149.

Type locality. “Teplica” [thermal springs Teplice (?)], Slovakia.

Remarks. Kobelt (1881) and Clessin (1890) attributed the name to Brot (1874), who did not name the variety but solely described it (*nana* = Latin “tiny”). Kobelt

(1881) referred the name to Brot's illustration, making it thereby available. However, the name is a junior homonym of *Melania nana* Lea & Lea, 1850.

***Melanopsis praemorsa variabilis* subvar. *nana* Nevill, 1884 [unavailable]**

Original source. Nevill 1884: 208.

Type locality. “Persia; [...] Shiraz”, Iran.

Remarks. Introduced infrasubspecific taxon (“subvariety”), which is not ruled by the provisions of the Code.

***Melanopsis costata* var. *nana* Paetel, 1888**

Original source. Paetel 1888: 400.

Type locality. “Sea of Galilee” (Nevill 1884: 212), Israel.

Remarks. Originally introduced as infrasubspecific taxon (“subvariety”) by Nevill (1884), but made available by Paetel (1888) who treated it as variety (Art. 45.5.1). Paetel clearly referred to the description of Nevill.

† *Melanopsis narzolina* d’Archiac in Viquesnel, 1846

Original source. Viquesnel 1846: 266.

Type horizon. Late Miocene.

Type locality. “Piémont” [no locality indicated], Italy.

Remarks. The name was based on Bonelli’s unpublished catalogue, but was first described in Viquesnel (1846) as “*M.[elanopsis] narsolina*” [sic]. However, the name *M. narzolina* - probably the “correct” spelling sensu Bonelli - has been in prevailing usage since then. For reasons of stability the name *narzolina* should be maintained according to Art. 33.3.1 (see Harzhauser et al. 2015: 10).

† *Melanopsis narzolinensis* Pallary, 1920 [invalid]

Original source. Pallary 1920b: 105, 112–113.

Type horizon. Late Miocene.

Remarks. Unjustified emendation and therefore junior objective synonym of *Melanopsis narzolina* d’Archiac in Viquesnel, 1846.

† *Melanopsis nassaeformis* Neumayr, 1880

Original source. Neumayr 1880b: 293, pl. 1, fig. 20.

Type horizon. Pliocene.

Type locality. “Bei Pylle” [near Pýli], Greece.

Remarks. Willmann (1981: 175) considered this taxon as a junior synonym of *Melanopsis gorceixi proteus* Tournouër, 1875.

† *Melanopsis navarroï* Vidal, 1917

Original source. Vidal 1917: 354, pl. 2, figs 11–14.

Type horizon. Cretaceous.

Type locality. “De Binisalém” [from Binissalem], Spain.

Melanopsis neolithica Pallary, 1911

Original source. Pallary 1911: 132, [unnumbered plate], figs 18, 23–24.

Type locality. “À Colomb-Béchar” [at Bechar], Algeria.

† *Melanopsis nereis* d’Orbigny, 1852

Original source. d’Orbigny 1852: 3.

Type horizon. Chattian, Oligocene.

Type locality. “Saint-Geours en Marensin, et ceux d’Abesse à Saint-Paul” (Grateloup 1828: 137) [Saint-Geours-de-Mareme and Abesse near Saint-Paul-lès-Dax], France.

Remarks. Introduced for *M. costata* sensu Grateloup, 1828, non Olivier, 1804.

Melanopsis neritiformis Deshayes, 1832

Original source. Deshayes 1830–1832: 438.

Type locality. “Dans l’Ohio et ses affluens” [in the Ohio river and its tributaries], United States.

Melanopsis neritoides Gassies, 1858

Original source. Gassies 1858: 371.

Type locality. “Des petits cours d’eau à Balade” [in small streams near Balade], New Caledonia.

Remarks. Brot (1879: 456) considered the species as a junior synonym of *M. brevis* Morelet, 1857 (which is a junior homonym of *Melanopsis brevis* Sowerby, 1826; see *M. moreleti* Pallary, 1916).

† *Melanopsis nesici* Brusina, 1893

Original source. Brusina 1893: 202 (Serbian part), 36 (Italian part).

Type horizon. Pannonian, zone D–E, late Miocene.

Type locality. “Ripanj”, Serbia.

Types. Milan et al. (1974: 95) indicated collection numbers for “syntypes” illustrated in Brusina (1902), but it is uncertain whether the specimens were part of the original type series. They are stored in the Croatian Natural History Museum, Zagreb, coll. no. 2533-179/1-4.

Remarks. The name “*nesici*” as mentioned in Stepanović (1938) is an incorrect subsequent spelling.

† *Melanopsis neumayri* Tournouër, 1874

Original source. Tournouër 1874: 303, pl. 9, fig. 5.

Type horizon. Mammal zone MN 11, late Miocene.

Type locality. “Visan”, France.

***Melanopsis nicollei* Pallary, 1923**

Original source. Pallary 1923: 43, [unnumbered plate], figs 17–18.

Type locality. “Gafsa, dans les eaux refroidies” [Gafsa, in cool waters], Tunisia.

***Melanopsis doriae* var. *nigra* Biggs, 1934**

Original source. Biggs 1934: 60.

Type locality. “Qanat at Ginehkan near Kerman, S. Persia”, Iran.

“*Melanopsis nivosa* Fer.” mentioned in Paetel (1888: 402) [unavailable]

Locality. “Mesopot.” [Mesopotamia, no locality indicated], Iraq.

Remarks. Nomen nudum, found only in the species list by Paetel (1888). He referred to Bourguignat (1884), but the name could not be found in that work.

† *Melanopsis nobilis* Seninski, 1905

Original source. Seninski 1905: 61, pl. 2, figs 1–2.

Type horizon. Duab Beds, middle to late Kimmerian, Pliocene.

Type locality. “Р. Дуабъ” [Duab river], Georgia.

***Melanopsis nobilis* Pallary, 1912 [invalid]**

Original source. Pallary 1912: 20, fig. 41.

Type locality. “Des bords du Chott Djerid, à Tozeur” [banks of the Chott el Djerid at Tozeur], Tunisia.

Remarks. Junior homonym of *Melanopsis nobilis* Seninski, 1905.

***Fagotia nocturna* Bourguignat, 1884**

Original source. Bourguignat 1884: 44.

Type locality. “La Save à Agram et la rivière entre Plaski et Ostaria (Croatie); le Danube à Ibraïla” [Sava river at Zagreb and river between Plaški and Oštarije (Croatia); Danube river at Brăila (Romania)].

Remarks. Note that Bourguignat denoted the authority as “Bourguignat, 1879”. Starobogatov et al. (1992: 60) considered the species as a junior synonym of *Fagotia* [= *Esperiana*] *esperi* (Férussac, 1823).

† *Melanopsis nodescens* Handmann, 1882

Original source. Handmann 1882: 557.

Type horizon. Pannonian, zone D, late Miocene.

Type locality. “Kottingbrunn [...] Ziegelei a”, Austria.

Remarks. Handmann (1887) listed a *M. varicosa nodescens*. It is unclear whether he actually introduced a new variety or ranked the species described in 1882 as variety of *M. varicosa* Handmann, 1882. Wenz (1929: 2674) considered *M. nodescens* Handmann, 1882 as a junior synonym of *Melanopsis bouei* Férussac, 1823.

† *Melanopsis nodicincta* Pallary, 1920

Original source. Pallary 1920b: 111.

Type horizon. Late Villafranchian, early Pleistocene.

Type locality. “Roccantica presso Poggio Mirteto; Fra Otricoli e le Vigne” (De Stefani 1880: 62–63) [Roccantica near Poggio Mirteto; between Otricoli and Le Vigne], Italy.

Remarks. Introduced for *M. nodosa* sensu De Stefani, 1880, non Férussac, 1822. Wenz (1929: 2654) considered the taxon as a junior synonym of “*M. affinis* Férussac”, which is not an available name.

† *Melanopsis (Canthidomus) nodifera* Handmann, 1887

Original source. Handmann 1887: 30, pl. 7, figs 4–6.

Type horizon. Pannonian, zone B–D, late Miocene.

Type locality. “Leobersdorf”, Austria.

† *Melanopsis nodosa* Férussac, 1822

Original source. Férussac 1819–1851: Mélanopsides fossiles, pl. 1 (1822), fig. 13.

Type horizon. Late Villafranchian, early Pleistocene.

Type locality. “Entre Otricoli et le Vigne, route de Rome à Foligno” [between Otricoli and Le Vigne, at the road from Rome to Foligno], Italy.

Remarks. The name first appeared in 1822 on the captions for plate 1 of the “Mélanopsides fossiles” in Férussac’s “Histoire naturelle” (see also introduction for details). While Férussac (1823) included also recent specimens under that name in his monograph on the Melanopsidae, the only specimen illustrated in 1822 in the “Histoire naturelle” was a fossil one from the Villafranchian of Italy. This fact remained widely unknown to biologists and paleontologists alike. The consequence, however, is that probably none of the specimens referred to as *M. nodosa* in the biological literature (e.g., Pallary 1939, Glaubrecht 1993, 1996) or the IUCN Red List (Van Damme 2014), which reports it from Anatolia and the Middle East, actually correspond to the real *M. nodosa*.

Fossil Italian specimens are often erroneously referred to as “*Melanopsis affinis* Férussac” or “*M. antiqua* Férussac” (Pallary 1916, Wenz 1929, Esu and Girotti 1975), both of which are unavailable names.

† *Melanopsis nodosa* Pecchioli, 1864 [invalid]

Original source. Pecchioli 1864: 522, pl. 5, figs 19–20.

Type horizon. Villafranchian, Plio-Pleistocene.

Type locality. “Orciano”, Italy.

Remarks. Junior homonym of *Melanopsis nodosa* Férussac, 1822. De Stefani (1877) introduced *M. semperi* as replacement name.

† *Melanopsis costata* var. *nodosa* Brusina, 1874 [invalid]

Original source. Brusina 1874: 41.

Type horizon. Cernikian, Pliocene.

Type locality. “Podvinje (Čaplja) [Čaplja trench near Slavonski Brod]; Kovačevac; Novska; Farkašić; Dubranjec”, Croatia.

Remarks. Junior homonym of *Melanopsis nodosa* Férussac, 1822. Wenz (1928a) introduced *M. nodosula* as replacement name.

† *Melanopsis nodosa* Handmann, 1882 [invalid]

Original source. Handmann 1882: 556.

Type horizon. Pannonian, zone D, late Miocene.

Type locality. “Kottingbrunn [...] Ziegelei a”, Austria.

Remarks. Junior homonym of *Melanopsis nodosa* Férussac, 1822. Pallary (1916: 80) introduced *M. venusta* as replacement name. Wenz (1929: 2673) considered the taxon as a junior synonym of *M. bouei* Férussac, 1823.

† *Melanopsis (Canthidomus) nodosa* Doncieux, 1908 [invalid]

Original source. Doncieux 1908: 203, pl. 11, figs 11a–d.

Type horizon. Middle Lutetian, Eocene.

Type locality. “Au Nord d’Albas”, France.

Remarks. Junior homonym of *Melanopsis nodosa* Férussac, 1822.

“*Melania nodosa*” mentioned in Reeve (1860) and Brot (1874–1879: 13) [unavailable]

Locality. Not indicated.

Remarks. Nomen nudum, apparently based on an unused manuscript name from Parreyss (see Reeve 1860) or Stentz (see Brot 1874). Reeve (1860) considered the taxon as a junior synonym of *Melania hollandri* [sic]. If available, it would be a junior homonym of *Melania nodosa* Münster, 1841. Note that the latter species is certainly no Melanopsidae as it derives from the marine deposits of the Triassic St. Cassian Fm.

† *Melanopsis nodosula* Wenz, 1928

Original source. Wenz 1928a: 119.

Type horizon. Cernikian, Pliocene.

Type locality. “Podvinje (Čaplja) [Čaplja trench near Slavonski Brod]; Kovačevac; Novska; Farkašić; Dubranjec” (Brusina 1874: 41), Croatia.

Remarks. Replacement name for *M. costata nodosa* Brusina, 1874, non Férussac, 1822 (see Note 1).

† *Melanopsis inconstans* var. *nodulosa* Brusina, 1874

Original source. Brusina 1874: 39.

Type horizon. Langhian, middle Miocene.

Type locality. “Miočić”, Croatia.

Types. The syntypes are stored in the Croatian Natural History Museum, Zagreb; no number indicated (Milan et al. 1974: 92).

Remarks. Wenz (1929: 2756) and Neubauer et al. (2016b: 23) considered this taxon as a junior synonym of *Melanopsis inconstans* Neumayr, 1869.

† *Melanopsis (Lyrcea) delessei* var. *nodulosa* Magrograssi, 1928 [invalid]

Original source. Magrograssi 1928: 259, pl. 6, fig. 16.

Type horizon. Plio-Pleistocene.

Type locality. “Coo: V. Armiri” [Kos island: Armiri valley (?)], Greece.

Remarks. Junior homonym of *Melanopsis inconstans nodulosa* Brusina, 1874 (see Note 1).

Melanopsis noetlingi Bourguignat in Noetling, 1886

Original source. Noetling 1886: 817, pl. 23, fig. 6.

Type locality. “Aus dem Jarmūk bei el Hammi [...] auch bei el-Hāwījān [fossil]” [from the Yarmuk river at Al Ḥammah and as fossil near Jisr al Ḥāwī], Jordan.

Remarks. Noetling clearly referred to “Bourguignat’s new species”, which is why the authority should read as “Bourguignat in Noetling, 1886”. Heller and Sivan (2002b: 615) considered the taxon as a junior synonym of *Melanopsis multiformis* Blanckenhorn, 1897. Heller et al. (2005: 247) in turn treated it as a junior synonym of *Melanopsis obliqua* Bourguignat, 1884.

† *Melanopsis constricta nowskaensis* Wenz, 1928

Original source. Wenz 1928a: 120.

Type horizon. Cernikian, Pliocene.

Type locality. “Novska (Bukovica)” (Brusina 1897: 7), Croatia.

Types. Milan et al. (1974: 88) indicated a holotype, but it is uncertain whether the specimen was the only one Brusina had at hand when describing *M. [constricta] subcostata* (holotype by monotypy, Art. 73.1.2). The specimen is stored in the Croatian Natural History Museum, Zagreb, coll. no. 2993-639.

Remarks. Replacement name for *M. constricta subcostata* Brusina, 1897, non *M. subcostata* d'Orbigny, 1850. The name “*novskaensis*” as mentioned in Wenz (1929: 2696) is an incorrect subsequent spelling.

***Melanopsis obediensis* Picard, 1934**

Original source. Picard 1934: 110, pl. 7, figs 30–44.

Type locality. “Obedieh” [El ‘Ubeidiya], Israel.

† ***Melania obeloides* Tausch, 1886**

Original source. Tausch 1886: 7, pl. 1, figs 16–19.

Type horizon. Late Santonian–early Campanian, late Cretaceous.

Type locality. “Csingerthal bei Ajka” [Csinger valley near Ajka], Hungary.

Remarks. The species was attributed to the genus *Esperiana* by Bandel and Riedel (1994: 20). That classification was, however, doubted by Neubauer et al. (2016a: 127) based on the large stratigraphic gap between this Cretaceous species and other representatives that appeared not before the late Miocene.

***Melanopsis praemorsa* var. *obesa* Gassies, 1856**

Original source. Gassies 1856: 12, figs 11–12.

Type locality. “Dans l’Oued-Lisser, sur la route de Sidi-bel-Abess, à Tlemcen” [in the Oued Lisser (?), on the road to Sidi Bel Abbès, at Tlemcen], Algeria.

Remarks. Bourguignat (1884) attributed the authority to himself, but even listed Gassies’ record in synonymy. Apparently, he did not consider a variety as a valid name that can result in homonymy (see also the introduction).

***Melanopsis obesa* Brot, 1868 [invalid]**

Original source. Brot 1868: 57, pl. 1, figs 14–15.

Type locality. “Prope Cehejin Prov. Murcia” [near Cehegín, prov. Murcia], Spain.

Remarks. The species was described as “*Melanopsis obesa* Guirao mss.”. Although Brot (1868) denoted “(Guirao)” after the description of the species as well, it is clear from the text that Brot described it. Junior homonym of *Melanopsis obesa* Gassies, 1856. Bourguignat (1884) introduced *M. guiraoi* as replacement name.

“*Melanopsis costata* var. *obesa*” mentioned in Locard (1883a: 202, 230) and Bourguignat (1884: 141) [unavailable]

Locality. “Le Jourdain et le lac de Tibériade” [river Jordan and Sea of Galilee], Israel.

Remarks. Nomen nudum. If available, it would be a junior homonym of *Melanopsis obesa* Gassies, 1856.

† ***Melanopsis (Smendovia) doumerguei* var. *obesa* Pallary, 1901 [invalid]**

Original source. Pallary 1901a: 177, pl. 2, fig. 29.

Type horizon. Late Miocene.

Type locality. “Smendou” [Zighoud Youcef], Algeria.

Remarks. Junior homonym of *Melanopsis obesa* Gassies, 1856. Wenz (1928b) introduced *M. doumerguei plena* as replacement name.

† ***Melanopsis pygmaea obesa* Brusina, 1902 [invalid]**

Original source. Brusina 1902: pl. 5, figs 39–41.

Type horizon. Middle Pannonian, late Miocene.

Type locality. “Markuševac”, Croatia.

Types. The illustrated syntypes are stored in the Croatian Natural History Museum, Zagreb, coll. no. 2492-138, 2493-139 (Milan et al. 1974: 96).

Remarks. Junior homonym of *Melanopsis obesa* Gassies, 1856. It was considered as a junior synonym of *M. pygmaea* Hörnes, 1856 by Wenz (1929: 2814).

***Melanopsis laevigata* var. *obesula* Pallary, 1904**

Original source. Pallary 1904: 37.

Type locality. “De l’Aïn bou Smelal près de Tétouan” [‘Aïn bou Smelal (?) near Tetouan], Morocco.

***Melanopsis obliqua* Bourguignat, 1884**

Original source. Bourguignat 1884: 138.

Type locality. “Le Bélus, près de Saint-Jean-d’Acre” [in the Na’aman river, near Acre], Israel.

Remarks. Bourguignat (1884) denoted the authority as “Letourneux, 1882”, but there is no evidence that the description really derived from that author.

***Melanopsis costata* var. *obliquata* Pallary, 1939**

Original source. Pallary 1939: 91, pl. 5, fig. 34.

Type locality. “Djishr ech Chegour” [Jisr Ash-Shughur], Syria.

Remarks. Heller and Sivan (2002a: 49) considered the taxon as a junior synonym of *Melanopsis multiformis* Blanckenhorn, 1897. Heller et al. (2005: 244) in turn treated it as a junior synonym of *Melanopsis costata* (Olivier, 1804).

† ***Melanopsis oblonga* Blanckenhorn, 1897**

Original source. Blanckenhorn 1897: 137, pl. 10, fig. 25.

Type horizon. Plio-Pleistocene.

Type locality. “In der Dreissensiaschicht von Dschisr esch-Schurr” [in the *Dreissena* layers at Jisr Ash-Shughur], Syria.

† ***Melanopsis obsoleta* Fuchs, 1873**

Original source. Fuchs 1873: 20, pl. 4, figs 14–15.

Type horizon. Transdanubian, Pannonian, late Miocene.

Type locality. “Radmanest” [Rădmănești], Romania.

***Melanopsis costata* var. *obsoleta* Martens, 1874 [invalid]**

Original source. Martens 1874: 33, p. 5, fig. 39.

Type locality. “Quellen des Chabur bei Ras-el-Ain” [source of Chabur river near Ra’s al ‘Ayn], Syria.

Remarks. Junior homonym of *Melanopsis obsoleta* Fuchs, 1873.

***Melanopsis saulcyi* var. *obsoleta* Dautzenberg, 1894 [invalid]**

Original source. Dautzenberg 1894: 345.

Type locality. “Palmyre, dans un ruisseau et dans la rivière Ephéca” [Palmyra, in a stream and the river Ephéca (?)], Syria.

Remarks. Junior homonym of *Melanopsis obsoleta* Fuchs, 1873.

† ***Melanopsis obtusa* Deshayes, 1825**

Original source. Deshayes 1824–1837: 123, pl. 14, figs 22–23.

Type horizon. Cuisian, late Ypresian, Eocene.

Type locality. “Retheuil près de Pierre-Fonds” [Retheuil near Pierrefonds, Dép. Aisne], France.

Remarks. Sandberger (1872: 202) classified the species within his new genus *Copstylus* (Thiaridae), which was followed by Wenz (1929: 2570).

***Melanopsis obtusa* Pallary, 1920 [invalid]**

Original source. Pallary 1920a: 30.

Type locality. “Meknès, près de anciennes écuries” [Meknes, near the old stables], Morocco.

Remarks. Junior homonym of *Melanopsis obtusa* Deshayes, 1824.

***Melanopsis praerosa* [sic] var. *ocuticarinata* Paetel, 1888**

Original source. Paetel 1888: 403.

Type locality. “Persia” (Nevill 1884: 209), Iran.

Remarks. Nevill (1884) first mentioned the name “*acuticarinata*” [sic] for an infrasubspecific taxon (“subvariety”), which he briefly described. Paetel (1888) made the name available by ranking it as species-group name and referring to the description of Nevill (Art. 45.5.1). Although he obviously “misspelt” the name, *ocuticarinata* remains the valid spelling because Nevill’s “*acuticarinata*” is not available as species-group name.

† *Melanopsis ogerieni* Locard, 1883

Original source. Locard 1883b: 92, 140, pl. 3, figs 3–4.

Type horizon. Mammal zone MN 9–12, late Miocene.

Type locality. “Priay; [...] Niquedet” (p. 150), France.

† “*Melanopsis oliva*” mentioned in Hoeninghaus (1831: 143) [unavailable]

Horizon. Burdigalian, early Miocene (?).

Locality. “Dax”, France.

Remarks. Nomen nudum. If available, it would be a junior homonym of *M. oliva* De Cristofori & Jan, 1832.

***Melanopsis oliva* De Cristofori & Jan, 1832**

Original source. De Cristofori and Jan 1832: 4.

Type locality. “Am. mer.” [South America], indicated in the previous part of the same work (“Conchylia terrestria et fluviatilia [...]”, p. 7; there, the name is a nomen nudum).

Remarks. Certainly not a *Melanopsidae*.

***Melanopsis olivieri* Bourguignat, 1884**

Original source. Bourguignat 1884: 98.

Type locality. “Entre Ain-Taib et Alep, à Sadjour-Sou; [...] du Nahr-el-Kelb, près Beyrouth; de divers cours d’eau du Liban; de Serghaia dans l’ouady Baradah près de Damas; de la fontaine Jérémie près de Jéricho; environs de Constantinople” [Gaziantep (Turkey) and Aleppo (Syria), at Sadjour-Sou (?); in Nahr el-Kalb near Beirut (Lebanon); in various streams of Lebanon; in Serghaia (?) in River Barada near Damascus (Syria); Jeremiah Fountain (?) near Jericho (Palestine); near Istanbul (Turkey)].

† ***Melanopsis buccinoidea* var. *olivula* Grateloup, 1828**

Original source. Grateloup 1828: 137.

Type horizon. Burdigalian, early Miocene.

Type locality. “Mandillot”, France.

***Melanopsis olivula* Letourneux & Bourguignat, 1887 [invalid]**

Original source. Letourneux and Bourguignat 1887: 156.

Type locality. “À Nefta, dans les canaux d’arrosage et à El-Hammam, près Tozer” [at Nefta, in the irrigation channel, and at El-Hamma-Djerid near Tozeur], Tunisia.

Remarks. Junior homonym of *Melanopsis olivula* Grateloup, 1838. Pallary (1916: 84) was aware of the homonymy issue but did not introduce a replacement name for he considered the taxon as a junior synonym of *M. doumeti* Letourneux & Bourguignat, 1887.

† ***Melanopsis oltszakadatensis* Halaváts, 1914**

Original source. Halaváts 1914: 421, six unnumbered textfigures.

Type horizon. Sarmatian (sensu stricto), middle Miocene.

Type locality. “Graben unterhalb der Gemeinde-Baumschule östlich von Oltszakadát” [trench below communal tree nursery, east of Săcădate], Romania.

† *Melanopsis (Lyrcea) onusta* Stefanescu, 1896

Original source. Stefanescu 1896: 131, pl. 11, figs 38–41.

Type horizon. Pliocene.

Type locality. “À Plostina, à Leurda et à Stângaceana, dans la vallée de Motru; à Valea-lui-Câne, dans la vallée de Gilortu; à Breasta, à Bocovatz et à Bâzdâna, dans la vallée de Jiu; à Beceni, dans la vallée de Slanic de Buzau; à Plaiu et à Chiojdeni, dans la vallée de Râmnicu-Sarat” [at Ploștina, at Leurda and at Stângăceaua, in the valley of the river Motru; at Gilortu in the valley of the river Gilortu; at Breasta, at Bucovăț and at Bâzdâna, in the valley of the river Jiu; at Beceni, in the valley of the river Slănic; at Plaiu and at Chiojdeni, in the valley of the river Râmnic], Romania.

† *Melanopsis onychia* Brusina, 1874

Original source. Brusina 1874: 30, pl. 1, figs 3–4.

Type horizon. Cernikian, Pliocene.

Type locality. “Bečić; Podvinje (Čaplja) [Čaplja trench near Slavonski Brod]; Sibinj”, Croatia.

Types. Milan et al. (1974: 95) indicated a holotype, but it is uncertain whether the specimen was the only one Brusina had at hand (holotype by monotypy, Art. 73.1.2). The specimen is stored in the Croatian Natural History Museum, Zagreb, coll. no. 3206-852/1.

† *Melanopsis oomorpha* De Stefani, 1877

Original source. De Stefani 1877: 284.

Type horizon. Villafranchian, Plio-Pleistocene.

Type locality. “Spoleto [...], Orciano [...], fra S. Gemini e Carsoli” [Spoleto, Orciano Pisano, Carsoli near San Gemini], Italy.

Remarks. Introduced for *M. buccinoidea* sensu Férussac, 1823, non Olivier, 1801 (partim) as well as for formerly misidentified *M. narzolina*. Esu and Girotti (1975: 251) considered this taxon as a junior synonym of “*Melanopsis affinis* Férussac”, which is not an available name.

† *Melanopsis oranensis* Pallary, 1926

Original source. Pallary 1926c: 286, figs 1–4.

Type horizon. Pliocene?

Type locality. “D’un puits, profond de 25 mètres, situé dans la propriété Lamur, sur la rive gauche d’un ravinement creusé par les eaux pluviales dans une dépression de terrain, perpendiculaire au chemin d’Aïn Beïda” (p. 284) [from a well, 25 m deep, located in the village Lamur (near Oran) on the left bank of a ravine carved by rainwater in a depression in the ground, perpendicular to the path from ‘Aïn el Beïda], Algeria.

Remarks. Pallary cited in the synonymy list his paper on the fauna of the “Berbérie”, which appeared in the *Journal de Conchyliologie*. That work, however, was not published before March 1928.

† *Melanopsis orientalis* Bukowski, 1892

Original source. Bukowski 1892: 249.

Type horizon. Apolakkia/Monolithos Formation, Pliocene.

Type locality. “Rhodos” (locality specified as “Monolithos” in Bukowski 1893), Greece.

“*Melanopsis orientalis* Von dem Busch” mentioned in Mousson (1854: 50) [unavailable]

Locality. Not indicated (except from “de l’Orient”).

Remarks. Nomen nudum, listed by Mousson without any explanation.

† *Melanopsis ornata* Deshayes, 1862

Original source. Deshayes 1861–1864: 474, pl. 31, figs 27–28.

Type horizon. Sparnacian, early Ypresian, Eocene.

Type locality. “Sainceny”, France.

Remarks. Type species of *Eginea* Pacaud & Harzhauser, 2012 (Pachychilidae).

† *Paludina (Vivipara) ornata* Fuchs, 1877

Original source. Fuchs 1877: 6, pl. 1, fig. 2.

Type horizon. Pliocene.

Type locality. “Kalamaki” [near Corinth], Greece.

Remarks. Since *Paludina* Férussac, 1812 is a junior objective synonym of *Viviparus* Montfort, 1810, this species is a junior homonym of *V. ornata* Neumayr in Neumayr & Paul, 1875. Wenz (1928a) recombined the species with *Amphimelania* and introduced *A. fuchsi* as replacement name.

***Melanopsis orontis* Pallary, 1939**

Original source. Pallary 1939: 92, pl. 6, figs 70–72.

Type locality. “L’Oronte à Djisir ech Chogour” [in the Orontes at Jisir Ash-Shughur], Syria.

***Melanopsis orophila* Pallary, 1920**

Original source. Pallary 1920b: 117.

Type locality. “M. Mario: Farnesina” (Cerulli-Irelli 1914: 183), Italy.

Remarks. Introduced for *M. praemorsa* sensu Cerulli-Irelli, 1914, non Linnaeus, 1758.

† *Melanopsis (Canthidomus) oroposi* Papp, 1979

Original source. Papp 1979: 672, pl. 3, figs 1–15.

Type horizon. Mammal zone MN 10, late Miocene.

Type locality. “Milessi” [Milesi], Greece.

Types. Institute of Paleontology, University of Vienna; no number indicated.

† *Melanopsis buccinoidea* var. *ovalis* Grateloup, 1838

Original source. Grateloup 1838: 146.

Type horizon. Sparnacian, early Ypresian, Eocene.

Type locality. “Épernay”, France.

Remarks. Introduced for “*M. buccinoidea* var. a” in Férussac (1823: 149, pl. 1 [= 7], fig. 1), non Olivier, 1801.

***Melanopsis ovata* Dunker, 1862**

Original source. Dunker 1862: 150.

Type locality. “Rotoitisee” [Lake Rotoiti], New Zealand.

Remarks. Suter (1893: 139) considered the taxon as a junior synonym of *Zemelanopsis trifasciata* (Gray in Dieffenbach, 1843).

† *Melanopsis ovicona* Deffner & Fraas, 1877

Original source. Deffner and Fraas 1877: 16.

Type horizon. Mammal zone MN 6, middle Miocene.

Type locality. “Trendel” [in Nördlinger Ries], Germany.

Remarks. Wenz (1929: 2763), who erroneously considered the taxon as a nomen nudum, listed it in synonymy of *Melanopsis kleinii* Kurr, 1856.

***Melanella ovoidaea* Bourguignat, 1884**

Original source. Bourguignat 1884: 22.

Type locality. “Le Danube au-dessus de Routschouk (Bulgarie)” [in the Danube river below Ruse], Bulgaria.

Remarks. Note that Bourguignat denoted the authority as “Bourguignat, 1879”.

***Melanopsis nodosa* var. *ovoidalis* Cerulli-Irelli, 1914**

Original source. Cerulli-Irelli 1914: 184, pl. 15 (47), figs 5–6.

Type locality. “M. Mario: Farnesina; Acuatraversa”, Italy.

Remarks. Girotti (1972: 232) considered the taxon as a junior synonym of “*Melanopsis affinis* Férussac”, which is not an available name.

† *Melanopsis ovosimilis* Willmann, 1981

Original source. Willmann 1981: 143.

Type horizon. Vasilios Formation, middle Tortonian, late Miocene.

Type locality. “Dermen Deressi 400 m westlich des Asklepion, ca. 3 km westlich von Kos-Ort/Kos” [Dermen Deressi 400 m west of Asklepion, ca. 3 km west of Kos city, Kos Island], Greece.

Types. Geological-Paleontological Institute, University of Kiel, Germany; no number indicated.

***Melanopsis ovula* Bourguignat, 1884**

Original source. Bourguignat 1884: 162.

Type locality. “Le Guadalquivir entre Séville et Cordoue” [in the Guadalquivir river between Sevilla and Córdoba], Spain.

† *Melanopsis ovularis* Watelet, 1853

Original source. Watelet 1853: 18, pl. 2, figs 18–19.

Type horizon. Cuisian, late Ypresian, Eocene.

Type locality. “Mercin” [Mercin-et-Vaux], France.

Remarks. Watelet attributed the authority to Deshayes, but there is no evidence that the description really derived from that author.

***Melanopsis ovum* Bourguignat, 1884**

Original source. Bourguignat 1884: 143.

Type locality. “See Tiberias” (Kobelt 1880: 17) [Sea of Galilee], Israel.

Remarks. Appeared first as a nomen nudum in Locard (1883a: 202). Introduced for “*Melanopsis costata* var.” in Kobelt (1880: 17, pl. 188, fig. 1906). Heller et al. (2005: 246) considered the species as a junior synonym of *Melanopsis jordanica* Roth, 1839.

† ***Melanopsis oxyacantha* Brusina, 1902**

Original source. Brusina 1902: pl. 7, figs 6–7.

Type horizon. Transdanubian, Pannonian, late Miocene.

Type locality. “Tihany”, Hungary.

Types. The illustrated syntypes are stored in the Croatian Natural History Museum, Zagreb, coll. no. 2534-180/1-2 (Milan et al. 1974: 95).

Remarks. The name “*oxicanthe*” as mentioned in Miletic-Spajic (1961: 252) is an incorrect subsequent spelling.

† ***Melanopsis (Lyrcea) pachecoi* Royo Gómez, 1922**

Original source. Royo Gómez 1922: 108.

Type horizon. Late Miocene.

Type locality. “Sierra de la Pinada en Sayatón (Guadalajara)”, Spain.

***Melanopsis pachya* Pallary, 1939**

Original source. Pallary 1939: 86, pl. 5, figs 10, 12.

Type locality. “Dans les sources de Mézéril, au N. O. de Derâa (Syrie méridionale)” [in the sources of the Muzayrib, northwest of Dar’a], Syria.

***Microcolpia pachystoma* Bourguignat, 1884**

Original source. Bourguignat 1884: 66.

Type locality. “La Save près d’Agram” [Sava river at Zagreb], Croatia.

Remarks. Starobogatov et al. (1992: 65) considered the species as a junior synonym of *Microcolpia cornea* (Pfeiffer, 1828).

† *Melanopsis (Melanosteira) pagodaeformis* Pavlović, 1927

Original source. Pavlović 1927: 62, pl. 7, fig. 11.

Type horizon. Middle Pannonian, late Miocene.

Type locality. “Из Караџача” [from Karagača near Vrčin], Serbia.

Types. The illustrated syntype is stored in the Natural History Museum, Belgrade, coll. no. 213 (Milošević 1962: 23).

† *Melanopsis pallaryi* Wenz, 1919

Original source. Wenz 1919a: 73.

Type horizon. Early Pleistocene.

Type locality. “Du puits Karoubi” (Pallary 1901a: 178) [from the well Karoubi], Algeria.

Remarks. Replacement name for *M. acuminata* Pallary, 1901, non Gümbel, 1861.

Melanopsis doriae pallida Biggs, 1962

Original source. Biggs 1962: 71.

Type locality. “Bagh-i-Zerisf near Kerman”, Iran.

Melanopsis palmyrensis Pallary, 1939

Original source. Pallary 1939: 96, pl. 6, figs 60–62.

Type locality. “Palmyre”, Syria.

Remarks. Heller et al. (2005: 248) considered the species as a junior synonym of *Melanopsis saulcyi* Bourguignat, 1853.

† *Melanopsis panciciana* Brusina, 1874

Original source. Brusina 1874: 44.

Type horizon. Langhian, middle Miocene.

Type locality. “Ribarić”, Croatia.

Types. The illustrated syntypes are stored in the Croatian Natural History Museum, Zagreb, coll. no. 3660-1300/1-2 (Milan et al. 1974: 95).

Remarks. Neubauer et al. (2013: 135) considered this taxon as a junior synonym of *Melanopsis lyrata* Neumayr, 1869.

† *Melanopsis neumayri* var. *papiolensis* Almera & Bofill y Poch in Almera, 1894

Original source. Almera 1894: 171, pl. 4, figs 5a–b.

Type horizon. Late Messinian–early Zanclean, late Miocene–early Pliocene.

Type locality. “Castellbisbal, Papiol”, Spain.

† *Melanopsis papkesiensis* Bandel, 2000

Original source. Bandel 2000: 162, figs 52–85.

Type horizon. Late Pannonian, late Miocene.

Type locality. “Sand pit near Papkesi [Papkeszi]”, Hungary.

Types. Geological-Palaeontological Institute and Museum University of Hamburg, coll. no. 4269.

† *Melanoptychia paradoxa* Brusina, 1892

Original source. Brusina 1892: 144.

Type horizon. Middle Pannonian, late Miocene.

Type locality. “Markuševac”, Croatia.

Types. The illustrated syntypes are stored in the Croatian Natural History Museum, Zagreb, coll. no. 2540-186/1-4 (Milan et al. 1974: 100).

Melanopsis pardalis Brot, 1878

Original source. Brot 1874–1879: 373, pl. 38, fig. 5.

Type locality. “Krain, im Gurkfluss” (Clessin 1890: 686) [in the river Gurk], Austria.

Remarks. Brot (1878) introduced it as “*Melanopsis pardalis*. Mhlf. Mss.” in synonymy of *Hemisinus esperi* (Férussac, 1823) [currently in *Esperiana*]. Clessin (1890) made the name available by treating it as valid name (see Note 2).

† *Melanopsis parkinsoni* Deshayes, 1825

Original source. Deshayes 1824–1837: 123, pl. 17, figs 3–4.

Type horizon. Cuisian, late Ypresian, Eocene.

Type locality. “Cuise-la-Mothe” [Cuise-la-Motte], France.

Remarks. Wenz (1929: 2566) considered the species as a junior synonym of *Coprostylus albidus* (Lamarck, 1822) (Thiaridae).

***Melania (Melanopsis) parreyssii* Philippi, 1847**

Original source. Philippi 1847: 176, pl. 4, fig. 15.

Type horizon. Late Pleistocene to Recent.

Type locality. “Hungaria” [no locality indicated], Hungary? (at that time, the kingdom Hungary covered large parts of Romania and northern Croatia).

Remarks. Philippi attributed the authority to Johann Georg Megerle von Mühlfeld, apparently based on an “in schedis” determination. The name “*parreyssi*” as mentioned in Brusina (1903), Kormos (1905) and Paucă (1937) is an incorrect subsequent spelling (see also Neubauer et al. 2014d).

† ***Melanopsis parva* Pallary, 1916**

Original source. Pallary 1916: 78.

Type horizon. Burdigalian, early Miocene (?).

Type locality. “Dax”, France.

Remarks. Pallary (1916, 1920) erroneously treated “*parva*” as available name attributed to Férussac (1823), who solely used the word as descriptive term (Latin “small”). The name became nevertheless available from Pallary (1916) who associated the name with an illustration in Férussac (1823: pl. 8, fig. 5). Cossmann and Peyrot (1918: 485) considered the taxon as a junior synonym of *Melanopsis subbuccinoides* d’Orbigny, 1852.

† ***Melanopsis olivula* var. *parvula* Grateloup, 1838**

Original source. Grateloup 1838: 143, pl. 4, fig. 56.

Type horizon. Burdigalian, early Miocene.

Type locality. “Dax. [...] Mandillot”, France.

***Melania parvula* Brot, 1874**

Original source. Brot 1874–1879: 13, pl. 1, figs 2–2a.

Type locality. “Krain” [Carniola, a historical region that comprised parts of present-day Slovenia; no locality indicated].

Remarks. Appeared first as a nomen nudum in Brot (1862). The name was not found in Rossmässler (1839), to whom Brot (1862, 1874) referred.

***Melanopsis jordanica* var. *parvula* Bourguignat, 1884 [invalid]**

Original source. Bourguignat 1884: 142.

Type locality. “Le Jourdain, à 4 kilomètres au-dessus de la Mer Morte; Ain-el-Mellaha, dans la plaine du Bahr-el-Houlé” [Jordan river, 4 km north of the Dead Sea; Ain Mallahah, in the plains of the Hula valley], Israel.

Remarks. Junior homonym of *Melanopsis olivula parvula* Grateloup, 1838 (see Note 1). Heller et al. (2005: 245) considered the variety a synonym of *M. lampra* Bourguignat, 1884.

† *Melania patula* Bellardi & Michelotti, 1841

Original source. Bellardi and Michelotti 1841: 163, pl. 7, figs 8–9.

Type horizon. Late Miocene.

Type locality. “Del Tortonese” [from the region of Tortona], Italy.

Remarks. Bellardi and Michelotti attributed the authority to Bonelli (1827), which is an unpublished collection catalogue. Considered to belong in the genus *Amphimelania* by Wenz (1929: 2876).

† *Melanopsis (Lyrcaea)* [sic] *vindobonensis paucai* Lubenescu, 1981

Original source. Lubenescu 1981: 176, pl. 12, figs 8–12.

Type horizon. Middle Pannonian, late Miocene.

Type locality. “Comuna Vingard, 15 km NE de Sebeș-Alba”, Romania.

Types. Institute of Geology and Geophysics, University of Bucharest; no number indicated.

† *Melanopsis pauli* Bandel & Riedel, 1994

Original source. Bandel and Riedel 1994: 19, pl. 12, figs 2–6.

Type horizon. Late Santonian–early Campanian, late Cretaceous.

Type locality. “Csingertal, near Ajka (Bakony Mountains)”, Hungary.

Types. Geological-Paleontological Department, Natural History Museum Vienna, Austria, coll. no. 1994/148.

† *Melanopsis paulovici* Bourguignat, 1880

Original source. Bourguignat 1880: 37.

Type horizon. Langhian, middle Miocene.

Type locality. “Vallée de la Cettina” [Cetina river valley], Croatia.

† *Melanopsis pavlovici* Brusina, 1902

Original source. Brusina 1902: pl. 5, figs 45–47.

Type horizon. Pannonian, zone D–E, late Miocene.

Type locality. “Ripanj”, Serbia.

Types. The illustrated syntypes are stored in the Croatian Natural History Museum, Zagreb, coll. no. 2495-141/1-3 (Milan et al. 1974: 95).

Remarks. Appeared first as a nomen nudum in Brusina (1893). Not a homonym of *M. paulovici* [sic] as considered by Pallary (1916), who invalidly introduced *Melanopsis ripajensis* as replacement name. Note that Art. 58.4. (regarding the use of u or v for the same Latin letter) does not apply here, because the names are not based on Latin words but on the names of two different persons.

† *Melania pecchiolii* Hörnes, 1856

Original source. Hörnes 1851–1856: 604, pl. 49, fig. 24.

Type horizon. Badenian, middle Miocene.

Type locality. “Forchtenau in Ungarn” [Forchtenstein, Burgenland], Austria.

Remarks. Considered to belong in the genus *Amphimelania* by Wenz (1929: 2877). Note that Wenz gave the name as “*pecchioli*”, which is an incorrect subsequent spelling.

Melanopsis pechaudi Bourguignat, 1884

Original source. Bourguignat 1884: 160.

Type locality. “De la source de la Moulouiah, au nord de Lalla-Maghnia, près des frontières du Maroc” [in the source of the river Moulouya, north of Maghnia, near the border to Morocco], Morocco or Algeria.

Remarks. Note that Bourguignat denoted the authority as “Bourguignat, 1882”.

† *Melanopsis pedemontana* Sacco, 1889

Original source. Sacco 1889: 66, pl. 2, figs 1–8.

Type horizon. Late Burdigalian, early Miocene.

Type locality. “Colli torinesi, [...] presso Tetti Varetto” [Torino hills, near Tetti Varetto], Italy.

Melanopsis penchinati Bourguignat, 1868

Original source. Bourguignat 1868: 432.

Type locality. “Agora, en Aragon (Espagne)”, Spain.

Remarks. The name “*pinchinati*” as mentioned in Brot (1879: 440) is an incorrect subsequent spelling.

† *Melanopsis pentagona* Brusina, 1892

Original source. Brusina 1892: 138.

Type horizon. Middle Pannonian, late Miocene.

Type locality. “Markuševac”, Croatia.

Types. The illustrated syntypes are stored in the Croatian Natural History Museum, Zagreb, coll. no. 2532-178/1-4 (Milan et al. 1974: 95).

Microcolpia peracuta Bourguignat, 1884

Original source. Bourguignat 1884: 63.

Type locality. “Environs de Krapina-Toeplitz (Croatie)” [surroundings of Krapinske toplice], Croatia.

Remarks. Starobogatov et al. (1992: 65) considered the species as a junior synonym of *Microcolpia acicularis* (Férussac, 1823).

Melanopsis praemorsa f. *perbrevis* Pérès, 1946 [invalid]

Original source. Pérès 1946: 113.

Type locality. Not indicated.

Remarks. First mentioned as nomen nudum in Pérès (1939), but described and made available by Pérès (1946). In both works Pérès obviously used the name not as a separate taxon but rather as descriptive term to fit existing species into his morphological concept. The name is in fact a junior objective synonym of several different species.

† *Melanopsis percallosa* Sandberger, 1875

Original source. Sandberger 1870–1875: 574.

Type horizon. Middle Miocene.

Type locality. “Locle”, Switzerland.

Remarks. Wenz (1929: 2763) considered the taxon as a junior synonym of *Melanopsis kleinii* Kurr, 1856.

† *Melanopsis percarinata* Förster, 1892

Original source. Förster 1892: 225, pl.11, fig. 6.

Type horizon. Early Rupelian, Oligocene.

Type locality. “Kleinkems, [...] Tagolsheim”, Germany.

Melanopsis percarinata Pallary, 1918 [invalid]

Original source. Pallary 1918: 150.

Type locality. “Dans les séguis du Guers” [in the irrigation channel of Guers], Morocco.

Remarks. Junior homonym of *Melanopsis percarinata* Förster, 1892.

† *Melanopsis pergamena* Calvert & Neumayr, 1880

Original source. Calvert and Neumayr 1880: 375, pl. 2, figs 18–19.

Type horizon. Late Sarmatian (sensu lato), Khersonian, late Miocene.

Type locality. “Renkiöi” [north of İntepe], Turkey.

Remarks. The name “*pergamenica*” as mentioned in Wenz (1929: 2804) is an incorrect subsequent spelling.

† *Melanopsis permutabilis* Pallary, 1920

Original source. Pallary 1920b: 118.

Type horizon. Cernikian, Pliocene.

Type locality. “Bečić; Podvinje (Čaplja) [Čaplja trench near Slavonski Brod]; Sibinj; Kovačevac; Novska; Moslavina” (Brusina 1874: 41–42), Croatia.

Remarks. Probably a junior synonym: introduced for *M. costata* sensu Brusina, 1874, non Olivier, 1804, which has been considered to be the same species as *M. costata* sensu Neumayr, 1869 (e.g., Neumayr and Paul 1875: 41). For that misidentified record, Brusina (1884a: 168) had already introduced *M. croatica* as new name (see also discussion of *M. pseudocostata* Oppenheim, 1890). Not included in the Fossilium Catalogus of Wenz (1929).

Melanopsis douttei var. *perornata* Pallary, 1920

Original source. Pallary 1920c: 150, pl. 4, fig. 3.

Type locality. “Oued Fès, à 1.500 mètres en amont de la ville” [Oued Fes, at 1.500 m above the city], Morocco.

† *Melanopsis petkovici* Pavlović, 1931

Original source. Pavlović 1931: 19, pl. 10, figs 19–20.

Type horizon. Badenian, middle Miocene.

Type locality. “Из села Врмџе” [Vrmdža], Serbia.

Types. The illustrated syntype is stored in the Natural History Museum, Belgrade, coll. no. 2888 (Milošević 1962: 24).

Remarks. Appeared first as a nomen nudum in Pavlović (1922: 49).

† *Melanopsis petrovici* Brusina, 1893

Original source. Brusina 1893: 58 (only described in Italian part).

Type horizon. Transdanubian, Pannonian, late Miocene.

Type locality. “Orešac”, Serbia.

Types. Lectotype designation by Milan et al. (1974: 84) invalid: it is uncertain whether the specimen illustrated by Brusina (1902) was part of the original type series. The specimen is stored in the Croatian Natural History Museum, Zagreb, coll. no. 2484-130.

***Fagotia pfeifferi* Bourguignat, 1884**

Original source. Bourguignat 1884: 36.

Type locality. “Rivières de Carniole; la Save à Steinbrück” [rivers in Carniola, a historical region that comprised parts of present-day Slovenia; in the Sava river near Zidani Most], Slovenia.

Remarks. Note that Bourguignat denoted the authority as “Bourguignat, 1880”. Starobogatov et al. (1992: 60) considered the species as a junior synonym of *Fagotia* [= *Esperiana*] *acroxia* Bourguignat, 1884.

† *Melanopsis phanesiana* Bukowski, 1892

Original source. Bukowski 1892: 249.

Type horizon. Salakos Formation, Pliocene.

Type locality. “Rhodos” (locality specified as “Kalavarda” in Bukowski 1893), Greece.

Remarks. Willmann (1981: 127) considered this taxon as a synonym of *Melanopsis vandevelde* Bukowski, 1892.

***Melanopsis phoeniciaca* Bourguignat, 1884**

Original source. Bourguignat 1884: 133.

Type locality. “Dans le Bélus [près de Saint-Jean-d’Acre, en Syrie]” [in the Na’aman river, near Acre], Israel.

Remarks. The spelling “*phaeniciaca*” on p. 76 is obviously a typesetting mistake regarding the ligature (æ instead of œ). Heller et al. (2002: 596) considered the species (misspelt as “*phoeneciaca*”) a junior synonym of *Melanopsis costata* (Olivier, 1804). Heller et al. (2005: 244) treated it as a junior synonym of *Melanopsis lampra* Bourguignat, 1884.

† “*Melanopsis pichleri*” mentioned in (Pichler 1856: 735–736)

Horizon. Late Cretaceous.

Locality. “Auf zwei Puncten der Brandenberger Ache. Ungefähr $\frac{3}{4}$ Stunde nördlich von Binneck [...]. Etwa eine Viertelstunde unter Binneck [...] am linken Ufer der Ache” [At two points along the Brandenberger Ache: about three quarters of an hour north of Pinegg and a quarter of an hour south of Pinegg in Tyrol], Austria.

Remarks. First mentioned as nomen nudum in Pichler (1856), who attributed the authority to Hörnes. It was validly described by Stoliczka (1860: 487, pl. 1, figs 6–9), but not considered a Melanopsidae there or later (see, e.g., Yen 1958: 206).

† *Melanopsis pictus* Hörnes, 1856

Original source. Hörnes 1851–1856: 600, pl. 49, fig. 14.

Type horizon. Badenian, middle Miocene.

Type locality. “Grund”, Austria.

Remarks. After Wenz (1929: 2505) this species belongs in the genus *Semisinus* P. Fischer, 1885, which is an unjustified emendation of *Hemisinus* Swainson, 1840 (Thiaridae).

Fagotia pilariana Bourguignat, 1884

Original source. Bourguignat 1884: 46.

Type locality. “Dans la Save à Agram et à Sissek. Dans la rivière, entre Plaski et Ostaria (Croatie)” [Sava river at Zagreb and Sisak. In the river between Plaški and Oštarije], Croatia.

Remarks. Note that Bourguignat denoted the authority as “Bourguignat, 1882”. Starobogatov et al. (1992: 60) considered the species as a junior synonym of *Fagotia* [= *Esperiana*] *servainiana* Bourguignat, 1884.

Melanella pilariana Bourguignat, 1884

Original source. Bourguignat 1884: 15.

Type locality. “En Croatie, au pont de la Save à Agram, ainsi qu’à Steinbrück; [...] en Bosnie, dans les rivières de Zenica et de la Migliaska à Sérajewo” [in Croatia, at the bridge of the Sava river in Zagreb, as well as Zidani Most (Slovenia); in Bosnia, in the rivers Zenica and Miljacka near Sarajevo].

Remarks. Appeared first as a nomen nudum in Servain (1884: 379). Note that Bourguignat denoted the authority as “Bourguignat, 1880”.

† *Melanopsis planoidea* Aldrich, 1895

Original source. Aldrich 1895: 15, pl. 2, fig. 7.

Type horizon. Early Eocene.

Type locality. “Gregg’s Landing, Alabama”, United States.

Types. Harris (1899: 77) designated a lectotype, which is stored in the collection of the National Museum of Natural History, Smithsonian Institution, Washington, DC, coll. no. 638788 (Palmer and Brann 1966: 755).

Remarks. Palmer and Brann (1966: 755) considered the taxon as a junior synonym of *Melanopsis anita* Aldrich, 1886. According to Allmon (1990: 63), neither of both species belongs to the Melanopsidae.

† *Melanopsis doumerguei plena* Wenz, 1928

Original source. Wenz 1928b: 219.

Type horizon. Late Miocene.

Type locality. “Smendou” [Zighoud Youcef], Algeria.

Remarks. Replacement name for *M. doumerguei obesa* Pallary, 1901, non Gassies, 1856 (see Note 1).

† *Fagotia (Sasykiana) plena* Gozhik in Gozhik & Datsenko, 2007

Original source. Gozhik and Datsenko 2007: 98, pl. 89, figs 2–3.

Type horizon. Pliocene.

Type locality. “Оз. Сасык” [Lake Sasyk], Ukraine.

Types. Institute of Geological Sciences of the National Academy of Sciences of Ukraine, coll. no. 4603.

† *Melanoptychia pleuroplagia* Bourguignat, 1880

Original source. Bourguignat 1880: 32.

Type horizon. Langhian, middle Miocene.

Type locality. “Vallée de la Cettina” [Cetina river valley], Croatia.

***Melanopsis pleuroplagia* Bourguignat, 1884 [invalid]**

Original source. Bourguignat 1884: 155.

Type locality. “Dans le Viar et le Carbanès, entre Cordoue et Séville (Espagne)” [in the Río Viar and Río Corbones, between Córdoba and Sevilla], Spain.

Remarks. Junior secondary homonym of *Melanoptychia pleuroplagia* Bourguignat, 1880. It was considered as a junior synonym of *M. sevillensis* by Pallary (1916: 86).

***Melanopsis pleurotomoidaea* Bourguignat, 1884**

Original source. Bourguignat 1884: 105.

Type locality. “Ruisseaux de l’île d’Ivice, dans les Baléares” [stream on the island of Ibiza], Spain.

† *Melanopsis (Melanoptychia) pleurotomoides* Pavlović, 1927

Original source. Pavlović 1927: 60, pl. 7, figs 5–6.

Type horizon. Pannonian, zone D–E, late Miocene.

Type locality. “У zaseoky Рамачу [...] у Рипњу” [from the village Ramača and from Ripanj], Serbia.

Types. The illustrated syntype is stored in the Natural History Museum, Belgrade, coll. no. 794 (Milošević 1962: 24).

***Melanopsis barbini* var. *plicata* Pallary, 1911**

Original source. Pallary 1911: 130, [unnumbered plate], fig. 6.

Type locality. “Tout près d’Oudjda, à 4 kilom. S.-E., sourdent les belles sources de Sidi-Yahia qui alimentent une véritable oasis, puis la ville d’Oudjda, et vont finalement se déverser dans l’oued Isly” [near Oujda, 4 km southeast, at the sources of Sidi Yahya that feed an oasis and the city of Oujda, and ultimately will flow into the Oued Isly], Morocco.

***Melanopsis subimpressa* var. *plicata* Pallary, 1928 [invalid]**

Original source. Pallary 1928a: 263.

Type locality. “Dans une source tiède à proximité de la Moulouïa” [in a warm source close to the river Moulouya], Morocco.

Remarks. Junior homonym of *Melanopsis plicata* Pallary, 1911.

***Melanopsis (Mesopotamia) khabourensis* var. *plicata* Pallary, 1939 [invalid]**

Original source. Pallary 1939: 103, pl. 5, fig. 19.

Type locality. “Ras el ‘Ain du Khabour” [Chabur river near Ra’s al ‘Ayn], Syria.

Remarks. Junior homonym of *Melanopsis barbini plicata* Pallary, 1911 (see Note 1). Heller et al. (2005: 254) considered the variety as a junior synonym of *Melanopsis infracincta* Martens, 1874.

† *Melanopsis plicatella* Neumayr, 1880

Original source. Neumayr 1880c: 477, pl. 7, fig. 2.

Type horizon. Pannonian, late Miocene; late Burdigalian to early Langhian, early–middle Miocene.

Type locality. “Posušje; [...] Seonica”, Bosnia and Herzegovina.

† *Melanopsis inconstans* var. *plicatula* Brusina, 1874

Original source. Brusina 1874: 39.

Type horizon. Langhian, middle Miocene.

Type locality. “Miočić”, Croatia.

Types. Milan et al. (1974: 96) indicated a holotype, but it is uncertain whether the specimen is part of the original type series and whether it was the only one Brusina had at hand (holotype by monotypy, Art. 73.1.2). The specimen is stored in the Croatian Natural History Museum, Zagreb, coll. no. 2976-622/1-2.

Remarks. Neubauer et al. (2016b: 23) considered this taxon as a junior synonym of *Melanopsis inconstans* Neumayr, 1869.

† “*Melanopsis themaki* var. *plicatula*” mentioned in Brusina (1903: 110) [unavailable]

Horizon. Late Pleistocene–early Holocene.

Locality. “Bischofsbad” [Püspökfürdő, Băile 1 Mai, Lake Pețea], Romania.

Remarks. Nomen nudum. If available, it would be a junior objective synonym of *M. themaki*, because Brusina (1903) indicated it as the typical form of the species, as well as a junior homonym of *Melanopsis inconstans plicatula* Brusina, 1874 (see Note 1). Neubauer et al. (2014d: 125) considered this taxon as a junior synonym of *Microcolpia parreyssii sikorai* (Brusina, 1903).

† *Melanopsis vidovici* var. *plicatula* Brusina, 1903 [invalid]

Original source. Brusina 1903: 113.

Type horizon. Late Pleistocene–early Holocene.

Type locality. “Bischofsbad” [Püspökfürdő, Băile 1 Mai, Lake Peșea], Romania.

Remarks. Junior homonym of *Melanopsis inconstans plicatula* Brusina, 1874 (see Note 1). Neubauer et al. (2014d: 125) considered this taxon as a junior synonym of *Microcolpia parreyssii sikorai* (Brusina, 1903).

† *Melanopsis (Canthidomus) plicatula* Handmann, 1887 [invalid]

Original source. Handmann 1887: 30, pl. 7, figs 1–3.

Type horizon. Pannonian, zone B–D, late Miocene.

Type locality. “Leobersdorf”, Austria.

Remarks. Originally the gender was indicated as masculine (“*plicatulus*”), but *Melanopsis* is feminine, which is why the name must be “*plicatula*”. Junior homonym of *Melanopsis inconstans plicatula* Brusina, 1874 (see Note 1). Pallary (1916: 84) introduced *M. similis* as replacement name. Wenz (1929: 2674) listed both names in synonymy of *M. bouei* Férussac, 1823.

† *Melanopsis polita* Pallary, 1916

Original source. Pallary 1916: 79.

Type horizon. Sparnacian, early Ypresian, Eocene.

Type locality. “Pourcy” (Cossmann 1909: captions of pl. 3), France.

Remarks. Introduced for *M. buccinoidea* sensu Cossmann, 1909 (p. 171, pl. 3, figs 21–22) and Cossmann, 1910 (in Cossmann and Pissarro 1906–1913: pl. 19, fig. 118–1). Wenz (1929: 2662) considered it as a junior synonym of *M. antediluviana*.

Melanopsis vanrossomi var. *polita* Pallary, 1936 [invalid]

Original source. Pallary 1936: 59.

Type locality. “Tanalt”, Morocco.

Remarks. Junior homonym of *Melanopsis polita* Pallary, 1916.

† *Melanopsis polyptycha* Neumayr, 1880

Original source. Neumayr 1880b: 294, pl. 1, fig. 26.

Type horizon. Plio-Pleistocene.

Type locality. “Zwischen Pyllé und Antimachia” [between Pýli and Antimácheia, Kos Island], Greece.

Remarks. The name “*polyptychia*” as mentioned in Wenz (1929: 2806) is an incorrect subsequent spelling.

† *Ancillaria (Amalda?) pomahaka* Hutton, 1873

Original source. Hutton 1873: 6.

Type horizon. Pomahaka Formation, Duntroonian (? = late Oligocene).

Type locality. “Pomahaka, Otago” [Pomahaka River], New Zealand.

Remarks. The species was attributed to the *Melanopsis* (*Stilospirula?*) by Beu and Maxwell (1990: 141).

Melanella ponderosa Bourguignat, 1884

Original source. Bourguignat 1884: 23.

Type locality. “En Carmiole [sic], en Croatie, et en Dalmatie, notamment dans la Cettina” [in Carniola (a historical region that comprised parts of present-day Slovenia), Croatia and Dalmiata, especially in the Cetina river].

Remarks. Appeared first as a nomen nudum in Villa and Villa (1841: 36) and Brusina (1867: 86). Note that Bourguignat denoted the authority as “Bourguignat, 1877”.

† *Melanopsis (Mingrelicana) pontica* Gozhik in Gozhik & Datsenko, 2007

Original source. Gozhik and Datsenko 2007: 92, pl. 96, figs 2–3.

Type horizon. Middle Pontian (Dacian Basin), late Miocene.

Type locality. “Вулканешты” [Vulcănești], Moldova.

Types. Institute of Geological Sciences of the National Academy of Sciences of Ukraine, coll. no. 3190.

† *Melanopsis porumbari* Porumbaru, 1881

Original source. Porumbaru 1881: 29, pl. 7, fig. 5, pl. 9, fig. 3.

Type horizon. Early–middle Romanian, Pliocene.

Type locality. “Cretzesi et Podari” [Crețești and Podari], Romania.

Remarks. Porumbaru attributed the authority to Brusina, but there is not clear evidence that the description really derived from that author. The name “*porumbarui*” as mentioned in Fontannes (1887: 336), Stefanescu (1896: 134) and Wenz (1929: 2806) is an incorrect subsequent spelling.

† *Melanopsis impressa posterior* Papp, 1953

Original source. Papp 1953b: 133, pl. 9, figs 19–23.

Type horizon. Pannonian, zone B, late Miocene.

Type locality. “Leobersdorf Sandgrube” [Leobersdorf sand pit], Austria.

Remarks. Appeared first as *nomen nudum* in Papp (1951: 107, 108, 124, 127). Although he mentioned differences from other melanopsids, he did not state any morphological characters.

† *Melanopsis eleis posterior* Schütt in Symeonidis et al., 1986 [invalid]

Original source. Symeonidis et al. 1986: 342, pl. 4, figs 8–9.

Type horizon. Gelasian, early Pleistocene.

Type locality. “Weganrisse und Regenrisse 3 Km N Antirrhion in Akarnanien” [outcrops 3 km north of Antirrhio], Greece.

Types. Senckenberg Forschungsinstitut und Naturmuseum Frankfurt; no number indicated.

Remarks. Junior homonym of *Melanopsis impressa posterior* Papp, 1953 (see Note 1). Neubauer et al. (2014a) introduced *M. gearyae* as replacement name.

Melanopsis potamactebia Bourguignat, 1870

Original source. Bourguignat 1870: 67.

Type locality. “Dans le Danube. [...] non-seulement de Brahilov, mais encore des environs de Belgrade” [in the Danube; not only from Brăila (Romania), but also from the surroundings of Belgrade (Serbia)].

† *Melanopsis (Coptostylus) pourcyensis* Cossmann, 1907

Original source. Cossmann 1907: 236, pl. 7, fig. 118–12.

Type horizon. Sparnacian, early Ypresian, Eocene.

Type locality. “Pourcy”, France.

Remarks. *Coptostylus* is at present considered to belong in the family Thiaridae (Pacaud and Le Renard 1995: 156).

Microcolpia praeclara Bourguignat, 1884

Original source. Bourguignat 1884: 54.

Type locality. “La Save à Agram, à Sissek et près de Belgrade” [Sava river at Zagreb, at Sisak (Croatia) and near Belgrade (Serbia)].

Remarks. Starobogatov et al. (1992: 65) considered the species as a junior synonym of *Microcolpia cornea* (Pfeiffer, 1828).

† *Melanopsis praecursor* Schütt in Schütt & Ortal, 1993

Original source. Schütt and Ortal 1993: 93, pl. 2, fig. 30.

Type horizon. Early Pleistocene.

Type locality. “The type-locality of the Upper Pliocene sediments or the ‘Erq el-Ahmar Formation in the Jordan Valley south of the Sea of Galilee” [i.e. ‘Erq el-Ahmar (also known as Gesher)], Israel.

Types. Paleontology Collection of the Hebrew University of Jerusalem; no number indicated.

Buccinum praemorsum Linnaeus, 1758

Original source. Linnaeus 1758: 740.

Type locality. “In Europa australiore” [southern Europe].

Remarks. Sometimes incorrectly listed as the type species of *Melanopsis* (e.g., Wenz 1929: 2647; Heller et al. 1999: 56; Esu and Girotti 2015b: 150).

Buccinum praerosum Linnaeus, 1767 [invalid]

Original source. Linnaeus 1767: 1203.

Remarks. The name *Buccinum praerosum*, commonly combined as *Melanopsis praerosa*, is an unjustified emendation and therefore a junior objective synonym of *Buccinum praemorsum* Linnaeus, 1758.

† “*Melanopsis pregorceisci*” mentioned in Gillet et al. (1978: 58) [unavailable]

Horizon. Khersonian, late Sarmatian, late Miocene.

Locality. “Hauptstraße bei Avcılar, nahe der Abzweigung nach Ambarliköy” [main road at Avcılar, near the diversion to Ambarlı, west of Istanbul], Turkey.

Remarks. Nomen nudum; mentioned as “new species” in Gillet et al. (1978) but left undescribed.

Melanopsis princeps Lea, 1837

Original source. Lea 1837: 82, pl. 19, fig. 74.

Type locality. “Cape of Good Hope”, South Africa.

Remarks. Type species of *Faunopsis* Gill, 1863. It was considered as a junior synonym of *Faunus ater* (Linnaeus, 1758), the type species of the genus *Faunus* Montfort, 1810 (Pachychilidae), by Starmühlner (1974: 132).

† *Melanopsis (Canthidomus) prionodonta* Handmann, 1887

Original source. Handmann 1887: 33, pl. 7, fig. 15.

Type horizon. Pannonian, zone B–D, late Miocene.

Type locality. “Leobersdorf”, Austria.

Remarks. Wenz (1929: 2675) considered the taxon as a junior synonym of *Melanopsis bouei* Férussac, 1823.

† *Melanopsis proboscidea* Deshayes, 1862

Original source. Deshayes 1861–1864: 471, pl. 31, figs 18–24.

Type horizon. Bartonian, Eocene.

Type locality. “Chery-Chartreuve”, France.

† *Melanopsis (Martinia) martiniana* var. *proclivis* Handmann, 1887

Original source. Handmann 1887: 25, pl. 4, figs 1–2.

Type horizon. Pannonian, zone B–D, late Miocene.

Type locality. “Leobersdorf”, Austria.

Remarks. Papp (1953b: 134) considered the taxon as a junior synonym of *Melanopsis fossilis* (Gmelin, 1791).

Melanopsis chehirensis var. *producta* Pallary, 1939

Original source. Pallary 1939: 95, pl. 6, figs 55, 75.

Type locality. “Yeni Chehir” [Yenişehir], Turkey.

† *Melania (Balanocochlis) propatula* Sacco, 1886

Original source. Sacco 1886: 451, pl. 1, fig. 10.

Type horizon. Late Burdigalian, early Miocene.

Type locality. “Collina di Torino” [Torino hills], Italy.

Remarks. Considered to belong in the genus *Amphimelania* by Wenz (1929: 2876).

Melanopsis prophetarum Bourguignat in Locard, 1883

Original source. Locard 1883a: 265, pl. 23, figs 52–55.

Type locality. “Dans le eaux de la fontaine de l’Élysée, à Jéricho; [...] des environs de Beyrouth; [...] dans les eaux du lac d’Antioche” [from the spring of Elysium (?) at Jericho (Palestine); from the surroundings of Beirut (Lebanon); from the waters of Lake Anuk (also as Amik) (Turkey)].

† *Melanopsis (Martinia) martiniana* var. *propinqua* Handmann, 1887

Original source. Handmann 1887: 25, pl. 4, figs 3–4.

Type horizon. Pannonian, zone B–D, late Miocene.

Type locality. “Leobersdorf”, Austria.

Remarks. Wenz (1929: 2719) considered this taxon as a junior synonym of *Melanopsis fossilis* (Gmelin, 1791).

† *Melanopsis proteus* Tournouër, 1875

Original source. Tournouër 1875: 77.

Type horizon. Tafi Formation, early Pleistocene.

Type locality. “Prope vicum Antimaki” [near the village Antimácheia, Kos Island], Greece.

Remarks. The species epithet is a noun in apposition (named after the Greek God Proteus) and needs not to agree in gender with the generic name (Art. 31.2.1).

† *Melanopsis protopygmaea* Halaváts, 1914

Original source. Halaváts 1914: 422, six unnumbered textfigures.

Type horizon. Sarmatian (sensu stricto), middle Miocene.

Type locality. “Graben unterhalb der Gemeinde-Baumschule östlich von Oltszakadát” [trench below communal tree nursery, east of Săcădate], Romania.

† *Fagotia prysjzhnjuki* Gozhik, 2002

Original source. Gozhik 2002: 54, pl. 3, figs 15, 17.

Type horizon. Middle Pontian (Dacian Basin), late Miocene.

Type locality. “Виноградівка” [Vynohradivka, Odes’ka Oblast’], Ukraine.

Types. Institute of Geological Sciences of the National Academy of Sciences of Ukraine, coll. no. 3186.

† *Melanopsis pseudoaustriaca* Sauerzopf, 1952

Original source. Sauerzopf 1952: 13, pl. 2, fig. 5.

Type horizon. Pannonian, zone D, late Miocene.

Type locality. “Burgau”, Austria.

Remarks. The name “*pseudoaustriaca*” as mentioned in Lueger (1980: 117) is an incorrect subsequent spelling.

† *Melanopsis pseudobesa* Bandel, 2000

Original source. Bandel 2000: 155, figs 33–36.

Type horizon. Late Pannonian, late Miocene.

Type locality. “Papkesi [Papkeszi] near the eastern shore of Lake Balaton”, Hungary.

Types. Geological-Palaeontological Institute and Museum University of Hamburg, coll. no. 4266.

† *Melanopsis pseudocostata* Oppenheim, 1890

Original source. Oppenheim 1890b: 591.

Type horizon. Cernikian, Pliocene.

Type locality. “Cigelnik; [...] Graben zwischen Čapla und der Podwiner Kirche; [...] Thal hinter der Podwiner Kirche; [...] Sibin; [...] Repušnica” (Neumayr and Paul 1875: 41) [Ciglenik; trench between Čapljia and the church of Podvinje; valley behind the church of Podvinje; Sibinj; Repušnica], Croatia.

Remarks. Different concepts of this species were applied in the literature and created considerable confusion. Oppenheim introduced the name clearly for the misidentified record of *M. costata* sensu Neumayr in Neumayr & Paul, 1875 (p. 41), non Olivier, 1804 from Slavonia, despite referring about Greek material. Later, however, he considered the name as rectification also for *M. costata* sensu Fuchs, 1877 from Megara and sensu Tournouër, 1876 from Kos Island (Oppenheim 1891: 465). This fact remained unnoticed by Wenz (1929: 2808), who treated *M. pseudocostata* as replacement name of the latter two records (Fuchs, Tournouër) but not of Neumayr and Paul’s species.

In addition, there are three other new names proposed for misidentified *M. costata* from the Pliocene of Slavonia, i.e., *M. croatica* Brusina, 1884 (for *M. costata* sensu Neumayr, 1869), *M. cosmanni* Pallary, 1916 [= *cossmanni*, just. emend.] (for *M. costata* sensu Cossmann, 1909) and *M. permutabilis* Pallary, 1920 (for *M. costata* sensu Brusina, 1874). Note that Wenz (1929) erroneously considered *M. cosmanni* a replacement name of Neumayr’s (1869) record.

Wenz obviously considered all erroneous “*M. costata*” records from Slavonia to represent the same species and synonymized them with *M. cosmanni*, but was unaware that *M. croatica* Brusina, 1884 is the first available name for them. Although the four names (*croatica*, *pseudocostata*, *cosmanni*, *permutabilis*) likely refer to the very same species they are based on different specimens and thus no objective synonyms.

For the record by Tournouër (1876) from Kos Island, Pallary (1929: 119) introduced *M. sculptilis* as replacement name. The misidentified *Melanopsis costata* sensu Fuchs, 1877 from Megara, however, is still nameless.

The name “*pseudodecorata*” mentioned by Cossmann (1909: 178) is an incorrect subsequent spelling.

† *Melanopsis praemorsa* var. *pseudofallax* Sacco, 1895

Original source. Sacco 1895: 9, pl. 1, fig. 14.

Type horizon. Early Messinian, late Miocene.

Type locality. “S. Marzano Oliveto”, Italy.

Remarks. Wenz (1929: 2733) considered the taxon as a junior synonym of *Melanopsis fusulatina* Sacco, 1895.

Melanopsis pseudoferussaci Pallary, 1899

Original source. Pallary 1899: 139, pl. 9, figs 11–12.

Type locality. “Environs de Tétouan” [surroundings of Tétouan], Morocco.

† *Melanopsis fossilis pseudoimpressa* Papp, 1953

Original source. Papp 1953b: 135, pl. 11, figs 5–8.

Type horizon. Pannonian, zone E, late Miocene.

Type locality. “Vösendorf (Sandriff)” [Vösendorf (sandbar)], Austria.

Remarks. Appeared first as a nomen nudum in Papp (1951: 115, 130 [as species], 147–148, 150 [as subspecies]).

† *Melanopsis impressa pseudonarzolina* Papp, 1953

Original source. Papp 1953b: 132, pl. 9, figs 14–18.

Type horizon. Pannonian, zone B, late Miocene.

Type locality. “Leobersdorf Sandgrube” [Leobersdorf sand pit], Austria.

Remarks. Appeared first as nomen nudum in Papp (1951: 107, 123, 124, 127).

† *Melanopsis pseudopigmea* Magrograssi, 1928

Original source. Magrograssi 1928: 260, pl. 6, fig. 21.

Type horizon. Plio-Pleistocene.

Type locality. “Coo: vicino a Cardamena” [Kos island: close to Kardámaina], Greece.

† *Melanopsis pseudopraerosa* Sacco, 1889

Original source. Sacco 1889: 67, pl. 2, fig. 14.

Type horizon. Late Burdigalian, early Miocene.

Type locality. “Colli torinesi” [Torino hills], Italy.

† *Melanopsis pseudopygmaea* Jekelius, 1944

Original source. Jekelius 1944: 131, pl. 49, figs 11–17.

Type horizon. Early Pannonian, late Miocene.

Type locality. “Turislav-Tal bei Soceni” [Turislav valley near Soceni], Romania.

† *Melanopsis pseudoscalaria* Sandberger, 1886

Original source. Sandberger 1886: 119.

Type horizon. *Oncophora* Beds, middle Burdigalian, early Miocene.

Type locality. Rather unprecisely given as “Kirchberger Schichten Mährens” [Kirchberg Formation in Moravia] by Sandberger (1886). Later specified as “Oslawan” [Oslavany] by Rzehak (1893: 174), Czech Republic.

† *Melanopsis pseudosubulata* Newton, 1891

Original source. Newton 1891: 203.

Type horizon. Bartonian?–Rupelian, late Eocene–early Oligocene.

Type locality. “Hempstead [...]. Headon Hill; Hordwell. [...] High Cliff”, United Kingdom.

Remarks. Introduced (as “*pseudo-subulata*”) for a part of Sowerby’s material for *M. fusiformis* Sowerby, 1822 (Sowerby 1821–1823: pl. 332, figs 6–7) as well as for misidentified *M. subulata* sensu Forbes, 1856, non Sowerby, 1822.

† *Melanopsis pterochila* Brusina, 1874

Original source. Brusina 1874: 30, pl. 1, figs 5–6.

Type horizon. Cernikian, Pliocene.

Type locality. “Podvinje (Čaplja) [Čaplja trench near Slavonski Brod]; Kovačevac”, Croatia.

Types. Milan et al. (1974: 96) indicated a holotype, but it is uncertain whether the specimen was the only one Brusina had at hand (holotype by monotypy, Art. 73.1.2). The specimen is stored in the Croatian Natural History Museum, Zagreb, coll. no. 3741-1381.

Remarks. The name “*pterochyla*” as mentioned in Pană (2003: 322) is an incorrect subsequent spelling.

Melanopsis costata var. *pulchella* Bourguignat, 1884 [unresolved]

Original source. Bourguignat 1884: 141.

Type locality. “Du Lac d’Homs” (Lorcard 1883: 288) [in Lake Homs], Syria.

Remarks. Introduced to replace - for whatever reason - the variety “*gracilis*” mentioned by Locard (1883a: 288), making *pulchella* its junior objective synonym. *M. gracilis* Locard, 1883 is a junior homonym of *M. gracilis* Brusina, 1874, which makes *pulchella* the next available name. However, *M. costata pulchella* is a homonym of the simultaneously introduced *M. seignetti pulchella* Bourguignat, 1884 (see Note 1). The action of a First Reviser is required to determine which of both *pulchella* is to be treated as valid.

Melanopsis seignetti var. *pulchella* Bourguignat, 1884 [unresolved]

Original source. Bourguignat 1884: 104.

Type locality. “[Dans les oasis du sud de la province d’Oran; [...] l’oasis Sidi Youcef, à l’extrême sud de la frontière du Maroc; [...] ruisseau d’eau chaude à Ouargla]” [in the oases in the south of the province Oran; oasis Sidi Youcef, at the far southern border of Morocco; hot water stream at Ouargla], Algeria.

Remarks. Homonym of the simultaneously introduced *M. costata pulchella* Bourguignat, 1884 (see Note 1). The action of a First Reviser is required to determine which of both *pulchella* is to be treated as valid.

Melanopsis edrissiana var. *pulchella* Pallary, 1920 [invalid]

Original source. Pallary 1920a: 33.

Type locality. “Dans une source entre Sidi Abdallah et Koudiat (région de Taza) et dans l’Innaouen, à Sidi Abdallah” [in a source between Douar Sidi Abdallah and Douar El Koudiat (Taza region) and in the Oued Abiod (?) in Douar Sidi Abdallah], Morocco.

Remarks. Junior homonym of *Melanopsis costata pulchella* Bourguignat, 1884 and *M. seignetti pulchella* Bourguignat, 1884 (simultaneously published; no priority fixed yet; see Note 1).

† *Lyrcaea* [sic] *pumila* Brusina, 1902

Original source. Brusina 1902: pl. 5, figs 37–38.

Type horizon. Middle Pannonian, late Miocene.

Type locality. “Markuševac”, Croatia.

Types. The illustrated syntypes are stored in the Croatian Natural History Museum, Zagreb, coll. no. 2491-137/1-2 (Milan et al. 1974: 84).

† *Melanopsis punctata* Stoliczka, 1860

Original source. Stoliczka 1860: 485, pl. 1, figs 5a–b.

Type horizon. Late Turonian–Coniacian, late Cretaceous.

Type locality. “Abtenau”, Austria.

Remarks. Kollmann (1984: 57) considered this species as a junior synonym of *Megalonoda reussi* (Hörnes, 1855).

† *Melanopsis pusilla* Handmann, 1882

Original source. Handmann 1882: 561.

Type horizon. Pannonian, zone D, late Miocene.

Type locality. “Kottingbrunn [...] Ziegelei a”, Austria.

Fagotia pusilla Bourguignat, 1884

Original source. Bourguignat 1884: 37.

Type locality. “Rives de la Save près Sissek, en Slavonie” [Sava river near Sisak in Slavonia], Croatia.

Remarks. Bourguignat denoted the authority as “Servain, 1884”, but there is no evidence that the description really derived from that author. Starobogatov et al. (1992: 60) considered the species as a junior synonym of *Fagotia* [= *Esperiana*] *acroxia* Bourguignat, 1884.

† *Melanopsis pygmaea* Hörnes, 1856

Original source. Hörnes 1851–1856: 599, pl. 49, fig. 13.

Type horizon. Early–middle Pannonian, late Miocene.

Type locality. “Brunn, Gumpoldskirchen, Guntramsdorf, Inzersdorf, Arsenal in Wien, Kroisbach bei Oedenburg” [Brunn am Gebirge, Gumpoldskirchen, Guntramsdorf, Inzersdorf and Arsenal in Vienna (Austria), Fertőrákos (Hungary)].

Remarks. Hörnes attributed the authority to Partsch, apparently based on an “in schedis” determination.

***Microcolpia pyramidalis* Bourguignat, 1884**

Original source. Bourguignat 1884: 61.

Type locality. “Dans le Danube, où elle a été rencontrée çà et là depuis Buda-Pesth jusqu’à Ibraila; [...] Pregrada, près de Krapina, en Croatie” [in the Danube river, where it was found here and there between Budapest (Hungary) and Brăila (Romania); Pregrada, near Krapina (Croatia)].

Remarks. Bourguignat (1884) referred to a publication by Lang (1823: 430) [not 1833 as stated by Brot 1879], who mentioned “*Melanopsis pyramidalis* (mihi)” [nomen nudum] in a species list. Starobogatov et al. (1992: 65) considered the species as a junior synonym of *Microcolpia acicularis* (Férussac, 1823).

† *Melanopsis pyramidalis* Pallary, 1920

Original source. Pallary 1920b: 117.

Type horizon. Early Rupelian, Oligocene.

Type locality. “Grossalmerode, Oberzwehren, Mardorf bei Wabern, Frielendorf, Kirchhain, Dannerod, Ofleiden, Mardorf an der Ohm” (Ludwig 1865: 71), Germany.

Remarks. Introduced for *M. praerosa* [= *M. praemorsa*] sensu Ludwig, 1865, non Linnaeus, 1758. It was regarded a junior synonym by Wenz (1929: 2740) because Sandberger (1873: 315) had already introduced *M. hassiaca* as replacement name for misidentified *M. praemorsa* from that region and time period. Since Sandberger did not explicitly refer to Ludwig’s material, *pyramidalis* and *hassiaca* are not objective synonyms.

† “*Melanopsis pyrguloeformis*” mentioned in Hermite (1879: 184) [unavailable]

Horizon. Early Eocene.

Locality. “De Binisalem et de Selva” [from Binissalem and Selva, Mallorca], Spain.

Remarks. Nomen nudum. Given as “*Melania pyrguloeformis*” on p. 328. The name was probably intended as “*pyrgulaeformis*”, but the ligature was mixed up during typesetting. Hermite listed the name in a section called “Espèces nouvelles citées et non décrites” [= “new species identified and not described”], where he listed 18 new names that he intended to describe in the second volume of his “Études géologiques

sur les îles Baléares”. That part, however, has never been published, probably because Hermite died in 1880.

† *Melanopsis (Homalia) pyrula* Handmann, 1887

Original source. Handmann 1887: 16, pl. 1, figs 16–24.

Type horizon. Pannonian, zone B–D, late Miocene.

Type locality. “Leobersdorf”, Austria.

Remarks. Wenz (1929: 2758) considered the taxon as a junior synonym of *Melanopsis inermis* Handmann, 1882.

† *Melanopsis impressa* var. *pyrulaeformis* Pavlović, 1927

Original source. Pavlović 1927: 79, pl. 10, figs 1–4.

Type horizon. Middle Pannonian, late Miocene.

Type locality. “Из Карагача” [from Karagača near Vrčin], Serbia.

Types. The illustrated syntype is stored in the Natural History Museum, Belgrade, coll. no. 222 (Milošević 1962: 22).

† *Melanopsis pyrum* Neumayr in Neumayr & Paul, 1875

Original source. Neumayr and Paul 1875: 48, pl. 8, fig. 33.

Type horizon. Cernikian, Pliocene.

Type locality. “Čaplthal bei Podwin” [Čaplja trench near Slavonski Brod], Croatia.

Remarks. The species epithet is a noun in apposition (*pyrum* = Latin “pear”) and needs not to agree in gender with the generic name (Art. 31.2.1).

† “*Melanopsis tothi* var. *quadrifilosa*” mentioned in Brusina (1903: 114) [unavailable]

Horizon. Late Pleistocene–early Holocene.

Locality. “Bischofsbad” [Püspökfürdő, Băile 1 Mai, Lake Pețea], Romania.

Remarks. Nomen nudum (Brusina apparently considered the term self-explanatory). Neubauer et al. (2014d: 125) considered this taxon as a junior synonym of *Microcolpia parreyssii sikorai* (Brusina, 1903).

† *Melanopsis raphidia* Pallary, 1920 [invalid]

Original source. Pallary 1920b: 110.

Type horizon. Plio-Pleistocene.

Type locality. “Du puits Karoubi” (Pallary 1901a: 178) [from the well Karoubi], Algeria.

Remarks. Introduced as replacement name for *Melanopsis acuminata* Pallary, 1901, non “Sandberger” [act. Gümbel, 1861], for which Wenz (1919a: 73) had already introduced the replacement name *M. pallaryi*. Thus, *M. raphidia* is a junior objective synonym of *M. pallaryi*.

† *Melanopsis (Macrospira) rapiformis* Sandberger, 1871

Original source. Sandberger 1870–1875: pl. 13, figs 3–3b.

Type horizon. Bartonian, Eocene.

Type locality. “Castres (Tarn)”, France.

Remarks. Plate 13 of Sandberger’s monograph appeared in 1871, whereas the description on p. 222 was issued in 1872 (Woodward 1906). Erroneously given as “*Melanopsis proboscidea* Desh.” in plate captions.

† *Microcolpia rara* Gozhik in Gozhik & Datsenko, 2007

Original source. Gozhik and Datsenko 2007: 95, pl. 89, fig. 6.

Type horizon. Late Pleistocene.

Type locality. “Аллювия VII террасы р. Днестр у с. Рогу” [Alluvial terrace VII of the Dniestr river near Rogi], Ukraine.

Types. Institute of Geological Sciences of the National Academy of Sciences of Ukraine, coll. no. 5243.

† *Melanoptychia rarinodosa* Brusina, 1892

Original source. Brusina 1892: 145.

Type horizon. Middle Pannonian, late Miocene.

Type locality. “Markuševac”, Croatia.

Types. The illustrated syntypes are stored in the Croatian Natural History Museum, Zagreb, coll. no. 2541-187/1-2 (Milan et al. 1974: 100).

† *Melanopsis rarispina* Lörenthey, 1902

Original source. Lörenthey 1902: 215, pl. 17, figs 18–30.

Type horizon. Transdanubian, Pannonian, late Miocene.

Type locality. “Budapest-Köbánya und Tinnye”, Hungary.

Remarks. Papp (1953b) gave *M. bouei spinosa* Handmann, 1882 and *M. bouei ventricosa* Handmann, 1882 in synonymy of *M. rarispina*, although both were published earlier. The name *ventricosa* Handmann is a junior homonym of *M. ventricosa* Neumayr, 1880 and thus invalid, but *spinosa* is available and has priority over *rarispina*.

***Melanopsis sevillensis* var. *rectituda* Pallary, 1924**

Original source. Pallary 1924: 252, pl. 25, fig. 24.

Type locality. “De la petite rivière de Guadaira qui se jette dans le Guadalquivir” [in the little river Guadaira which flows into the Guadalquivir], Spain.

† *Melanopsis recurrens* Brusina, 1874

Original source. Brusina 1874: 42.

Type horizon. Cernikian, Pliocene.

Type locality. “Bečić; Podvinje (Čaplja) [Čaplja trench near Slavonski Brod]; Sibinj; Nova Gradiška; Kovačevac; Moslavina; Farkašić”, Croatia.

† *Melanopsis gorceixi reducta* Willmann, 1981

Original source. Willmann 1981: 178.

Type horizon. Upper Elia Formation, early Pleistocene.

Type locality. “Oberes Vokasia-Tal südöstlich von Kos-Ort/Kos (Profil K 1)” [upper Vokasia valley, section K 1 (near Paradeísi)], Greece.

Types. Geological-Paleontological Institute, University of Kiel, Germany; no number indicated.

† ?*Melanopsis reineri* Penecke, 1885

Original source. Penecke 1885: 363, pl. 4, figs 8–9.

Type horizon. Paleocene.

Type locality. “Im südlichen Muldenflügel des Sonnberges (p. 334–335) [on the southern mountainside of the Sonnberg (near Guttaring, Carinthia)], Austria.

† *Melanopsis requenensis* Pallary, 1925

Original source. Pallary 1925: 257.

Type horizon. Late Miocene.

Type locality. “Meseta de Requena y Ayora (Valencia)” (Royo Gomez 1922: 110) [Plateau of Requena and Ayora], Spain.

Remarks. Introduced for *M. costata* sensu Royo Gomez, 1922, non Olivier, 1804.

***Melanopsis retoutiana* Gassies, 1861**

Original source. Gassies 1861: 293, pl. 6, fig. 9.

Type locality. “L’intérieur de la Nouvelle-Calédonie” [the interior of New Caledonia].

Remarks. Brot (1879: 454) considered the species as a junior synonym of *M. carinata* Gassies, 1861, which is a junior homonym of *Melanopsis carinata* Sowerby, 1826 (see *M. ducosi* Pallary, 1916).

† *Melanopsis retusa* Brusina, 1904

Original source. Brusina 1904: 495, pl. 1, fig. 3.

Type horizon. Langhian, middle Miocene.

Type locality. “Varcar-Vakufa” [Mrkonjić Grad], Bosnia and Herzegovina.

† *Purpuroidea reussi* Hörnes, 1855

Original source. Hörnes 1855: 177, pl. 2, figs 1a–b.

Type horizon. Late Turonian, late Cretaceous.

Type locality. “Aus der Gams, nordwestlich von Hieflau in Steiermark” [from Gams, northwest of Hieflau in Styria], Austria.

Types. Kollmann (1984: 57) designated a lectotype, which is stored in the collection of the Geological Survey Austria, Vienna (coll. no. 1856/3/4).

Remarks. Type species of the melanopsid genus *Megalonoda* Kollmann, 1984.

† *Melanopsis revelata* Pallary, 1920

Original source. Pallary 1920b: 110.

Type horizon. Pliocene.

Type locality. “Megara” (Fuchs 1877: 11), Greece.

Remarks. Replacement name for *M. incerta* Fuchs, 1877, non Férussac, 1822.

† *Melanopsis rhodanica* Locard, 1883

Original source. Locard 1883b: 44, pl. 3, fig. 6.

Type horizon. Mammal zone MN 10–12, late Miocene.

Type locality. “Boulées” [brook Boulées near Miribel], France.

Remarks. Appeared first as a nomen nudum in Falsan and Locard (1879).

***Melanopsis ricardi* Pallary, 1918**

Original source. Pallary 1918: 150.

Type locality. “Fès, dans les séguías. Ras el Mâ, à 16 kilom. de Fès” [Fes, in irrigation channels; Ras el Ma, 16 km from Fes], Morocco.

† *Melania ricinus* Neumayr in Neumayr & Paul, 1875

Original source. Neumayr and Paul 1875: 36, pl. 7, fig. 34.

Type horizon. Cernikian, Pliocene.

Type locality. “Cigelnik [Ciglenik]; [...] Novska”, Croatia.

Remarks. Neumayr mentioned uncertainties regarding the locality Ciglenik because he had lost the original collection label. The species epithet is a noun in apposition and needs not to agree in gender with the generic name (Art. 31.2.1). Considered to belong in the genus *Amphimelania* by Brusina (1897: 6), Cossmann (1909: 126) and Wenz (1929: 2876).

***Melanopsis rifi* Ahuir Galindo, 2014**

Original source. Ahuir Galindo 2014a: 14, unnumbered figure.

Type locality. “Spring from the Southeastern Rich region”, Morocco.

Types. Museo Malacologico di Cupra Marittima, Italy; no number indicated.

† *Melanopsis ripajensis* [sic] Pallary, 1916 [invalid]

Original source. Pallary 1916: 86.

Type horizon. Pannonian, zone D–E, late Miocene.

Type locality. “Ripanj” (Brusina 1902: captions of pl. 5), Serbia.

Remarks. Introduced by Pallary (1916) for the alleged junior homonym “*M. pavlovici* Brusina, 1902”, non Bourguignat, 1880. However, Brusina (1902) had named his species “*pavlovici*”, which makes *M. ripajensis* its junior objective synonym. Note that Art. 58.4. (regarding the use of u or v for the same Latin letter) does not apply here, because the names are not based on Latin words but on the names of two different persons.

† *Melanopsis haueri ripanjensis* Neubauer, Harzhauser, Kroh, Georgopoulou & Mandic, 2014

Original source. Neubauer et al. 2014c: 16.

Type horizon. Pannonian, zone D–E, late Miocene.

Type locality. “Ripanj” (Brusina 1902: captions of pl. 6), Serbia.

Types. Syntypes are stored in the Croatian Natural History Museum, Zagreb, coll. no. 2530-176/1-2 (Milan et al. 1974: 86).

Remarks. Replacement name for *M. haueri serbica* Brusina, 1902, non Brusina, 1893 (see Note 1).

Melanopsis robusta Gassies, 1870

Original source. Gassies 1870: 147.

Type locality. “Insula Ouen” [Île Ouen], New Caledonia.

Microcolpia rochebruniana Bourguignat, 1884

Original source. Bourguignat 1884: 57.

Type locality. “Lac Sabandja” [Lake Sapanca], Turkey.

Remarks. Starobogatov et al. (1992: 68) considered the species as a junior synonym of *Microcolpia coutagniana* (Pfeiffer, 1828).

† *Melanopsis subulata* var. *romejacensis* Fontannes, 1884

Original source. Fontannes 1884: 29, pl. 4, figs 3–4.

Type horizon. Early Rupelian, Oligocene.

Type locality. “Roméjac, près de Barjac (Gard)”, France.

Melanopsis roseni Izzatullaev & Starobogatov, 1984

Original source. Izzatullaev and Starobogatov 1984: 1479, fig. 1 (11).

Type locality. “Закаспийская область” [Transcaspiian Region], Russia.

Types. Zoological Institute of Russian Academy of Sciences, St.-Petersburg; no number indicated.

† *Fagotia roseni* Starobogatov in Starobogatov et al., 1992

Original source. Starobogatov et al. 1992: 62, figs 2 (3), 3 (4–5).

Type horizon. Quaternary.

Type locality. “Кулеви (быв. Регым-Кале)” [Q’ulevi (also read as Kulevi), former Redut-Kale], Georgia.

Types. Zoological Institute of Russian Academy of Sciences, St.-Petersburg; no number indicated.

Melanopsis rossiteri Gassies, 1880

Original source. Gassies 1880: 85.

Type locality. “Prope Kanala [...]; insula Ouen” (Gassies 1870: 148) [near Canala; Île Ouen], New Caledonia.

Remarks. Replacement name for *M. fusiformis* Gassies, 1870, non Sowerby, 1822.

Fagotia rosmaessleri Bourguignat, 1884

Original source. Bourguignat 1884: 34.

Type locality. “Ruisseau de Pregrada, près Krapina-Toeplilz, en Croatie” [Pregrada river near Krapinske toplice], Croatia.

Remarks. Note that Bourguignat denoted the authority as “Bourguignat, 1880”. Starobogatov et al. (1992: 60) considered the species as a junior synonym of *Fagotia* [= *Esperiana*] *esperi* (Férussac, 1823).

Melanopsis rosmaessleri Bourguignat, 1884

Original source. Bourguignat 1884: 158.

Type locality. “Spanien” [Spain, no locality indicated by Rossmässler 1839: 42].

Remarks. Introduced for *M. cariosa* sensu Rossmässler, 1839, non Linnaeus, 1767.

“*Melanopsis rothii* Ziegl.” mentioned in Brot (1874–1879: 419) [unavailable]

Locality. Not indicated.

Remarks. Nomen nudum, apparently based on an unpublished manuscript name from Ziegler and listed in synonymy of *Melanopsis buccinoidea* (Olivier, 1801) by Brot (1879).

† *Melanopsis hybostoma* var. *rotundata* Magrograssi, 1928

Original source. Magrograssi 1928: 260, pl. 6, fig. 20.

Type horizon. Plio-Pleistocene.

Type locality. “Coo: V. Iracli, V. S. Giorgio” [Kos island: Irakli valley, Ágios Geórgios valley (?)], Greece.

† *Melanopsis rudis* Brusina, 1902

Original source. Brusina 1902: pl. 5, figs 51–54.

Type horizon. Early–middle Pannonian, late Miocene.

Type locality. “Jazvine” [Jazvina], Croatia.

Types. The syntypes are stored in the Croatian Natural History Museum, Zagreb; no number indicated (Milan et al. 1974: 96).

Remarks. Wenz (1929: 2824) considered the taxon as a junior synonym of *M. senatoria* Handmann, 1887.

† *Melanopsis rugosa* Matheron, 1842

Original source. Matheron 1842: 293, pl. 37, fig. 11.

Type horizon. Early Campanian, Cretaceous.

Type locality. “Les Martigues”, France.

Remarks. D’Orbigny (1850) considered this species a *Melania* and a junior secondary homonym of *Melania rugosa* Lea, 1842 (which in turn is a replacement name for *M. corrugata* Lea, 1841, non Lamarck, 1804). However, the first one was published earlier (August 1842 vs. November/December 1842; for details on publication dates of Matheron’s work see note at the beginning of the Reference section).

† *Melanopsis (Martinia) martiniana* var. *rugosa* Handmann, 1887 [invalid]

Original source. Handmann 1887: 26, pl. 5, figs 5–7.

Type horizon. Pannonian, zone B–D, late Miocene.

Type locality. “Leobersdorf”, Austria.

Types. It is uncertain whether the single specimen from Wittmannsdorf near Leobersdorf stored in the Geological Survey Austria, Vienna, is really the only specimen left of the original type series (i.e., holotype by monotypy), as was considered by Fischer (1996b: 23). The statement by Fischer (1996b) did not suffice for a valid lectotype designation.

Remarks. Junior homonym of *Melanopsis rugosa* Matheron, 1842. Neubauer et al. (2014c: 17) introduced *M. wolfgangfischeri* as replacement name.

† *Melanopsis lanzaeana* f. *rugosa* Brusina, 1897 [invalid]

Original source. Brusina 1897: 12.

Type horizon. Langhian, middle Miocene.

Type locality. “Ribarić”, Croatia.

Types. Milan et al. (1974: 94) indicated a holotype, but it is uncertain whether the specimen is part of the original type series and whether it was the only one Brusina had at hand (holotype by monotypy, Art. 73.1.2). The specimen is stored in the Croatian Natural History Museum, Zagreb, coll. no. 2983-629.

Remarks. Junior homonym of *Melanopsis rugosa* Matheron, 1842. Neubauer et al. (2011: 205) considered the taxon as a junior synonym of *M. lanzaeana* Brusina, 1874.

† *Melanopsis doboi rugosa* Schréter, 1975 [invalid]

Original source. Schréter 1975: 11, pl. 3, figs 15–18.

Type horizon. Riss/Würm end to early Würm Ice Age, Pleistocene.

Type locality. “Eger, az egri vár Zárkándy bástyájának vasúti átmetszése” [Eger, section at the railway at the Zarkandy bastion of the fortress Eger], Hungary.

Types. Magyar Állami Földtani Intézet (Hungarian Geological Museum), Budapest; no number indicated.

Remarks. Junior homonym of *Melanopsis rugosa* Matheron, 1842. Neubauer et al. (2016a) attributed the species *M. doboi* to the genus *Microcolpia*.

† *Melanopsis matheroni* var. *rugosocarinata* Fontannes, 1880

Original source. Fontannes 1879–1882: 176.

Type horizon. Miocene or Pliocene.

Type locality. Rhône Basin? (no exact locality given), France.

Remarks. Originally written as “*rugoso-carinata*”. Based on material from the Rhône Basin, not Italy as claimed by Wenz (1929); probably the taxon was mixed up by Italian authors cited by Wenz (1929).

† *Melanopsis tothi* var. *rugosula* Brusina, 1903 [invalid]

Original source. Brusina 1903: 114.

Type horizon. Late Pleistocene–early Holocene.

Type locality. “Bischofsbad” [Püspökfürdő, Băile 1 Mai, Lake Peșea], Romania.

Remarks. Junior objective synonym of *M. tothi*: Brusina (1903) indicated it as the typical form of the species. Neubauer et al. (2014d: 125) considered this taxon as a junior synonym of *Microcolpia parreyssii sikorai* (Brusina, 1903).

“*Melanopsis ruinarum*” mentioned in Brot (1874–1879: 419–420, pl. 45, fig. 4) [unavailable]

Locality. “Balbeck” [Baalbek], Lebanon.

Remarks. Introduced in synonymy of *M. buccinoidea* by Brot (1879), referring to an “in schedis” name by Tarnier (see Note 2).

† ***Melanopsis rumana* Tournouër, 1880**

Original source. Tournouër 1880: 97.

Type horizon. Early Romanian, Pliocene.

Type locality. “[In loco Bukovatzu dicto, prope urbem Rumaniae Craiova]” [Bucovăț, near Craiova], Romania.

“*Melania sabljari* Kucik” mentioned in Brusina (1867: 86) [unavailable]

Locality. Not indicated.

Remarks. Nomen nudum, based on an “in schedis” name in the collection of Kucik (also read as “Kutschig”).

† ***Melanopsis sabolici* Brusina, 1902**

Original source. Brusina 1902: pl. 6, figs 6–7.

Type horizon. Cernikian, Pliocene.

Type locality. “Kovačevac” [east of Nova Gradiška], Croatia.

Types. The illustrated syntypes are stored in the Croatian Natural History Museum, Zagreb, coll. no. 2505-151/1-2 (Milan et al. 1974: 96).

***Melanopsis liocephala* var. *sabulosa* Pallary, 1936**

Original source. Pallary 1936: 57.

Type locality. “D’Agourai” [Agourai, south of Meknes], Morocco.

***Melanopsis maroccana saharica* Bourguignat, 1864**

Original source. Bourguignat 1864: 260.

Type locality. Rather unspecifically indicated in text as “Cours d’eau du Sahara” [rivers of the Sahara], but more precisely as “D’Aïn-Sidi-Taifour; Ouargla; fontaine

d'Oumach près de Biskra" [Sidi Taifour; Ouargla; Oumache spring near Biskra, Algeria] in the plate captions.

***Fagotia saintsimoniana* Bourguignat, 1884**

Original source. Bourguignat 1884: 47.

Type locality. "Dans la Save, au-dessous d'Agram, en Croatie" [in the Sava river below Zagreb], Croatia.

Remarks. Originally written as "*Saint-Simoniana*". Note that Bourguignat denoted the authority as "Bourguignat, 1879". Starobogatov et al. (1992: 60) considered the species as a junior synonym of *Fagotia* [= *Esperiana*] *acroxia* Bourguignat, 1884.

***Melanopsis salomonis* Bourguignat, 1884**

Original source. Bourguignat 1884: 95.

Type locality. "Fossés d'eau stagnante au camp des Pins, et çà et là dans le Liban (Syrie)" [ditches of stagnant water in the "camp des Pins" (not found, said to be near Beirut), and here and there in Lebanon].

Remarks. Appeared first as a nomen nudum in Locard (1883a: 201). Note that Bourguignat denoted the authority as "Bourguignat, 1880".

***Melanopsis sancta* Bourguignat, 1884**

Original source. Bourguignat 1884: 129.

Type locality. "Eliasbrunnen bei Jericho" (Kobelt 1880: 17) [Naḥal Elisha' (?) near Jericho], Palestine.

Remarks. Introduced for *M. costata* sensu Kobelt, 1880 (p. 17, pl. 188, fig. 1901), non Olivier, 1804. Bourguignat denoted authority as "Letourneux, 1882", but there is no evidence that the description really derived from that author. Heller et al. (2005: 248) considered the species as a junior synonym of *Melanopsis saulcyi* Bourguignat, 1853.

† ***Melanopsis sandbergeri* Neumayr, 1869**

Original source. Neumayr 1869: 372, pl. 13, fig. 5.

Type horizon. Cernikian, Pliocene.

Type locality. "Repušnica", Croatia.

***Fagotia (Fagotia) sangarica* Schütt & Bilgin, 1974**

Original source. Schütt and Bilgin 1974: 60, figs 1–4.

Type locality. “Sakarya başı, main spring of Sakarya river near village Çifteler, 60 km SE Eskişehir, 160 km WSW of Ankara”, Turkey.

Types. Senckenberg Forschungsinstitut und Naturmuseum Frankfurt, coll. no. SMF 232008.

† ?*Melanopsis sanmigueli* Royo Gómez, 1929

Original source. Royo Gómez 1929: 242, pl. 17, fig. 5.

Type horizon. Lutetian, Eocene.

Type locality. “De Santo Domingo de Silos (Burgos)”, Spain.

Remarks. Originally written as “*san-migueli*”. Probably not a Melanopsidae.

***Melanopsis saulcyi* Bourguignat, 1853**

Original source. Bourguignat 1853: 66.

Type locality. “Artouze, en Syrie” [Artouz (or ‘Arṭūz), 15 km W of Damascus, Rif Dimashq], Syria.

† ?*Melanopsis scabrida* Youlue, 1978

Original source. Youlue 1978: 155, pl. 25, figs 5–6.

Type horizon. Shahejie Formation (second Member), Eocene.

Type locality. “山东垦利” [Kenli, Shandong Province], China.

Remarks. Originally the gender was indicated as masculine (“*scabridus*”), but *Melanopsis* is feminine, which is why the name must be “*scabrida*”. Probably not a Melanopsidae (perhaps a Hydrobiidae or Pomatiopsidae?).

† *Melanopsis (Canthidomus) proteus* var. *scalarata* Magrograssi, 1928

Original source. Magrograssi 1928: 260, pl. 6, fig. 18.

Type horizon. Plio-Pleistocene.

Type locality. “Coo: V. Iracli e varie località della zona centrale fossilifera” [Kos island: Irakli valley and different locations around the central fossiliferous zone], Greece.

† *Melanopsis affinis* var. *scalariformis* Handmann, 1882

Original source. Handmann 1882: 558.

Type horizon. Pannonian, zone D, late Miocene.

Type locality. “Kottingbrunn [...] Ziegelei a”, Austria.

Remarks. Not included in the Fossilium Catalogus by Wenz (1929).

† *Melanopsis scalariformis* Papp, 1953 [invalid]

Original source. Papp 1953b: 153, pl. 3, figs 27–28.

Type horizon. Pannonian, zone C, late Miocene.

Type locality. “Leobersdorf Sandgrube” [Leobersdorf sand pit], Austria.

Remarks. Appeared first as nomen nudum in Papp (1951: 109, 147). Junior homonym of *Melanopsis affinis scalariformis* Handmann, 1882 (see Note 1).

Melanopsis scalaris Gassies, 1856

Original source. Gassies 1856: 12, figs 7–8.

Type locality. “L’Aïn-Fekan, source d’eau chaude, située entre Mascara et Saïda [...] l’Oued-M’llouya, frontière du Maroc” [‘Aïn Fekan, hot spring between Mascara and Saïda; river Moulouya, at the border to Morocco], Algeria.

† *Melanopsis scalaris* Handmann, 1882 [invalid]

Original source. Handmann 1882: 559.

Type horizon. Pannonian, zone D, late Miocene.

Type locality. “Kottingbrunn [...] Ziegelei a”, Austria.

Remarks. Junior homonym of *Melanopsis scalaris* Gassies, 1856. Pallary (1916: 83) introduced *M. limbata* as replacement name. Wenz (1929: 2741) considered the taxon as a junior synonym of *M. haueri* Handmann, 1882.

Melanella elegans var. *scalaris* Bourguignat, 1884

Original source. Bourguignat 1884: 15.

Type locality. “Au pont de la Save, près d’Agram” [at the bridge of the Sava river near Zagreb], Croatia.

***Melanopsis parreyssi* f. *scalaris* Westerlund, 1886 [invalid]**

Original source. Westerlund 1886: 123.

Type locality. “Siebenbürgen b. Deva” [Deva], Romania.

Remarks. Not available from Brot (1879: 431), who listed the name (“*Melanopsis scalaris* Parreyss in sched.”) as synonym of *Melanopsis Parreyssii* var. β , which he introduced there with a short Latin description. Further below on p. 431, Brot even associated *scalaris* with an illustration of a specimen from Deva in Romania (pl. 46, fig. 15). However, the name was explicitly referred to as synonym of the (unnamed) variety β , which is not a valid name, and thus the requirements of Art. 11.6 (and therefore Art. 12.1) are not met. Westerlund (1886) was the first to describe it and attributed the authority to Parreyss. It is a junior homonym of *M. scalaris* Gassies, 1856. Neubauer et al. (2014: 125) considered it as a junior synonym of *Microcolpia parreyssii*.

† *Melanopsis* (*Canthidomus*) *scalaritesta* Papp, 1953

Original source. Papp 1953a: 109, pl. 23, figs 9–12.

Type horizon. Gelasian, early Pleistocene.

Type locality. “Rómezi (Elis)” [Romésion near Pyrgos], Greece.

† *Melania* (*Campylostylus*) *galloprovincialis* var. *scalaroides* Oppenheim, 1892 [invalid]

Original source. Oppenheim 1892: 765, pl. 30, fig. 4.

Type horizon. Early Campanian, Cretaceous.

Type locality. “Plan de Campagne bei Septème (B.-du-Rhône)” [Plan de Campagne near Septèmes-les-Vallons], France.

Remarks. Junior homonym of *Melania scalaroides* Briart & Cornet, 1882.

***Melanopsis foleyi* var. *scalata* Pallary, 1928**

Original source. Pallary 1928a: 270.

Type locality. “Aïn Mélias, près de Figuig” [Ain Melias near Figuig], Algeria.

† *Melanopsis pterochila* var. *scansoria* Stefanescu, 1896

Original source. Stefanescu 1896: 130, pl. 11, figs 26, 28, 30–31.

Type horizon. Pliocene (Dacian?).

Type locality. “À Breasta, dans la vallée de Jiu” [near Breasta, in the valley of the river Jiu], Romania.

Remarks. The name “*scansorie*” as mentioned in Pană et al. (1981: 117) is an incorrect subsequent spelling.

***Microcolpia schileykoi* Starobogatov in Starobogatov et al., 1992**

Original source. Starobogatov et al. 1992: 67, fig. 3 (18).

Type locality. “Фёслау близ Вены” [Vöslau near Vienna], Austria.

Types. Zoological Institute of Russian Academy of Sciences, St.-Petersburg; no number indicated.

† ***Melanopsis schmidtii* Neumayr, 1880**

Original source. Neumayr 1880b: 289, pl. 1, figs 7–8.

Type horizon. Plio-Pleistocene.

Type locality. “Zwischen Pylle und Antimachia” [between Pýli and Antimácheia, Kos Island], Greece.

Remarks. Willmann (1981: 171) considered this taxon as a junior synonym of *Melanopsis sporadum* Tournouër, 1876.

† ***Melanopsis scripta* Fuchs, 1870**

Original source. Fuchs 1870b: 544, pl. 22, figs 1–2.

Type horizon. Middle Pannonian, late Miocene.

Type locality. “Kúp”, Hungary.

† ***Melanopsis sculptilis* Pallary, 1920**

Original source. Pallary 1920b: 119.

Type horizon. Plio-Pleistocene.

Type locality. “Antimaki” (Tournouër 1876: 449, 455) [Antimácheia], Greece.

Remarks. Introduced for *M. costata* sensu Tournouër, 1876. It was indirectly considered as a junior synonym by Willmann (1981: 177), who was unaware of Pallary’s replacement name but synonymized *M. costata* sensu Tournouër, 1876 with *M. heldreichi* Neumayr, 1880.

***Melanopsis seignetti* Bourguignat, 1884**

Original source. Bourguignat 1884: 104.

Type locality. “Dans les oasis du sud de la province d’Oran; [...] l’oasis Sidi Yousef, à l’extrême sud de la frontière du Maroc; [...] ruisseau d’eau chaude à Ouargla” [in the oases in the south of the province Oran; oasis Sidi Youcef, at the far southern border of Morocco; hot water stream at Ouargla], Algeria.

Remarks. Note that Bourguignat denoted the authority as “Bourguignat, 1872”.

† *Melanoptychia lyrata semicostata* Olujić, 1999

Original source. Olujić 1999: 20, 48, pl. 1, figs 2–4.

Type horizon. Langhian, middle Miocene.

Type locality. It is unclear from the original work in which of the studied localities/sections along the valleys of the Sutina, Batarelov and Vojskava rivers (4 km W of Sinj) the taxon occurred and in which not, Croatia.

Remarks. Neubauer et al. (2011: 207) considered the taxon as a junior synonym of *Melanopsis lyrata* Neumayr, 1869.

***Melanopsis semigranulosa* Deshayes, 1832**

Original source. Deshayes 1830–1832: 438.

Type locality. “Dans l’Ohio” [in the Ohio river], United States.

***Melanopsis eximia* var. *semilaevigata* Pallary, 1928 [unresolved]**

Original source. Pallary 1928a: 260.

Type locality. Not explicitly stated but probably the same as for the species (Agourai, south of Meknès, Morocco).

Remarks. Homonym of the simultaneously published name *Melanopsis seurati semilaevigata*. This case requires the action of a First Reviser.

***Melanopsis seurati* var. *semilaevigata* Pallary, 1928 [unresolved]**

Original source. Pallary 1928a: 268.

Type locality. Not explicitly stated but probably the same as for the species (“La Zousfana, à la hauteur de Figuig et à Beni Ounif” [Oued Zousfana, at the height of Figuig, and at Beni Ounif], Algeria).

Remarks. Homonym of the simultaneously published name *Melanopsis eximia semilaevigata*. This case requires the action of a First Reviser.

***Melanopsis vanrossomi* var. *semilaevigata* Pallary, 1936 [invalid]**

Original source. Pallary 1936: 59.

Type locality. “Tanalt”, Morocco.

Remarks. Junior homonym of *Melanopsis seurati semilaevigata* Pallary, 1928 and *M. eximia semilaevigata* Pallary, 1928 (simultaneously published; no priority fixed yet; see Note 1).

***Melanopsis (Mesopotamia) aroussiana* var. *semilaevigata* Pallary, 1939 [invalid]**

Original source. Pallary 1939: 101.

Type locality. Not explicitly stated but probably the same as for the species (“Ain Arouss” [‘Ayn al ‘Arūs, near Tall Abyaḍ], Syria).

Remarks. Pallary (1939) indicated an illustration (“Fig. 4”) of this taxon, which, however, does not appear as such in the plate captions. Junior homonym of *Melanopsis seurati semilaevigata* Pallary, 1928 and *M. eximia semilaevigata* Pallary, 1928 (simultaneously published; no priority fixed yet; see Note 1).

***Melanella holandri* var. *semiplicata* Brusina, 1870**

Original source. Brusina 1870: 14.

Type locality. “Dans la Glina près de Topusko” [in the Glina river near Topusko], Croatia.

† *Melanopsis semiplicata* Neumayr, 1880

Original source. Neumayr 1880b: 292, pl. 1, fig. 18.

Type horizon. Phoka to Elia Formation, Plio-Pleistocene.

Type locality. “Phuka” [Ákra Ágios Fokás], Greece.

***Melanopsis foucauldiana* var. *semiplicata* Pallary, 1928 [invalid]**

Original source. Pallary 1928a: 254, pl. 4, fig. 31.

Type locality. Not explicitly stated but probably the same as for the species (“Ougarta, à 60 kil. sud de Beni-Abbès” [Ougarta, prov. Béchar], Algeria).

Remarks. Junior homonym of *Melanopsis semiplicata* Neumayr, 1880.

***Melanopsis excoriata* var. *semisulcata* Pallary, 1920**

Original source. Pallary 1920c: 144.

Type locality. “Ait Brahim”, Morocco.

† ***Melanopsis semperi* De Stefani, 1877**

Original source. De Stefani 1877: 310.

Type horizon. Villafranchian, Plio-Pleistocene.

Type locality. “Orciano” (Pecchioli 1864: 524) [Orciano Pisano], Italy.

Remarks. Replacement name for *M. nodosa* Pecchioli, 1864, non Férussac, 1822.

† ***Melanopsis (Lyrcea) senatoria* Handmann, 1887**

Original source. Handmann 1887: 19, pl. 2, figs 10–11.

Type horizon. Pannonian, zone B–D, late Miocene.

Type locality. “Leobersdorf”, Austria.

“*Melanopsis magnifica* m. *senilis*” mentioned in Pérès (1939) [unavailable]

Locality. “Stations 117 et 117his: Oued Bou Fekrane à 13 kilomètres de Meknès sur la route d’El Hajeb” [Stations 117 and 117His: Oued Boufekrane, 13 km S of Meknes on the road to El Hajeb], Morocco.

Remarks. Introduced as infrasubspecific taxon (“mode”), which is not governed by the provisions of the Code. Moreover, the name is a nomen nudum. Pérès referred to an earlier publication of his (Pérès 1938a), claiming that he had described the taxon there already. In that work, however, he only mentioned a subfossil and a “forme modifiée” of *M. magnifica* Bourguignat, 1884.

† ***Melanopsis lanzae senilis* Olujić, 1999**

Original source. Olujić 1999: 22, 50, pl. 3, figs 30–32.

Type horizon. Langhian, middle Miocene.

Type locality. It is unclear from the original work in which of the studied localities/sections along the valleys of the Sutina, Batarelov and Vojskava rivers (4 km W of Sinj) the taxon occurred and in which not, Croatia.

Remarks. Neubauer et al. (2011: 205) considered it as a junior synonym of *M. lanzaeana* Brusina, 1874.

† *Melanopsis seninskii* Wenz, 1928

Original source. Wenz 1928a: 119.

Type horizon. Duab Beds, middle to late Kimmerian, Pliocene.

Type locality. “Моквинские пласты, р. Дуабъ” (Seninski 1905: 62) [Mokvi layers at Duab river], Georgia.

Remarks. Replacement name for *Melanopsis acuminata* Seninski, 1905, non Gümbel, 1861.

† *Melania serbica* Živković, 1893

Original source. Živković 1893: 165.

Type horizon. Early Langhian, middle Miocene.

Type locality. “Zvezdan, Crna Reka oberhalb Zaječar, Vražogrnac, Vrbiza, Sumrakovac, Šarbanovac”, Serbia.

Remarks. Considered to belong in the genus *Amphimelania* by Wenz (1929: 2878).

† *Melanopsis serbica* Brusina, 1893

Original source. Brusina 1893: 50 (only described in Italian part).

Type horizon. Pannonian, zone D–E, late Miocene.

Type locality. “Begaljica”, Serbia.

Types. Milan et al. (1974: 85) indicated a holotype, but it is uncertain whether the specimen actually derives from the original type series and whether it was the only specimen Brusina had at hand. The specimen is stored in the Croatian Natural History Museum, Zagreb, coll. no. 3013-659.

† *Melanopsis austriaca serbica* Brusina, 1902 [invalid]

Original source. Brusina 1902: pl. 6, figs 73–74.

Type horizon. Pannonian, zone D–E, late Miocene.

Type locality. “Ripanj”, Serbia.

Types. The illustrated syntypes are stored in the Croatian Natural History Museum, Zagreb, coll. no. 2530-176/1-2 (Milan et al. 1974: 86).

Remarks. Junior homonym of *Melanopsis serbica* Brusina, 1893. Neubauer et al. (2014c: 16) introduced *M. haueri ripanjensis* as replacement name, following Wenz (1929: 2741, 2743) who synonymized *M. austriaca* Handmann, 1882 with *M. haueri* Handmann, 1882 and listed *serbica* as a subspecies of *M. haueri*.

† *Melanopsis serchensis* Vidal, 1874

Original source. Vidal 1874: 236, pl. 3, figs 14–14a.

Type horizon. Maastrichtian, Cretaceous.

Type locality. “En Serchs y en Auzas (Alto Garona), pero principalmente en Isona” [in Cercs (Spain) and Auzas (Dép. Haute-Garonne, France), but mainly in Isona (Spain)].

Remarks. Considered to belong to the subgenus *Stilospirula* by Dominici and Kowalke (2014: 148).

Melanopsis magnifica var. *serira* Pallary, 1920

Original source. Pallary 1920c: 153.

Type locality. “Jardin public de bou Jeloud, oued Masmouda, avant son entrée à Fàs el bâli” [park of Bou Jeloud, river Sakiat Masmouda before entry to Fès al Bali], Morocco.

† *Melanopsis serrensis* Mayer-Eymar, 1903

Original source. Mayer-Eymar 1903: 318, pl. 13, fig. 9.

Type horizon. Sparnacian, early Ypresian, Eocene.

Type locality. “Du village de la Serre” [La Serre (?)], France.

Microcolpia servaini Bourguignat, 1884

Original source. Bourguignat 1884: 55.

Type locality. “La Save à Sissek (Slavonie) et dans la rivière de Zenica (Bosnie)” [Sava river at Sisak (Croatia) and in the river at Zenica (Bosnia and Herzegovina)].

Remarks. Appeared first as a nomen nudum in Servain (1884). Starobogatov et al. (1992: 65) considered the species as a junior synonym of *Microcolpia cornea* (Pfeiffer, 1828).

Fagotia servainiana Bourguignat, 1884

Original source. Bourguignat 1884: 44.

Type locality. “Rivières entre Plaski et Ostaria (Croatie)” [river between Plaški and Oštarije], Croatia.

Remarks. Note that Bourguignat denoted the authority as “Bourguignat, 1882”.

***Melanopsis sesteri* Bourguignat, 1884**

Original source. Bourguignat 1884: 119.

Type locality. “Petit cours d’eau à Sadjour-Sou, entre Aïn-Taïb et Alep [...]; Aïn-el-Bass, dans la plaine du Bahr-el-Houlé (Syrie)” [small brook at Sadjour-Sou between Gaziantep (Turkey) and Aleppo (Syria) [...]; Aïn el Bass, in the plains of the Hula valley (Israel)].

Remarks. Heller et al. (2005: 232) considered the species as a junior synonym of *Melanopsis buccinoidea* (Olivier, 1801).

***Melanopsis seurati* Pallary, 1920**

Original source. Pallary 1920a: 33.

Type locality. “La Zousfana, à la hauteur de Figuig et à Beni Ounif” [Oued Zousfana, at the height of Figuig, and at Beni Ounif], Algeria.

***Melanopsis sevillensis* Grateloup, 1840**

Original source. Grateloup 1840: 433, pl. 4, figs 10–11.

Type locality. “Séville; les bords de la petite rivière de Guadaira, qui se jette dans le Guadalquivir” [Sevilla; the banks of the small river Guadaira, which flows into the Guadalquivir], Spain.

Remarks. The “var. *minor* [...] de Grateloup” as discussed by Bourguignat (1884) was never declared as such by Grateloup (“var. b”). Brot (1874–1879: 441) considered the taxon as a junior synonym of *Melanopsis cariosa* (Linnaeus, 1767).

***Melanopsis sharhabili* Bandel, 2000**

Original source. Bandel 2000: 156, figs 119–123.

Type locality. “Spring and creek next to the Wadi Raiyan Plantation in the Jordan Valley near the town of Wadi Raiyan and close to the mosque of the grave of Sharhabil Ibn Hassana”, Jordan.

Types. Geological-Palaeontological Institute and Museum University of Hamburg, coll. no. 4267.

***Melanopsis sharpei* Pallary, 1922**

Original source. Pallary 1922: 211.

Type locality. “Aoullouz” [Alous], Morocco.

“*Melanopsis siamensis* Mrts.” mentioned in Paetel (1888: 404) [unavailable]

Locality. “Siam”, Thailand.

Remarks. Nomen nudum, appears only in the species list of Paetel (1888). Perhaps it is a lapsus calami of *Melampus siamensis* Martens, 1866 (Ellobiidae).

† *Melanopsis sigmocorrugata* Heller & Sivan, 2001 [unavailable]

Original source. Heller and Sivan 2001: 140, fig. 3G.

Type horizon. Pleistocene.

Type locality. “Gesher Benot Ya’aqov”, Syria.

Remarks. Since Heller and Sivan (2001) did not fix a type, the name is unavailable after Art. 16.4.

† *Melanopsis sikorai* Brusina, 1903

Original source. Brusina 1903: 111.

Type horizon. Late Pleistocene–Holocene.

Type locality. “Bischofsbad” [Püspökfürdő, Băile 1 Mai, Lake Pețea], Romania.

Remarks. Considered a subspecies of *Microcolpia parreyssii* by Neubauer et al. (2014d).

† *Melanopsis similis* Pallary, 1916

Original source. Pallary 1916: 84.

Type horizon. Pannonian, zone B–D, late Miocene.

Type locality. “Leobersdorf” (Handmann 1887), Austria.

Remarks. Replacement name for *M. plicatula* Handmann, 1887, non Brusina, 1874. Wenz (1929: 2674) considered the taxon as a junior synonym of *Melanopsis bouei* Férussac, 1823.

† “*Melanopsis sikorai* var. *siminina*” mentioned in Brusina (1903: 112) [unavailable]

Horizon. Late Pleistocene–early Holocene.

Locality. “Bischofsbad” [Püspökfürdő, Băile 1 Mai, Lake Pețea], Romania.

Remarks. Nomen nudum. Listed in synonymy of *Microcolpia parreyssii sikorai* (Brusina, 1903) by Neubauer et al. (2014d: 125).

† *Melanopsis simulata* Pallary, 1925 [invalid]

Original source. Pallary 1925: 257.

Type horizon. Cernikian, Pliocene.

Type locality. “Malino” (Brusina 1878: 348), Croatia.

Remarks. Pallary (1925) introduced *M. simulata* as replacement name for the presumed junior homonym *M. arcuata* Brusina, 1878, non Matheron, 1842. The alleged homonymy is, however, based on a reading error of *M. armata* Matheron, 1842 by Pallary. Therefore, *M. simulata* is a junior objective synonym of *M. arcuata* Brusina, 1897.

† *Melanopsis sinjana* Brusina, 1874

Original source. Brusina 1874: 32, pl. 1, figs 1–2.

Type horizon. Early Langhian, middle Miocene.

Type locality. “Sinj (Stuparuša)”, Croatia.

Types. Milan et al. (1974: 97) defined a “neotype” based on one of the specimens illustrated by Brusina (1897). It is uncertain, however, whether the specimen was part of the original type series, which would qualify it as lectotype. Nonetheless, the type fixation is insufficient with respect to Art. 75.3 of the Code. The specimen is stored in the Croatian Natural History Museum, Zagreb, coll. no. 2974-620/1.

† *Melanopsis sinzowi* Brusina, 1885

Original source. Brusina 1885: 160.

Type horizon. *Spaniodon* Beds, Karaganian, middle Miocene.

Type locality. “Лонушны” (Sinzov 1884: 5) [Lăpușna], Moldova.

Remarks. Introduced for *M. sinjana* sensu Sinzov, 1884, non Brusina, 1874.

† *Melanopsis sinzowi* Lörenthey, 1902 [invalid]

Original source. Lörenthey 1902: 213, pl. 17, figs 31–32.

Type horizon. Transdanubian, Pannonian, late Miocene.

Type locality. “Tinnye”, Hungary.

Remarks. Junior homonym of *Melanopsis sinzowi* Brusina, 1885. Wenz (1919a: 73) introduced *M. tinnyensis* as replacement name. Papp (1953b: 145) considered the taxon as a junior synonym of *Melanopsis rarispina* Lörenthey, 1902. However, Papp (1953b) gave the older names *M. bouei spinosa* Handmann, 1882 and *M. bouei ventricosa* Handmann, 1882 in synonymy of *M. rarispina*. While *ventricosa* Handmann is a junior homonym of *M. ventricosa* Neumayr, 1880 and thus invalid, the name *spinosa* is available and has priority over *rarispina*.

***Melanopsis sistanica* Izzatullaev & Starobogatov, 1984**

Original source. Izzatullaev and Starobogatov 1984: 1477, fig. 1 (15), fig. 2 (6), fig. 3 (5).

Type locality. “Восточная Персия” [Eastern Persia], Iran.

Types. Zoological Institute of Russian Academy of Sciences, St.-Petersburg; no number indicated.

Remarks. Type species of *Melanopsis* (*Sistaniana*) Izzatullaev & Starobogatov, 1984.

† *Melanopsis* (*Canthidomus*) *costatiformis skuphoi* Papp & Psarianos, 1955

Original source. Papp and Psarianos 1955: 148, pl. 23, figs 6–9.

Type horizon. Early Pleistocene.

Type locality. “Tárapsa” [Vasilákion], Greece.

Types. Museum of Palaeontology and Geology of the University of Athens; no number indicated.

† *Melanopsis* (*Melanosteira*) *skurensis* Papp, 1955

Original source. Papp 1955: 130, pl. 20, figs 27–30.

Type horizon. Akchagylian, latest Pliocene–earliest Pleistocene.

Type locality. “Skura bei Sparta” [Skoúra], Greece.

Types. Museum of Palaeontology and Geology of the University of Athens; no number indicated.

† *Melanopsis slavonica* Neumayr in Neumayr & Paul, 1875

Original source. Neumayr and Paul 1875: 45, pl. 8, fig. 25.

Type horizon. Cernikian, Pliocene.

Type locality. “Graben zwischen Podwin und der Čapla” [Čaplja trench near Slavonski Brod], Croatia.

† *Melanopsis sobrievskii* Rosen, 1914

Original source. Rosen 1914: 221, pl. 3, fig. 19.

Type horizon. Pleistocene?

Type locality. “Am Fusse des Černjajev’schen Berges bei Suchum” [at the foot of Mt. Černjajev (?) at Sokhumi], Georgia.

Remarks. On p. 221 Rosen gave the species as “*Melanopsis sobrievskii*” within the subgenus *Fagotia*. In the plate captions it appears as “*Fagotia sobrievskii*”.

† *Melanopsis soceni* Jekelius, 1944

Original source. Jekelius 1944: 73, pl. 16, figs 14–17, pl. 51, figs 1–11.

Type horizon. *Mohrensternia* Zone, early Sarmatian, middle Miocene.

Type locality. “Polițioanătal bei Soceni” [Polițioană valley near Soceni], Romania.

† *Boistelia soceni* Harzhauser, Kowalke & Mandic, 2002

Original source. Harzhauser et al. 2002: 99, pl. 7, fig. 6.

Type horizon. Pannonian, zone C–D, late Miocene.

Type locality. “St. Margarethen (Burgenland)” [more precisely, it is the Zollhaus sandpit near St. Margarethen], Austria.

Types. Geological-Paleontological Department, Natural History Museum Vienna, Austria, coll. no. 2001/0126/0049.

Remarks. Neubauer et al. (2014a: 463) considered the taxon as a junior synonym of *Melanopsis soceni* Jekelius, 1944.

† *Melanopsis sodalis* Deshayes, 1862

Original source. Deshayes 1861–1864: 470, pl. 31, figs 14–15.

Type horizon. Thanetian, Paleocene.

Type locality. “Châlons-sur-Vesles, Gueux, Jonchery, Noailles”, France.

† *Melanopsis soldaniana* De Stefani in Pantanelli, 1877

Original source. Pantanelli 1877: 5.

Type horizon. Late Messinian, late Miocene.

Type locality. “Casino, e specialmente presso le Gallozzole” [Casino, and especially at Gallozzole], Italy.

Remarks. Pantanelli (1879: 315, pl. 2, figs 4–5) re-described the species and provided illustrations.

Melanopsis sordida Gassies, 1871

Original source. Gassies 1871: pl. 6, fig. 9.

Type locality. New Caledonia [no locality indicated].

Remarks. Gassies (1871) only illustrated but not described the species. Brot (1879: 444) considered the taxon as a junior synonym of *Melanopsis frustulum* Morelet, 1857.

† *Melanopsis sostarici* Brusina, 1897

Original source. Brusina 1897: 8, pl. 6, figs 9–12.

Type horizon. Latest Burdigalian, early Miocene.

Type locality. “Dugoselo”, Croatia.

Types. Milan et al. (1974: 97) indicated a holotype, but it is uncertain whether the specimen was the only one Brusina had at hand (holotype by monotypy, Art. 73.1.2). The specimen is stored in the Croatian Natural History Museum, Zagreb, coll. no. 2999-645.

† *Melanopsis soubeirani* Porumbaru, 1881

Original source. Porumbaru 1881: 28, pl. 9, fig. 1.

Type horizon. Early–middle Romanian, Pliocene.

Type locality. “Cretzeszi, Podari” [Crețești, Podari], Romania.

Remarks. The name “*soubeiranus*” as mentioned in Macarovici (1940: 344) is an incorrect subsequent spelling.

Melanopsis souverbieana Gassies, 1870

Original source. Gassies 1870: 148.

Type locality. “In Nova Caledonia” [New Caledonia; no locality indicated].

† *Melanopsis sowerbyi* Pallary, 1920 [invalid]

Original source. Pallary 1920b: 116.

Type horizon. Headon Beds, Priabonian, Eocene.

Type locality. “Hordwell” (Sowerby 1821–1823: 36), United Kingdom.

Remarks. Introduced for one of the illustrated specimens of *M. fusiformis* Sowerby, 1846 (pl. 221, fig. 9) [sic]. The name is based on an error of Pallary (1920b), who only had the French translation of Sowerby’s “Mineral conchology” from 1845 (not 1846) at hand, where figures were organized in a different way than in the original version. Fig. 9 in the 1845-version corresponds to Fig. 7 on pl. 332 of the original publication (Sowerby 1822). Newton (1891) had introduced the new name *M. pseudosubulata* for that specimen and another specimen already (Sowerby 1821–1823: pl. 332, figs 6–7). Since Pallary mixed up the figures, he considered Newton’s taxonomic separation misleading and, in order to settle the problem, introduced *M. sowerbyi*. Consequently, *M. sowerbyi* is a junior objective synonym of *M. pseudosubulata*.

† *Melanopsis sparnacensis* Pallary, 1916

Original source. Pallary 1916: 77.

Type horizon. Sparnacian, early Ypresian, Eocene.

Type locality. “Épernay”, France.

Remarks. Based on a specimen illustrated by Férussac (1823: pl. 7, fig. 7) as “*Melanopsis buccinoidea* var. γ) antiqua; *elongata*”. Note that *M. sparnacensis* was not included in the catalogus of Wenz (1929).

† *Melanopsis (Martinia) martiniana* var. *spatiosa* Handmann, 1887

Original source. Handmann 1887: 25, pl. 4, figs 5–7.

Type horizon. Pannonian, zone B–D, late Miocene.

Type locality. “Leobersdorf”, Austria.

Remarks. Wenz (1929: 2719) considered this taxon as a junior synonym of *Melanopsis fossilis* (Gmelin, 1791).

Melanella speciosa Bourguignat, 1884

Original source. Bourguignat 1884: 28.

Type locality. “Rivière entre Ostaria et Plaski, dans la Croatie méridionale” [river between Plaški and Oštarije], Croatia.

Remarks. Note that Bourguignat denoted the authority as “Bourguignat, 1879”.

Melanopsis sphaeroidaea Bourguignat, 1884

Original source. Bourguignat 1884: 78.

Type locality. “Dans l’Oronte (Syrie)” [in the Orontes river], Syria?

Remarks. Spelt as “*sphaeroidoea*” on p. 78 but as “*sphaeroidaea*” on p. 73. From the description it is clear that the name must be “*sphaeroidaea*” (see Art. 33.2.1).

† *Melanopsis bouei* var. *spinea* Stefanescu, 1896

Original source. Stefanescu 1896: 136, pl. 11, figs 65–68.

Type horizon. Khersonian, late Sarmatian, late Miocene.

Type locality. “À Sacel, dans la vallée de Blahnitza, district de Gorjiu” [at Săcelu, in the valley of the Blahnița river, Gorj county], Romania.

Remarks. Wenz (1929: 2678) considered the taxon as a junior synonym of *Melanopsis bouei* Férussac, 1823.

† *Melanopsis spinicostata* Rolle, 1860

Original source. Rolle 1860: 32, pl. 2, figs 6–8.

Type horizon. Late Pliocene to early Pleistocene.

Type locality. “Skalis prope Schoenstein” [Pesje near Šoštanj], Slovenia.

Remarks. The name “*spinicosta*” as mentioned in Wenz (1929: 2831) is an incorrect subsequent spelling.

† *Trochus spiniger* Sowerby in Sedgwick & Murchison, 1832

Original source. Sedgwick and Murchison 1832: 418, pl. 38, fig. 15.

Type horizon. Late Cretaceous.

Type locality. “Gosau”, Austria.

Types. Holotype (?) stored in the collection of the British Museum, coll. no. G 17908 (Kollmann 1984: 58).

Remarks. Kollmann (1984: 58) classified the species in the melanopsid genus *Megalonoda*, which was there described as new.

† *Melanopsis spinigera* Seninski, 1905

Original source. Seninski 1905: 60, pl. 1, figs 19–22.

Type horizon. Duab Beds, middle to late Kimmerian, Pliocene.

Type locality. “Моквинские пласты, р. Дуабъ” [Mokvi layers at Duab river], Georgia.

† *Melanopsis bouei* var. *spinosa* Handmann, 1882

Original source. Handmann 1882: 557.

Type horizon. Pannonian, zone D, late Miocene.

Type locality. “Kottingbrunn [...] Ziegelei a”, Austria.

Remarks. Papp (1953b) listed this variety together with *M. bouei ventricosa* Handmann, 1882 in synonymy of *M. rarispina* Lörenthey, 1902, although Handmann’s taxa were published earlier. The name *ventricosa* Handmann is a junior homonym of *M. ventricosa* Neumayr, 1880 and thus invalid, but *spinosa* is available and has priority over *rarispina*.

† *Melanopsis spiralis* Handmann, 1882

Original source. Handmann 1882: 555.

Type horizon. Pannonian, zone D, late Miocene.

Type locality. “Kottingbrunn [...] Ziegelei a”, Austria.

Remarks. Wenz (1929: 2718) considered this taxon as a junior synonym of *Melanopsis fossilis* (Gmelin, 1791).

***Melanopsis spirata* Chenu, 1859**

Original source. Chenu 1859: 297, fig. 2062.

Type horizon. Not stated; unclear if recent or fossil.

Type locality. Not indicated.

Remarks. The species is based on a single illustration, without description or any kind of explanation.

† ***Melanopsis avellana* var. *spirata* Leymerie, 1881 [invalid]**

Original source. Leymerie 1881: 777, pl. O, fig. 2.

Type horizon. Maastrichtian, Cretaceous.

Type locality. “Auzas”, France.

Remarks. Junior homonym of *Melanopsis spriata* Chenu, 1859.

† ***Melanopsis spiridioni* Pallary, 1916**

Original source. Pallary 1916: 81.

Type horizon. Latest Burdigalian, early Miocene.

Type locality. “Dugoselo” (Brusina 1897: 9), Croatia.

Remarks. Introduced for *M. praemorsa* sensu Brusina, 1897, non Linnaeus, 1758. Wenz (1929: 2831–2833) listed also other misidentified *M. praemorsa* from the late Miocene deposits of Lake Pannon and the Pliocene of Greece and Croatia under that name, but it is very unlikely that all of them belong to the same species.

† ***Melanopsis sporadum* Tournouër, 1876**

Original source. Tournouër 1876: 453, pl. 4, fig. 4.

Type horizon. Phoka to Elia Formation, Plio-Pleistocene.

Type locality. “Fouka” (p. 449) [Ákra Ágios Fokás], Greece.

† ***Melanopsis aetolica* var. *stamnana* Oppenheim, 1891**

Original source. Oppenheim 1891: 469, pl. 27, figs 3–4.

Type horizon. Gelasian, early Pleistocene.

Type locality. “Stamna”, Greece.

Remarks. Not available from Oppenheim (1890b), where he gave it as “mutation”. Probably by mistake, Papp (1955, in figure captions) designated a lectotype based on a specimen from Neumayr’s (1880a) (not Oppenheim’s) original material.

† *Melanopsis (Duabiana) starobogatovi* Anistratenko, 1993

Original source. Anistratenko 1993: 70, textfig. 1.

Type horizon. Duab Beds, middle to late Kimmerian, Pliocene.

Type locality. “Окр. с. Мокви, Очамчирский р-н” [near the village Mok’vi, Ochamchirskiy rayon], Georgia.

Types. Schmalhausen Institute of Zoology of National Academy of Sciences of Ukraine, Kiev; no number indicated.

Melanopsis starostini Izzatullaev & Starobogatov, 1984

Original source. Izzatullaev and Starobogatov 1984: 1480, fig. 1 (13), fig. 3 (7).

Type locality. “Гермоб (Центральный Конемгаз)” [Germob (central Kopetdag)], Iran.

Types. Zoological Institute of Russian Academy of Sciences, St.-Petersburg; no number indicated.

† *Melanopsis staubi* Brusina, 1903

Original source. Brusina 1903: 115.

Type horizon. Late Pleistocene–Holocene.

Type locality. “Bischofsbad” [Püspöckfürdő, Băile 1 Mai, Lake Peșea], Romania.

Remarks. Neubauer et al. (2014d: 125) considered this taxon as a junior synonym of *Microcolpia parreyssii sikorai* (Brusina, 1903).

Melanopsis (Mesopotamia) infracincta var. *stena* Pallary, 1939

Original source. Pallary 1939: 104.

Type locality. “Ras el ‘Ain du Khabour” [Chabur river near Ra’s al ‘Ayn], Syria.

Fagotia stenostoma Bourguignat, 1884

Original source. Bourguignat 1884: 42.

Type locality. “La Save au-dessous d’Agram; rivières au sud de Krapina-Toeplitz, et entre Plaski et Ostaria (Croatie)” [Sava river below Zagreb; rivers south of Krapinske toplice, and between Plaški and Oštarije], Croatia.

Remarks. Note that Bourguignat denoted the authority as “Bourguignat, 1879”. Starobogatov et al. (1992: 60) considered the species as a junior synonym of *Fagotia* [= *Esperiana*] *acroxia* Bourguignat, 1884.

***Melania parvula* f. *stenostoma* Westerlund, 1886**

Original source. Westerlund 1886: 108.

Type locality. “Der untere Lauf jenes kleinen Gebirgsbaches, welcher den Abhängen des Berges Kopitnig entspringt, am hochgelegenen Kurhause von Römerbad vorbeieilend, dessen Thermenabfluss aufnimmt, und in starkem Gefälle der den Fuss des Berges umspülenden Sann zufließt” (Tschapeck 1881: 105–106) [in the lower reaches of a small mountain brook, which originates at the slopes of Mt. Kopitnik, passes the kurhaus of Rimske Toplice, takes its thermal water runoff, and flows over a steep gradient into the Savinja river], Slovenia.

Remarks. Introduced for a specimen of *Melania holandri* illustrated by Tschapeck (1881: 106, pl. 5, fig. k).

***Melanopsis stephanota* Bourguignat, 1884**

Original source. Bourguignat 1884: 120.

Type locality. Rossmässler, to whom Bourguignat referred, indicated the locality only as “Palästina” [Palestine]. As the species was explicitly introduced for Rossmässler’s material only this is the type locality.

Remarks. Introduced for *M. costata* sensu Kobelt, 1880 (figs 1899–1900), non Olivier, 1804.

***Melanopsis (Mesopotamia) stolliana* Pallary, 1939**

Original source. Pallary 1939: 102, pl. 5, figs 44–48.

Type locality. “‘Ain Arouss” [‘Ayn al ‘Arūs (near Tall Abyaḍ)], Syria.

***Microcolpia stossichiana* Bourguignat, 1884**

Original source. Bourguignat 1884: 64.

Type locality. “Carniole” [Carniola, a historical region that comprised parts of present-day Slovenia; no locality indicated].

Remarks. Starobogatov et al. (1992: 65) considered the species as a junior synonym of *Microcolpia acicularis* (Férussac, 1823).

***Melanopsis strangei* Reeve, 1860**

Original source. Reeve 1860: Section *Melanopsis*, pl. 1, figs 3a–b.

Type locality. “New Zealand” [no locality indicated].

Remarks. Suter (1893: 139) considered the taxon as a junior synonym of *Zemelanopsis trifasciata* (Gray in Dieffenbach, 1843).

† *Melanopsis strangulata* Brusina, 1902

Original source. Brusina 1902: pl. 6, figs 46–50.

Type horizon. Cernikian, Pliocene.

Type locality. “Kovačevac” [east of Nova Gradiška], Croatia.

Types. The illustrated syntypes are stored in the Croatian Natural History Museum, Zagreb, coll. no. 2521-167/1-5 (Milan et al. 1974: 97).

† *Melanopsis olivula* var. *striata* Grateloup, 1838 [invalid]

Original source. Grateloup 1838: 143, pl. 4, fig. 55.

Type horizon. Burdigalian, early Miocene.

Type locality. “Dax. [...] Mandillot”, France.

Remarks. Junior objective synonym of *M. buccinoidea subventricosa* Grateloup, 1828, which Grateloup (1838) listed in synonymy.

***Melanopsis maroccana* var. *striata* Pantanelli, 1886**

Original source. Pantanelli 1886b: 72, pl. 3, figs 26–27.

Type locality. “Accesa”, Italy.

***Melania parvula* f. *striata* Westerlund, 1886 [invalid]**

Original source. Westerlund 1886: 108.

Type locality. “Steinbrück” [Zidani Most], Slovenia.

Remarks. Junior homonym of *Melania striata* Sowerby, 1818, described from the Paleogene (Eocene?) of the United Kingdom.

† *Melanopsis (Hyphantria) striata* Handmann, 1887 [invalid]

Original source. Handmann 1887: 38, pl. 8, fig. 18.

Type horizon. Pannonian, zone B–D, late Miocene.

Type locality. “Leobersdorf”, Austria.

Remarks. Junior homonym of *Melanopsis maroccana striata* Pantanelli, 1886 (see Note 1).

Melanopsis maroccana var. *striatacarinata* Pantanelli, 1886

Original source. Pantanelli 1886b: 72, pl. 3, figs 24–25.

Type locality. “Accesa”, Italy.

Remarks. Originally written as “*striata-carinata*”.

Melanopsis maroccana var. *striatasulcata* Pantanelli, 1886

Original source. Pantanelli 1886b: 72, pl. 3, fig. 28.

Type locality. “Accesa”, Italy.

Remarks. Originally written as “*striata-sulcata*”.

† *Melanopsis (Melanosteira) striatula* Pavlović, 1927

Original source. Pavlović 1927: 62, pl. 7, figs 9–10.

Type horizon. Middle Pannonian, late Miocene.

Type locality. “У караџачких жуму neckoџума” [from the yellow sands of Karagača near Vrčin], Serbia.

Types. The illustrated syntype is stored in the Natural History Museum, Belgrade, coll. no. 212 (Milošević 1962: 23).

Melanopsis dufourii var. *stricta* Pallary, 1924

Original source. Pallary 1924: 249, pl. 15, fig. 4.

Type locality. Spain [no locality indicated].

Melanopsis buccinoidea var. *stricta* Pallary, 1939 [invalid]

Original source. Pallary 1939: 85, pl. 6, figs 64–65.

Type locality. “Environs de Beyrouth” [surroundings of Beirut], Lebanon.

Remarks. Junior homonym of *Melanopsis dufourii stricta* Pallary, 1924 (see Note 1).

† *Melanopsis stricturata* Brusina, 1892

Original source. Brusina 1892: 138.

Type horizon. Middle Pannonian, late Miocene.

Type locality. “Markuševac”, Croatia.

Types. The illustrated syntype is stored in the Croatian Natural History Museum, Zagreb, coll. no. 2496-142/1 (Milan et al. 1974: 97).

† *Boistelia stricturata* Jekelius, 1944 [invalid]

Original source. Jekelius 1944: 136, pl. 55, figs 1–14.

Type horizon. Early Pannonian, late Miocene.

Type locality. “Turislav-Tal bei Soceni” [Turislav valley near Soceni], Romania.

Remarks. Junior secondary homonym and junior synonym of *M. stricturata* Brusina, 1892 (see discussion in Neubauer et al. 2014a: 463).

Melanopsis costata var. *strigosa* Pallary, 1939

Original source. Pallary 1939: 90, pl. 5, figs 35–36.

Type locality. “Djishr ech Chegour” [Jisr Ash-Shughur], Syria.

Remarks. Heller and Sivan (2002a: 49) considered the taxon as a junior synonym of *Melanopsis multiformis* Blanckenhorn, 1897. Heller et al. (2005: 244) in turn considered it as a junior synonym of *Melanopsis costata* (Olivier, 1804).

† *Melanopsis sturii* Fuchs, 1873

Original source. Fuchs 1873: 21, pl. 4, figs 18–19.

Type horizon. Transdanubian, Pannonian, late Miocene.

Type locality. “Moosbrunn bei Wien; Tinnye bei Ofen” [Moosbrunn near Vienna (Austria); Tinnye (Hungary)].

Remarks. The name “*sturi*” as mentioned in Wenz (1929: 2835) is an incorrect subsequent spelling.

Fagotia (Microcolpia) daudebartii stussineri Schütt & Bilgin, 1974

Original source. Schütt and Bilgin 1974: 62, fig. 5.

Type locality. “River Pinios [Pineíos] in Thessalia, Greece, between Larissa and Tempe [Tempón] valley”, Greece.

Types. Senckenberg Forschungsinstitut und Naturmuseum Frankfurt, coll. no. SMF 111523a.

Remarks. The name “*situssineri*” as mentioned in Koşal-Şahin and Yıldırım (2007: 51) is an incorrect subsequent spelling.

***Melanopsis suarezi* Ahuir Galindo, 2016**

Original source. Ahuir Galindo 2016: 26.

Type locality. “Southeastern of Guelmin, at river Seyad basin”, Morocco.

Types. Museo Malacologico di Cupra Marittima, Italy; no number indicated.

† ***Melanopsis subaffinis* Pallary, 1916 [invalid]**

Original source. Pallary 1916: 82.

Type horizon. Pannonian, zone D, late Miocene.

Type locality. “Kottingbrunn [...] Ziegelei a” (Handmann 1882: 558), Austria.

Remarks. Replacement name for the alleged junior homonym *Melanopsis affinis* Handmann, 1882, “non Férussac, 1823”. The latter name is, however, unavailable from Férussac (1823) but was introduced validly by Pallary (1916), who referred to an illustration in Férussac (1823) (see discussion of *M. affinis*). This makes *M. subaffinis* an objective synonym of *M. affinis* Handmann, 1882 and *M. affinis* Pallary, 1916 a junior homonym of *M. affinis* Handmann, 1882. *M. subaffinis* was considered as a junior synonym of *M. bouei* by Wenz (1929: 2681).

***Melanopsis maroccana* var. *subangulata* Pallary, 1922**

Original source. Pallary 1922: 209.

Type locality. “Marrakech”, Morocco.

† ***Melanopsis subangulosa* Sandberger, 1875**

Original source. Sandberger 1870–1875: 559.

Type horizon. Kirchberg Formation, middle Burdigalian, early Miocene.

Type locality. “Kirchberg” [Illerkirchberg], Germany.

† ***Melanopsis pygmaea* f. *subaudebardi* Soós in Bartha, 1955**

Original source. Bartha 1955: 297, pl. 2, fig. 5.

Type horizon. Transdanubian, Pannonian, late Miocene.

Type locality. “Várpalota”, Hungary.

Remarks. Bartha clearly stated that the description was made by Lajos Soós.

† *Fagotia subbergeroni* Gozhik, 2002

Original source. Gozhik 2002: 54, pl. 3, figs 5, 7, 12, 16.

Type horizon. Middle Pontian (Dacian Basin), late Miocene.

Type locality. “Виноградівка” [Vynohradivka, Odes’ka Oblast’], Ukraine.

Types. Institute of Geological Sciences of the National Academy of Sciences of Ukraine, coll. no. 3185.

† *Melanopsis subbuccinoides* d’Orbigny, 1852

Original source. d’Orbigny 1852: 28.

Type horizon. Burdigalian, early Miocene.

Type locality. “Dax, St-Paul. Mandillot” (Grateloup 1840, in captions of plate “Mollusques terrestres et fluviatiles fossiles de Dax”), France.

Remarks. Introduced for *M. buccinoides* [sic] sensu Grateloup, 1840, non Olivier, 1801. The name “*sublucinoïdes*” as mentioned in Glibert (1962: 141) is an incorrect subsequent spelling.

† *Melanopsis pedemontana subcallosa* Wenz, 1928 [invalid]

Original source. Wenz 1928b: 220.

Type horizon. Late Burdigalian, early Miocene.

Type locality. “Colle torinesi” (Sacco 1889: 68) [Torino hills], Italy.

Remarks. Invalid replacement name for the homonym *M. taurinensis* Sacco, 1889, non *M. clava taurinensis* Sacco, 1889 (same work). According to Art. 24.1 and 57.7 the name proposed at higher rank takes precedence. This makes *subcallosa* an objective synonym of *M. taurinensis* Sacco, 1889, while the subspecies *M. clava taurinensis* Sacco, 1889 is still in need of a substitute name.

† *Melanopsis subcarinata* Deshayes in Férussac, 1839

Original source. Férussac 1819–1851: Mélanopsides fossiles, pl. 2, fig. 3 (altered caption by Deshayes delivered with Livraison 29; see introduction for details).

Type horizon. Late Villafranchian, Pleistocene (?).

Type locality. “D’Italie” (in the original captions, Férussac had given the locality as “entre Saint-Germinin et Carsoli” [between San Gemini and Carsoli, Italy]).

† *Melanopsis subcarinata* Morris in Forbes, 1856 [invalid]

Original source. Forbes 1856: 156, pl. 6, figs 5–6.

Type horizon. Headon Beds, Priabonian, Eocene.

Type locality. “Headon Hill”, United Kingdom.

Remarks. The description was evidently performed by Morris (see bottom of p. 156 in Forbes 1856). Junior homonym of *Melanopsis subcarinata* Deshayes in Férussac, 1851.

† *Melanopsis subcostata* d’Orbigny, 1850

Original source. d’Orbigny 1850: 301.

Type horizon. Sparnacian, early Ypresian, Eocene.

Type locality. d’Orbigny (1850) gave “Antheuil (Oise), Soissons”, Pacaud (2007) only “Antheuil”. Deshayes (1825), however, which the new name was based on, had only mentioned material from Soissons (along with occurrences of *M. costata* mentioned in other works), France.

Remarks. Introduced for *M. costata* sensu Deshayes, 1825, non Olivier, 1804.

Melanopsis subcostata Bourguignat, 1884 [invalid]

Original source. Bourguignat 1884: 137.

Type locality. “Dans l’Oronte” (Lamarck 1822: 168) [in the Orontes river], Syria?

Remarks. Introduced for *M. costata* sensu Lamarck, 1822, non Olivier, 1804. Junior homonym of *Melanopsis subcostata* d’Orbigny, 1850. Pallary (1920b: 110–111) synonymized the species with *M. costata* (Olivier, 1804).

† *Melanopsis constricta* f. *subcostata* Brusina, 1897 [invalid]

Original source. Brusina 1897: 7.

Type horizon. Late Cernikian, late Pliocene–early Pleistocene.

Type locality. “Novska (Bukovica)”, Croatia.

Types. Milan et al. (1974: 88) indicated a holotype, but it is uncertain whether the specimen was the only one Brusina had at hand (holotype by monotypy, Art. 73.1.2). The specimen is stored in the Croatian Natural History Museum, Zagreb, coll. no. 2993-639.

Remarks. Junior homonym of *Melanopsis subcostata* d’Orbigny, 1850. Wenz (1928a: 120) introduced *M. constricta nowskaensis* as replacement name.

***Melanopsis laevigata* var. *subcostulata* Pallary, 1904**

Original source. Pallary 1904: 37.

Type locality. “Sriratzel Cromfel sur la route de Rabat à Casablanca; Temslott, dans les canaux; Agagour dans l’Atlas” [Sriratzel Cromfel (?) at the road from Rabat to Casablanca; Temslott (?), in the channels; Agagour (?) in the Atlas Mts.], Morocco.

***Melanopsis saharica* var. *subcostulata* Pallary, 1912 [invalid]**

Original source. Pallary 1912: 15, figs 1–2.

Type locality. “De Ngoussa” [N’Goussa], Algeria.

Remarks. Junior homonym of *Melanopsis subcostulata* Pallary, 1904.

***Melanopsis praemorsa* f. *subcylindrica* Pérès, 1939**

Original source. Pérès 1939: 135.

Type locality. Several collection stations near Meknès, Fès, Aïn El Aouda and Aïn Chkef, Morocco (Pérès 1939: 140).

† *Melanopsis dufourii* var. *subfusiformis* Grateloup, 1838

Original source. Grateloup 1838: 142.

Type horizon. Burdigalian, early Miocene.

Type locality. “Dax. Mandillot, à Saint-Paul”, France.

Remarks. Introduced for a specimen of *M. dufourii* Férussac, 1822 from Dax illustrated in Férussac (1823: 154, pl. 8, fig. 5).

† *Melanopsis subfusiformis* Morris in Forbes, 1856

Original source. Forbes 1856: 155, pl. 6, figs 2–3.

Type horizon. Headon Beds, Priabonian, Eocene.

Type locality. “From the Headon series”, United Kingdom.

Remarks. The description was evidently performed by Morris (see bottom of p. 156 in Forbes 1856).

***Melanopsis maroccana* var. *subgraellsiana* Bourguignat, 1864**

Original source. Bourguignat 1864: 260.

Type locality. “Mascara; Oran”, Algeria.

Remarks. The name “*subgraellsii*” as used by Pallary (1920b: 106) is an incorrect subsequent spelling.

† *Melanopsis subhercynica* Mertin, 1939

Original source. Mertin 193: 207, pl. 6, fig. 5.

Type horizon. Heidelberg Formation, late Santonian, late Cretaceous.

Type locality. “Flugplatz Quedlinburg” [airfield at Quedlinburg], Germany.

Melanopsis subimpressa Pallary, 1928

Original source. Pallary 1928a: 262, pl. 6, figs 2–3.

Type locality. “Guefäit (Maroc oriental)”, Morocco.

Melanella elegans var. *sublaevis* Bourguignat, 1884

Original source. Bourguignat 1884: 15.

Type locality. “Au pont de la Save, près d’Agram” [at the bridge of the Sava river near Zagreb], Croatia.

† *Melanopsis sublanceolata* Kormos, 1905

Original source. Kormos 1905: 393, 440, pl. 2, fig. 9.

Type horizon. Late Pleistocene–early Holocene.

Type locality. “Püspökfürdő” [Băile 1 Mai, Lake Pețea], Romania.

Remarks. Neubauer et al. (2014d: 125) considered this taxon as a junior synonym of *Microcolpia parreyssii* (Philippi, 1847).

† *Melanopsis sublongata* Pallary, 1916

Original source. Pallary 1916: 82.

Type horizon. Lutetian, Eocene.

Type locality. “Au Nord d’Albas” (Doncieux 1908: 202), France.

Remarks. Replacement name for *M. elongata* Doncieux, 1908, non Férussac, 1822 (see Note 1). Wenz (1929: 2838) gave it as “*subelongata*”, which is an incorrect subsequent spelling.

***Melanopsis costata* var. *subnodata* Pallary, 1939**

Original source. Pallary 1939: 90, pl. 5, figs 30–32.

Type locality. “Djishr ech Chegour” [Jisr Ash-Shughur], Syria.

Remarks. Heller and Sivan (2002a: 49) considered the taxon as a junior synonym of *Melanopsis multiformis* Blanckenhorn, 1897. Heller et al. (2005: 244) in turn considered the species as a junior synonym of *Melanopsis costata* (Olivier, 1804).

***Melanopsis iraqensis* var. *subplicata* Pallary, 1939**

Original source. Pallary 1939: 89.

Type locality. Not explicitly stated but probably the same as for the species (“Tappah, à 3 km Est de Belad Sindjar et ‘Ain Haglan” [Tappah, 3 km east of Sinjar and Ain Haglan (?)], Iraq).

† *Melanopsis praemorsa* f. *subpraemorsa* Bogachev, 1908

Original source. Bogachev 1908: 246, 257, pl. 4, figs 8–13.

Type horizon. Late Sarmatian, Khersonian, late Miocene.

Type locality. “Chatma, district de Signakh, gouv. de Tiflis” [Chatma, district of Signaghi], Georgia.

Remarks. Introduced originally as “*sub-praemorsa*”.

† *Melanopsis subpraerosa* Andrusov, 1909

Original source. Andrusov 1909: 88, 160, pl. 5, figs 3–4.

Type horizon. Pontian (sensu stricto), late Miocene.

Type locality. “Adjipirdariaki, O. von Marasy, Tscharagan, [...] Sundi, [...] Chinasty” [Mount Adjipirdariaki, east of Marazy, Çarxan, Syundi, Chinasty canyon], Azerbaijan.

† *Melanopsis subpyrum* Penecke, 1884

Original source. Penecke 1884: 24, pl. 10, figs 19–20.

Type horizon. Cernikian, Pliocene.

Type locality. “Capla-Graben” [Čaplja trench near Slavonski Brod], Croatia.

† *Melania subrugosa* d'Orbigny, 1850 [invalid]

Original source. d'Orbigny 1850: 300.

Type horizon. Early Campanian, Cretaceous.

Type locality. “Martigues” (Matheron 1842: 293), France.

Remarks. D'Orbigny (1850) considered *Melanopsis rugosa* Matheron, 1842 a *Melania* and the species name a secondary homonym of *Melania rugosa* Lea, 1842. However, Matheron's species was published earlier (August 1842 vs. November/December 1842; for details on publication dates of Matheron's work see note at the beginning of the Reference section). *Melania subrugosa* is thus a junior objective synonym of *Melanopsis rugosa* Matheron, 1842.

Melanopsis subscalaris Bourguignat, 1884

Original source. Bourguignat 1884: 108.

Type locality. “Aïn-Fekan, source d'eau chaude entre Mascara et Saïda; dans Oued-M'Ilouya, sur la frontière du Maroc” (Bourguignat 1884: 260, 262) [‘Aïn Fekan, hot spring between Mascara and Saïda; river Moulouya, at the border to Morocco], Algeria.

Remarks. Introduced for *M. maroccana scalaris* sensu Bourguignat, 1884, non Gassies, 1856 (see Note 1).

† *Boistelia substricturata* Jekelius, 1944

Original source. Jekelius 1944: 136, pl. 55, figs 15–19.

Type horizon. Early Pannonian, late Miocene.

Type locality. “Turislav-Tal bei Soceni” [Turislav valley near Soceni], Romania.

Melanopsis subtilis Brot, 1878

Original source. Brot 1874–1879: 373, pl. 38, fig. 5d.

Type locality. “Perna Fluss in Ungarn” [not found], Hungary?

Remarks. Described and illustrated in synonymy of “*Hemisinus Esperi*” [now in *Esperiana*], based on a manuscript name by Parreyss. Treated as a valid name by Clessin (1890) and made thereby available (see Note 2).

† *Melanopsis subtilis* Pallary, 1920 [invalid]

Original source. Pallary 1920b: 110.

Type horizon. Transdanubian, Pannonian, late Miocene.

Type locality. “Tihany” (Fuchs 1870: 533), Hungary.

Remarks. Introduced for *M. aquensis* sensu Fuchs, 1870, non Grateloup, 1838. Junior homonym of *Melanopsis subtilis* Brot, 1879. That name was introduced in synonymy but became available from Clessin (1890), who treated it as valid species-group name (see Note 2). Wenz (1929: 2804) considered the taxon as a junior synonym of *M. petrovici* Brusina, 1893.

† *Melanopsis gorceixi* var. *subtilis* Magrograssi, 1928 [invalid]

Original source. Magrograssi 1928: 259, pl. 6, fig. 13.

Type horizon. Plio-Pleistocene.

Type locality. “Coo: molto frequente in tutte e due le zone fossilifere” [Kos island: very common in both areas rich in fossils, i.e., between Antimáchei and Pýli and in the northeast of the island, near Ágios Fokás], Greece.

Remarks. Junior homonym of *Melanopsis subtilis* Brot, 1879, which was originally introduced in synonymy but made available by Clessin (1890) who treated it as a valid name (see Note 2).

Melanopsis subtingitana Annandale, 1918

Original source. Annandale 1918: 163, pl. 20, figs 1–2.

Type locality. “Basra”, Iraq.

Types. Indian Museum, Calcutta, coll. no. 11390/2M.

Remarks. Annandale attributed the authority to Nevill, apparently based on a manuscript of that author.

† *Melanopsis subtuberculata* Pallary, 1916

Original source. Pallary 1916: 78.

Type horizon. Late Villafranchian, Pleistocene.

Type locality. “D’Italie” (Férussac 1823 had given the locality more precisely as “entre St.-Germini et Carsoli, et entre Otricoli et le Vigne” [between San Gemini and Carsoli, and between Otricoli and Le Vigne, Italy]).

Remarks. The name “*Melanopsis subtuberculata* Férussac” has been used as valid name by several authors (e.g., Wenz 1929: 2839), but was obviously not intended as species-group name by Férussac (1823). Esu and Girotti (1975: 251) considered the taxon as a junior synonym of “*M. affinis* Férussac, 1823”, which is not available either. The name *subtuberculata* became nevertheless available from Pallary (1916) who treated it as valid species-group name and associated it with an illustration in Férussac

(1823: pl. 7, fig. 11). See introduction for detailed discussion about the names used by Férussac (1823).

***Melanopsis subturrita* Azpeitia Moros, 1929**

Original source. Azpeitia Moros 1929: 192, pl. 4, figs 91–93, 95, pl. 5, figs 116, 118–120.

Type locality. “Alhama de Aragón, [...] principalmente en la fuente termal de las Dehesillas” [Alhama de Aragón, mainly in the hot spring of Dehesillas], Spain.

† ***Melanopsis subulata* Sowerby, 1822**

Original source. Sowerby 1821–1823: 36, pl. 332, fig. 8.

Type horizon. Eocene?

Type locality. “Isle of Wight”, United Kingdom.

Remarks. Originally the gender was indicated as masculine (“*subulatus*”), but *Melanopsis* is feminine, which is why the name must be “*subulata*”.

† ***Melanopsis dufourii* var. *subulata* Grateloup, 1838 [invalid]**

Original source. Grateloup 1838: 142, pl. 4, fig. 51.

Type horizon. Burdigalian, early Miocene.

Type locality. “Dax. Mandillot, à Saint-Paul”, France.

Remarks. Junior homonym of *Melanopsis subulata* Sowerby, 1822.

† ***Melanopsis buccinoidea* var. *subventricosa* Grateloup, 1828**

Original source. Grateloup 1828: 137.

Type horizon. Burdigalian, early Miocene.

Type locality. “Mandillot”, France.

† ***Melanopsis suskalovici* Pavlović, 1903**

Original source. Pavlović 1903: 156, pl. 3, figs 1–2.

Type horizon. Middle Miocene.

Type locality. “Из Бабиног Дола близу Скопља” (p. 155) [Babin Dol near Skopje], Macedonia.

Types. The illustrated syntype is stored in the Natural History Museum, Belgrade, coll. no. 1447 (Milošević 1962: 24).

† *Melanopsis synaniae* Esu & Girotti, 2015

Original source. Esu and Girotti 2015a: 68, figs 5–8.

Type horizon. Synania Formation, Pleistocene.

Type locality. “S-SW of Neos Erineos” [near Aigio], Greece.

Types. Senckenberg Forschungsinstitut und Naturmuseum Frankfurt, coll. no. SMF 345712.

† *Melanopsis szontaghi* Kormos, 1905

Original source. Kormos 1905: 392, 439, pl. 2, fig. 2.

Type horizon. Late Pleistocene–early Holocene.

Type locality. “Püspökfürdő” [Băile 1 Mai, Lake Peștea], Romania.

Remarks. Neubauer et al. (2014d: 125) considered this taxon as a junior synonym of *Microcolpia parreyssii* (Philippi, 1847).

† *Melanopsis tabulata* Hörnes, 1856

Original source. Hörnes 1851–1856: 600, pl. 49, fig. 15.

Type horizon. Badenian, middle Miocene.

Type locality. “Grund”, Austria.

Remarks. After Cossmann (1909: 152) this species belongs in the genus *Semisinus* P. Fischer, 1885, which is an unjustified emendation of *Hemisinus* Swainson, 1840 (Thiaridae).

“*Melanopsis tachitensis* Fér.” mentioned in De Cristofori and Jan (1832: Conchyliia terrestria et fluviatilia, p. 7) [unavailable]

Locality. “Austr.” [Australia].

Remarks. Nomen nudum, found only in the species list of De Cristofori and Jan (1832).

Melanopsis tanousi Bourguignat, 1884

Original source. Bourguignat 1884: 137.

Type locality. “Dans les cours d’eau de la plaine du Bahr-el-Houlé, non loin d’Ain-el-Mellaha” [in the rivers of the plains of the Hula valley, near Aïn Mallahah], Israel.

Remarks. Bourguignat (1884) denoted the authority as “Letourneux, 1882”, but there is no evidence that the description really derived from that author. Heller et al. (2005: 244) considered the species as a junior synonym of *Melanopsis lampra* Bourguignat, 1884.

† *Melanopsis clava* var. *taurinensis* Sacco, 1889 [invalid]

Original source. Sacco 1889: 67, pl. 2, fig. 9.

Type horizon. Late Burdigalian, early Miocene.

Type locality. “Colline torinesi” [Torino hills], Italy.

Remarks. Homonym of the simultaneously described *M. taurinensis* Sacco, 1889. Wenz (1928b) invalidly selected the variety as valid and introduced a replacement name for the species, but the taxon of higher rank takes precedence (Art. 24.1 and 57.7). No valid replacement name exists for the variety at present.

† *Melanopsis taurinensis* Sacco, 1889

Original source. Sacco 1889: 68, pl. 2, fig. 15.

Type horizon. Late Burdigalian, early Miocene.

Type locality. “Colle torinesi” [Torino hills], Italy.

Remarks. Homonym of the simultaneously described *M. clava taurinensis* Sacco, 1889. Wenz (1928b) invalidly selected the subspecies as valid and introduced *M. subcallosa* as replacement name for this species (junior objective synonym; see Art. 24.1 and 57.7).

† *Balanocochlis patula* var. *taurostriata* Sacco, 1895

Original source. Sacco 1895: 6, pl. 1, fig. 9.

Type horizon. Late Burdigalian, early Miocene.

Type locality. “Colli torinese” [Torino hills], Italy.

Remarks. Considered to belong in the genus *Amphimelania* by Wenz (1929: 2876). Note that Wenz gave the name as “*taurostricta*”, which is an incorrect subsequent spelling.

† *Melanopsis tchernovi* Heller & Sivan, 2002

Original source. Heller and Sivan 2002b: 619, fig. 3F.

Type horizon. Early Pleistocene.

Type locality. “Erq el-Ahmar” [locality also known as Gesher], Israel.

Types. Paleontology Collection of the Hebrew University of Jerusalem, coll. no. HUJ 9016.

† *Melanopsis tenuiplicata* Neumayr, 1880

Original source. Neumayr 1880c: 477, pl. 7, fig. 4.

Type horizon. Late Burdigalian–early Langhian, early–middle Miocene.

Type locality. “Seonica”, Bosnia and Herzegovina.

† *Melanopsis vidovici* var. *tenuis* Brusina, 1903 [invalid]

Original source. Brusina 1903: 114.

Type horizon. Late Pleistocene–early Holocene.

Type locality. “Bischofsbad” [Püspökfürdő, Băile 1 Mai, Lake Pețea], Romania.

Remarks. Junior objective synonym of *M. vidovici*: Brusina (1903) indicated it as the typical form of the species. Neubauer et al. (2014d: 125) considered this taxon as a junior synonym of *Microcolpia parreyssii sikorai* (Brusina, 1903).

† *Melanopsis tessellata* Brusina, 1902

Original source. Brusina 1902: pl. 6, figs 22–25.

Type horizon. Transdanubian, Pannonian, late Miocene.

Type locality. “Radmanest” [Rădmănești], Romania.

Types. The illustrated syntypes are stored in the Croatian Natural History Museum, Zagreb, coll. no. 2512-158/1-2 (Milan et al. 1974: 98).

Remarks. The name “*tesselata*” as mentioned in Wenz (1929: 2841) is an incorrect subsequent spelling.

† *Melanopsis (Homalia) textilis* Handmann, 1887

Original source. Handmann 1887: 15, pl. 1, figs 12–14.

Type horizon. Pannonian, zone B–D, late Miocene.

Type locality. “Leobersdorf”, Austria.

Remarks. Wenz (1929: 2758) considered the taxon as a junior synonym of *Melanopsis inermis* Handmann, 1882.

† *Melanopsis themaki* Brusina, 1903

Original source. Brusina 1903: 110.

Type horizon. Late Pleistocene–Holocene.

Type locality. “Bischofsbad” [Püspökfürdő, Băile 1 Mai, Lake Pețea], Romania.

Remarks. Neubauer et al. (2014d: 125) considered this taxon as a junior synonym of *Microcolpia parreyssii sikorai* (Brusina, 1903).

Hemisinus thermalis Brot, 1868

Original source. Brot 1868: 52, pl. 3, figs 14–15.

Type locality. “Carpazi [...], Miskolz [...]” [Carpazi (Italy); Miskolcz (Hungary)].

Remarks. Brot attributed the authority to “Titius (?) (Parreyss)”, probably based on an “in schedis” determination.

† *Melanopsis thomasi* Tournouër, 1877

Original source. Tournouër 1877: 275.

Type horizon. Late Pannonian, late Miocene.

Type locality. “Smendou” [Zighoud Youcef], Algeria.

Melanopsis tiaraeformis Pallary, 1936

Original source. Pallary 1936: 60, pl. 3, fig. 2.

Type locality. “Cascade d’Imouzer d’Agadir” [waterfall of Imouzzet at Agadir], Morocco.

“*Melanopsis tigris* Féruss.” mentioned in Adams and Adams (1853–1858: 310) [unavailable]

Locality. Not indicated.

Remarks. *Nomen nudum*, appears only in Adams and Adams (1854) without explanation.

† *Melanopsis tihanyensis* Wenz, 1928

Original source. Wenz 1928b: 219.

Type horizon. Transdanubian, Pannonian, late Miocene.

Type locality. “Tihany” (Fuchs 1870: 533), Hungary.

Remarks. Replacement name for the junior homonym *M. gradata* Fuchs, 1870. Both taxa were listed as junior synonyms of *M. brusinai* Lörentz, 1902 by Gillet and Marinescu (1971: 57).

† *Melanopsis timacensis* Živković, 1893

Original source. Živković 1893: 166.

Type horizon. Badenian, middle Miocene.

Type locality. “Vražognac unterhalb des Einflusses des Alapin in den Timok” [Vražognac, below confluence of the rivers Alapin and Timok], Serbia.

***Melanopsis tingitana* Morelet, 1864**

Original source. Morelet 1864: 155.

Type locality. “In Marocco” [no locality indicated].

† *Melanopsis tinnyensis* Wenz, 1919

Original source. Wenz 1919a: 73.

Type horizon. Middle Pannonian, late Miocene.

Type locality. “Tinnye” (Lörenthey 1902: 214), Hungary.

Remarks. Replacement name for *M. sinzowi* Lörenthey, 1902, non Brusina, 1885. Papp (1953b: 145) considered the taxon as a junior synonym of *Melanopsis rarisipina* Lörenthey, 1902. However, Papp (1953b) gave the older names *M. bouei spinosa* Handmann, 1882 and *M. bouei ventricosa* Handmann, 1882 in synonymy of *M. rarisipina*. While *ventricosa* Handmann is a homonym of *M. ventricosa* Neumayr, 1880 and thus invalid, the name *spinosa* is available and has priority over *rarisipina*.

† *Melanopsis parkinsoni* var. *titestiensis* Popescu-Voitești, 1910

Original source. Popescu-Voitești 1910: 362, pl. 21 (4), fig. 5.

Type horizon. Lutetian, Eocene.

Type locality. “Gropile Vulpilor près Titești” [Gropile Vulpilor (?) near Titești], Romania.

Remarks. Wenz (1929: 2568) ranked the variety as a subspecies of *Coptostylus albidus* (Lamarck, 1822) (Thiaridae).

† “*Melanopsis toriensis*” mentioned in Bogachev (1938: 13, 33, 53) [unavailable]

Horizon. Miocene.

Locality. “Tori bei Borshomi” [Tori near Borjomi], Georgia.

Remarks. The name was only mentioned in a species list by Bogachev without description or illustration.

† *Melanopsis oblonga* var. *torosa* Blanckenhorn in Blanckenhorn & Oppenheim, 1927

Original source. Blanckenhorn and Oppenheim 1927: 355, pl. 21, fig. 16.

Type horizon. Plio-Pleistocene.

Type locality. “Aus der pliocänen Dreissensiaschicht von Djisr esch-Schughr” [from the pliocene *Dreissena* layer at Jisr Ash-Shughur], Syria.

***Melanopsis torquilla* Pallary, 1920**

Original source. Pallary 1920a: 32.

Type locality. “Ras el Mà de Fès” [Ras el Ma near Fes], Morocco.

† *Melanopsis tortispina* Papp, 1953

Original source. Papp 1953b: 147, pl. 12, figs 18–20.

Type horizon. Pannonian, zone F–G, late Miocene.

Type locality. “Moosbrunn, N.-Ö.”, Austria.

Remarks. Appeared first as nomen nudum (as *M. bouei tortispina*) in Papp (1951: 167).

† *Melanopsis tothi* Brusina, 1903

Original source. Brusina 1903: 114.

Type horizon. Late Pleistocene–Holocene.

Type locality. “Bischofsbad” [Püspöckfürdő, Băile 1 Mai, Lake Peșea], Romania.

Remarks. Neubauer et al. (2014d: 125) considered this taxon as a junior synonym of *Microcolpia parreyssii sikorai* (Brusina, 1903).

† *Melanopsis tournoueri* Pallary, 1920

Original source. Pallary 1920b: 112.

Type horizon. Tafi Formation, early Pleistocene.

Type locality. “Antimaki” (Tournouër 1876: 449) [Antimácheia, Kos Island], Greece.

Remarks. Introduced for *M. cariosa* sensu Tournouër, 1876, non Linnaeus, 1758.

***Melanopsis transcaspia* Izzatullaev & Starobogatov, 1984**

Original source. Izzatullaev and Starobogatov 1984: 1478, fig. 1 (9).

Type locality. “Ручей Гяуре, колхоз им. 1 Мая Ашхабадского р-на” [Gyaure creek, Ashgabat district], Turkmenistan.

Types. Zoological Institute of Russian Academy of Sciences, St.-Petersburg; no number indicated.

† *Melanopsis transiens* Blanckenhorn, 1897

Original source. Blanckenhorn 1897: 135, pl. 10, fig. 19.

Type horizon. Plio-Pleistocene.

Type locality. “In der Dreissensiaschicht von Dschisr esch-Schurr” [in the *Dreissena* layers at Jisr Ash-Shughur], Syria.

***Melanopsis nodosa* var. *transiens* Cerulli-Irelli, 1914 [invalid]**

Original source. Cerulli-Irelli 1914: 185, pl. 15 (47), figs 8–9.

Type locality. “M. Mario: Farnesina”, Italy.

Remarks. Junior homonym of *Melanopsis transiens* Blanckenhorn, 1897. Pallary (1920b) introduced *M. cerullii* as replacement name. Girotti (1972: 232) considered the taxon as a junior synonym of “*M. affinis* Férussac”, which is, however, not an available name.

† *Melanopsis recurrens* var. *transitans* Brusina, 1874

Original source. Brusina 1874: 42.

Type horizon. Cernikian, Pliocene.

Type locality. “Bečić; Podvinje (Čaplja) [Čaplja trench near Slavonski Brod]; Novska; Kovačevac; Moslavina”, Croatia.

Types. The illustrated syntypes are stored in the Croatian Natural History Museum, Zagreb, coll. no. 2520-166/1-2 (Milan et al. 1974: 98).

† *Melanopsis trempensis* Bandel, 2000

Original source. Bandel 2000: 182, figs 143–145.

Type horizon. Valcarga Formation, Campanian, Cretaceous.

Type locality. “Pumanous slump exposed near Torallola”, Spain.

Types. Geological-Palaeontological Institute and Museum University of Hamburg, coll. no. 4270.

***Buccinum tricarinatum* Bruguière, 1789**

Original source. Bruguière 1789: 280.

Type locality. Not indicated.

Remarks. Commonly combined as *Melanopsis tricarinata* or as *M. dufourii tricarinata* (e.g., Azpeitia Moros 1929, Glaubrecht 1996, Ahuir Galindo 2015).

† *Melanopsis tricarinata* Sowerby in Fitton, 1836 [invalid]

Original source. Fitton 1836: 228, 346, pl. 22, fig. 4.

Type horizon. Weald Clay, early Cretaceous.

Type locality. “Punfield”, United Kingdom.

Remarks. Appeared first as nomen nudum (as “*Melania tricarinata*”) in Fitton (1824: 376). Junior secondary homonym of *Melanopsis tricarinata* (Bruguière, 1789). Sandberger (1870: 55) considered this species as a junior synonym of *Pleurocera strombiforme* (Schlotheim, 1820) (Pleuroceridae).

Melanopsis trifasciata Gray in Dieffenbach, 1843

Original source. Gray in Dieffenbach 1843: 263.

Type locality. “New Zealand, Bay of Islands, Waitanga Falls”, New Zealand.

Remarks. Originally the gender was indicated as masculine (“*trifasciatus*”), but *Melanopsis* is feminine, which is why the name must be “*trifasciata*”. Type species of *Zemelanopsis* Finlay, 1926. The name “*bifasciata*” as mentioned in Pallary (1926a: 76) is an incorrect subsequent spelling.

† “*Melanopsis themaki* var. *triflora*” mentioned in Brusina (1903: 110) [unavailable]

Horizon. Late Pleistocene–early Holocene.

Locality. “Bischofsbad” [Püspökfürdő, Băile 1 Mai, Lake Pețea], Romania.

Remarks. Nomen nudum (Brusina apparently considered the term self-explanatory). Neubauer et al. (2014d: 125) considered this taxon as a junior synonym of *Microcolpia parreyssii sikorai* (Brusina, 1903).

† “*Melanopsis tothi* var. *triflora*” mentioned in Brusina (1903: 114) [unavailable]

Horizon. Late Pleistocene–early Holocene.

Locality. “Bischofsbad” [Püspökfürdő, Băile 1 Mai, Lake Pețea], Romania.

Remarks. Nomen nudum (Brusina apparently considered the term self-explanatory). Brusina (1903) introduced several varieties with this name, apparently considering it only a descriptive term. Neubauer et al. (2014d: 125) considered this taxon as a junior synonym of *Microcolpia parreyssii sikorai* (Brusina, 1903).

† *Melanopsis tripaloi* Bourguignat, 1880

Original source. Bourguignat 1880: 35.

Type horizon. Langhian, middle Miocene.

Type locality. “Vallée de la Cettina” [Cetina river valley], Croatia.

Remarks. Wenz (1929: 2734) considered this taxon as a junior synonym of *Melanopsis geniculata* Brusina, 1874.

“*Melanopsis tripoliana*” mentioned in Brot (1874–1879: 419) [unavailable]

Locality. Not indicated.

Remarks. Nomen nudum, “in schedis” name from Tarnier listed in synonymy of *M. buccinoidea* (Olivier, 1801) by Brot (1879).

† *Melanopsis bouei trispina* Pashko, 1968

Original source. Pashko 1968: 137, 151, pl. 2, figs 1–8.

Type horizon. Tortonian, late Miocene.

Type locality. “Suita e Burrelit, prerja e Zallit të Gërmanit” [at Burrel, outcrop at the river Zalli të Germanit], Albania.

† *Melanopsis trivortiana* Locard, 1883

Original source. Locard 1883b: 59, pl. 3, fig. 5.

Type horizon. Mammal zone MN 14, Pliocene.

Type locality. “Trévoux” (p. 53), France.

† *Melanopsis defensa* var. *trochiformis* Fuchs, 1870

Original source. Fuchs 1870a: 354, pl. 14, figs 77–78.

Type horizon. Transdanubian, Pannonian, late Miocene.

Type locality. “Radmanest” [Rădmănești], Romania.

† *Melanopsis trojana* Hoernes, 1876

Original source. Hoernes 1876: 18, figs 8–15.

Type horizon. Late Sarmatian, Khersonian, late Miocene.

Type locality. “Renkiöi” [north of İntepe], Turkey.

† *Melanopsis trstenjaki* Brusina, 1884

Original source. Brusina 1884b: 55.

Type horizon. Early Langhian, middle Miocene.

Type locality. “Potravlje” (p. 47), Croatia.

Types. Milan et al. (1974: 98) indicated a holotype, but it is uncertain whether the specimen was the only one Brusina had at hand (holotype by monotypy, Art. 73.1.2). The specimen is stored in the Croatian Natural History Museum, Zagreb, coll. no. 2973-619.

“*Melanopsis truncata*” mentioned in De Cristofori and Jan (1832: *Conchylia terrestria et fluviatilia*, p. 7) [unavailable]

Locality. “Am. mer.” [South America].

Remarks. Nomen nudum, only mentioned in a species list by De Cristofori and Jan (1832).

Amphimelania (Melania) holandri var. *tschapecki* Westerlund, 1886

Original source. Westerlund 1886: 105.

Type locality. “Uferstellen der Sann bei Römerbad” (Tschapeck 1881: 102) [riversides of the Savinja river near Rimske Toplice], Slovenia.

Remarks. Introduced for a specimen of *Melania holandri* illustrated by Tschapeck (1881: 102, pl. 5, fig. d).

† *Melanopsis (Melanoptychia) tuberculata* Pavlović, 1927

Original source. Pavlović 1927: 60, pl. 7, figs 1–2.

Type horizon. Middle Pannonian, late Miocene.

Type locality. “У Врчинском хатару (долини потока Карагача)” [from Karagača river valley near Vrčin], Serbia.

Types. Perhaps stored in the Natural History Museum, Belgrade, but not mentioned in the catalogue of Milošević (1962).

Melanopsis tuberculata Yen, 1939 [invalid]

Original source. Yen 1939: 58, pl. 5, fig. 10.

Type locality. “See von Ta-li-fu (Yünnan)” [Lake Er Hai, Yunnan province], China.

Remarks. Junior homonym of *Melanopsis tuberculata* Pavlović, 1927. Probably not a Melanopsidae.

† *Melanopsis flammulata* var. *tuberosa* De Stefani, 1880

Original source. De Stefani 1880: 49, pl. 3, fig. 14.

Type horizon. Late Villafranchian, Pleistocene.

Type locality. “Monticiano”, Italy.

Remarks. Esu and Girotti (1975: 251) considered the taxon as a junior synonym of “*Melanopsis affinis* Férussac”, which is not an available name.

“*Melanopsis tumida*” mentioned in Brot (1874–1879: 420, pl. 45, fig. 3) [unavailable]

Locality. “Taurus-Gebiete” [Taurus region], Turkey.

Remarks. Introduced in synonymy of *M. buccinoidea* by Brot (1879), referring to an “in schedis” name by Parreyss (see Note 2).

Melanopsis tumida Pallary, 1916

Original source. Pallary 1916: 86.

Type locality. Stated to be uncertain by Philippi (1887, p. 40): “vermutlich bei Lebu oder Tubul gefunden” [probably found near Lebu or Tubul, Región del Biobío], Chile.

Remarks. Pallary (1916) introduced this replacement for the presumed secondary homonym *Melanopsis obesa* (Philippi, 1887), non Gassies, 1856. Despite a valid nomenclatural act, the name is hardly useful because Philippi’s species is certainly no Melanopsidae.

Melanopsis tunetana Morlet, 1881

Original source. Morlet 1881: 396, pl. 6, figs 3–4.

Type locality. “Tozeur, Kriz”, Tunisia.

Remarks. The name “*tuneata*” as mentioned in Chevallier (1969: 286) is an incorrect subsequent spelling.

Melanopsis praemorsa f. *turbinata* Pérès, 1939

Original source. Pérès 1939: 134.

Type locality. “Station 31. Daïa Afourgagh près d’Annosseur. Station 144. Oued Sidi Raba à 6 kilomètres environ en amont de son confluent avec le Bou-Regreg” [Station 31 at Dayat Afergagh near Anosseur; station 144 at Oued Sidi Raho, approximately 6 km upstream of its confluence with the Bou Regreg], Morocco.

***Melanopsis costata* var. *turcica* Mousson, 1874**

Original source. Mousson 1874: 33.

Type locality. “Dans le Karasu, affluent du lac d’Antioche (p. 33); environs de Samava (p. 49; there treated as distinct species)” [in the Karasu river, tributary of the Lake Anuk (also as Amik) (Turkey); surroundings of As Samawah (Iraq)].

Remarks. Mousson attributed the authority to Parreyss, but there is no evidence that the description really derived from that author.

† ***Melanopsis turgida* Fischer-de-Waldheim, 1837**

Original source. Fischer-de-Waldheim 1837: 131, pl. 18 [in text err. as “VXIII”], fig. 13.

Type horizon. Jurassic?

Type locality. “Miatchkova” [Myachikovo?], Russia.

Remarks. Certainly not a *Melanopsis*.

***Melanopsis turgida* Brot, 1878 [invalid]**

Original source. Brot 1874–1879: 373, pl. 38, fig. 5c.

Type locality. “Glina Fluss in Ungarn” [Glina river], Croatia.

Remarks. Described and illustrated in synonymy of “*Hemisinus Esperi*” [now in *Esperiana*], based on a manuscript name by Parreyss. Clessin (1890) treated it as valid name and made it available thereby (see Note 2). Junior homonym of *Melanopsis turgida* Fischer-de-Waldheim, 1837. Starobogatov et al. (1992: 60) considered the species as a junior synonym of *Fagotia* [= *Esperiana*] *esperi* (Férussac, 1823).

***Melanopsis turgida* Pallary, 1928 [invalid]**

Original source. Pallary 1928a: 259, pl. 5, figs 11–12.

Type locality. “Moulai Taïeb et Taourirt du Zâ (Maroc oriental)” [Moulay Taïeb and in Oued Za at Taourirt (?) (eastern Morocco)], Morocco.

Remarks. Junior homonym of *Melanopsis turgida* Fischer-de-Waldheim, 1837.

† ***Melanopsis turislavica* Jekelius, 1944**

Original source. Jekelius 1944: 131, pl. 49, figs 5–6.

Type horizon. Early Pannonian, late Miocene.

Type locality. “Turislav-Tal bei Soceni” [Turislav valley near Soceni], Romania.

***Melanopsis turkmenica* Izzatullaev & Starobogatov, 1984**

Original source. Izzatullaev and Starobogatov 1984: 1478, fig. 1 (10), fig. 2 (4).

Type locality. “Закаспийская область” [Transcaspiian Region], Russia.

Types. Zoological Institute of Russian Academy of Sciences, St.-Petersburg; no number indicated.

† *Melanopsis turricula* Matheron, 1842

Original source. Matheron 1842: 294, pl. 37, figs 15–16.

Type horizon. Early Campanian, Cretaceous.

Type locality. Not stated after the description, as Matheron used to do it for the other species. Although he did not denote an exact locality, he mentioned earlier in text the occurrence of the species in deposits situated “vers Vitrolles et Martigues” (p. 148–149) [towards Vitrolles and Martigues, France].

***Melanopsis (Mesopotamia) mesopotamica* var. *turricula* Pallary, 1939 [invalid]**

Original source. Pallary 1939: 100, pl. 5, fig. 7.

Type locality. “‘Ain Arouss” [‘Ayn al ‘Arūs, near Tall Abyaḍ], Syria.

Remarks. Junior homonym of *Melanopsis turricula* Matheron, 1842.

† *Melanopsis (Canthidomus) phanesiana* var. *turriculata* Magrograssi, 1928

Original source. Magrograssi 1928: 260, pl. 6, fig. 19.

Type horizon. Plio-Pleistocene.

Type locality. “Coo” [Kos island], Greece.

***Melanopsis turriiformis* Picard, 1934**

Original source. Picard 1934: 123, pl. 8, figs 15–27.

Type locality. “Jarmukmündung” [Yarmouk river mouth], Jordan/Israel.

***Melanopsis cariosa* var. *turrita* Rossmässler, 1854 [invalid]**

Original source. Rossmässler 1854–1859: 33, pl. 68, fig. 846.

Type locality. “Im Gebiet des unteren Guadalquivir” [in the lower Guadalquivir river], Spain.

Remarks. Rossmässler (1854) based this new variety entirely on *M. sevilensis* Grateloup, 1840, arguing that “Grateloup elevated this variety to species level without reason”. Thus, *M. turrita* is an objective synonym of *M. sevilensis*.

† *Melanopsis (Canthidomus) turrita* Handmann, 1887 [invalid]

Original source. Handmann 1887: 32, pl. 7, fig. 13.

Type horizon. Pannonian, zone B–D, late Miocene.

Type locality. “Leobersdorf”, Austria.

Remarks. Junior homonym of *Melanopsis turrita* Rossmässler, 1854. Pallary (1916: 84) introduced *M. hispidula* as replacement name.

† *Melanoptychia turrita* Jekelius, 1944 [invalid]

Original source. Jekelius 1944: 138, pl. 57, figs 1–21.

Type horizon. Early Pannonian, late Miocene.

Type locality. “Turislav-Tal bei Soceni” [Turislav valley near Soceni], Romania.

Remarks. Junior secondary homonym of *Melanopsis turrita* Rossmässler, 1854 and junior synonym of *M. turrita* Handmann, 1887, for which Pallary (1916: 84) had introduced *M. hispidula* as replacement name (see also Neubauer et al. 2014a: 463).

† *Pseudofagotia turrita* Anistratenko, 1993

Original source. Anistratenko 1993: 73, textfig. 2.

Type horizon. Duab Beds, middle to late Kimmerian, Pliocene.

Type locality. “Okp. c. Mokʼvu, Очамчирский р-н” [near the village Mokʼvi, Ochamchirskiy rayon], Georgia.

Types. Schmalhausen Institute of Zoology of National Academy of Sciences of Ukraine, Kiev; no number indicated.

† *Melanopsis conemenosiana* var. *turritella* Pallary, 1920

Original source. Pallary 1920b: 112.

Type horizon. Plio-Pleistocene.

Type locality. “Preveza in Epirus” (Oppenheim 1891: 469), Greece.

Remarks. Replacement name for *Melanopsis boettgeri* Oppenheim, 1891, non Klika, 1891.

***Melanopsis tutulata* Pallary, 1928**

Original source. Pallary 1928a: 271, pl. 5, figs 8–10.

Type locality. “Berguent; Aoùllout; Ras el Mâ de Fès; O. Chkef près Fès” [Aïn Beni Mathar, Aïn Aoullout, Ras El Ma, Oued Aïn Chkef in Fes], Morocco.

***Microcolpia ucrainica* Starobogatov, Alexenko & Levina, 1992**

Original source. Starobogatov et al. 1992: 69, textfig. 3 (20).

Type locality. “Из Южногo Буга” [in the southern Bug river], Ukraine.

Types. Zoological Institute of Russian Academy of Sciences, St.-Petersburg; no number indicated.

† *Melanopsis themaki* var. *unicarinata* Brusina, 1903

Original source. Brusina 1903: 111.

Type horizon. Late Pleistocene–early Holocene.

Type locality. “Bischofsbad” [Püspökfürdő, Băile 1 Mai, Lake Peșea], Romania.

Remarks. Neubauer et al. (2014d: 125) considered this taxon as a junior synonym of *Microcolpia parreyssii sikorai* (Brusina, 1903).

† *Melanopsis uncinata* Blanckenhorn, 1897

Original source. Blanckenhorn 1897: 119, pl. 9, figs 41–44.

Type horizon. Plio-Pleistocene.

Type locality. “In der obersten Melanopsiden-Thonbank des rechten Orontesufers bei Dschir esch-Schurr” [in the uppermost clay bank at the right riverside of the Orontes near Jisr Ash-Shughur], Syria.

Remarks. Introduced as “n. mut.” but clearly as a binomen and hence not infrasubspecific in the sense of ICZN Art. 45.6.

† *Melanopsis tothi* var. *uningulata* Brusina, 1903

Original source. Brusina 1903: 114.

Type horizon. Late Pleistocene–early Holocene.

Type locality. “Bischofsbad” [Püspökfürdő, Băile 1 Mai, Lake Peșea], Romania.

Remarks. Neubauer et al. (2014d: 125) considered this taxon as a junior synonym of *Microcolpia parreyssii sikorai* (Brusina, 1903).

† *Melanopsis hazayi* var. *unifilosa* Brusina, 1903 [unresolved]

Original source. Brusina 1903: 113.

Type horizon. Late Pleistocene–early Holocene.

Type locality. “Bischofsbad” [Püspökfürdő, Băile 1 Mai, Lake Peșea], Romania.

Remarks. Brusina (1903) introduced several varieties with this name, apparently considering it only a descriptive term; the homonymy issue needs to be solved by a First Reviser. Currently, all of them are considered junior synonyms of *Microcolpia parreyssii sikorai* (Brusina, 1903) (Neubauer et al. 2014d: 125).

† *Melanopsis sikorai* var. *unifilosa* Brusina, 1903 [unresolved]

Original source. Brusina 1903: 112.

Type horizon. Late Pleistocene–early Holocene.

Type locality. “Bischofsbad” [Püspökfürdő, Băile 1 Mai, Lake Peșea], Romania.

Remarks. Brusina (1903) introduced several varieties with this name, apparently considering it only a descriptive term; the homonymy issue needs to be solved by a First Reviser. Currently, all of them are considered junior synonyms of *Microcolpia parreyssii sikorai* (Brusina, 1903) (Neubauer et al. 2014d: 125).

† *Melanopsis themaki* var. *unifilosa* Brusina, 1903 [unresolved]

Original source. Brusina 1903: 110.

Type horizon. Late Pleistocene–early Holocene.

Type locality. “Bischofsbad” [Püspökfürdő, Băile 1 Mai, Lake Peșea], Romania.

Remarks. Brusina (1903) introduced several varieties with this name, apparently considering it only a descriptive term; the homonymy issue needs to be solved by a First Reviser. Currently, all of them are considered junior synonyms of *Microcolpia parreyssii sikorai* (Brusina, 1903) (Neubauer et al. 2014d: 125).

† *Melanopsis tothi* var. *unifilosa* Brusina, 1903 [unresolved]

Original source. Brusina 1903: 114.

Type horizon. Late Pleistocene–early Holocene.

Type locality. “Bischofsbad” [Püspökfürdő, Băile 1 Mai, Lake Peșea], Romania.

Remarks. Brusina (1903) introduced several varieties with this name, apparently considering it only a descriptive term; the homonymy issue needs to be solved by a First Reviser. Currently, all of them are considered junior synonyms of *Microcolpia parreyssii sikorai* (Brusina, 1903) (Neubauer et al. 2014d: 125).

† *Melanopsis vacua* Vidal, 1874

Original source. Vidal 1874: 236, pl. 3, fig. 15.

Type horizon. Maastrichtian, Cretaceous.

Type locality. “Isona”, Spain.

† *Melanopsis valdeci* Brusina, 1902

Original source. Brusina 1902: pl. 6, figs 18–21.

Type horizon. Early Cernikian, early Pliocene.

Type locality. “Čerević”, Serbia.

Types. The illustrated syntypes are stored in the Croatian Natural History Museum, Zagreb, coll. no. 2511-157/1-4 (Milan et al. 1974: 98).

† *Melanopsis kleinii* var. *valentinensis* Fontannes, 1881

Original source. Fontannes 1881: 1015, pl. 1, figs 7–9.

Type horizon. Mammal zone MN 10–12, late Miocene.

Type locality. “Montvendre”, France.

† *Melanopsis vandevelde* Bukowski, 1892

Original source. Bukowski 1892: 249.

Type horizon. Salakos Formation, Pliocene.

Type locality. “Rhodos” (locality specified as “Kalavarda” in Bukowski 1893), Greece.

***Melanopsis vanrossomi* Pallary, 1936**

Original source. Pallary 1936: 58, pl. 3, fig. 7.

Type locality. “Souk el Hâd des Aït Souhab, dans l’oued Gough, et Tanalt, à quelques kilomètres au sud du précédent dans l’Anti Atlas occidental” [Souk el Hâd of Aït Souhab (?), in the Oued Gough (?), and Tanalt, a few kilometers south of the former in the western Anti-Atlas], Morocco.

***Melania* (*Melanopsis*) *variabilis* Von dem Busch in Philippi, 1847**

Original source. Philippi 1847: 175, pl. 4, figs 7–8, 10.

Type locality. “Schiraz et Persepolis Persiae” [Shiraz and Persepolis, Fars Province], Iran.

“*Melania crassa* var. *variceata* Sandri” mentioned in Brusina (1866: 106) [unavailable]

Locality. “Cetina”, Croatia.

Remarks. Nomen nudum, listed by Brusina without any explanation.

† ***Melanopsis varicosa* Handmann, 1882**

Original source. Handmann 1882: 553.

Type horizon. Pannonian, zone D, late Miocene.

Type locality. “Kottingbrunn [...] Ziegelei a”, Austria.

***Melanopsis variegata* Morelet, 1857**

Original source. Morelet 1857: 33.

Type locality. “[Ad Sanctam-Mariam de Balade]” [Balade], New Caledonia.

Remarks. Brot (1879: 444) considered the taxon as a junior synonym of *Melanopsis frustulum* Morelet, 1857.

“*Melania holandri* var. *variegata* Grimmer” mentioned in Brot (1874–1879: 12) and Clessin (1890: 676) [unavailable]

Locality. Not indicated.

Remarks. Nomen nudum, listed in synonymy of “*Melania holandri*” [sic] (Brot 1874) and “*Melania holandri* var. *aequata*” (Clessin 1890), respectively.

***Melanopsis pseudoferussaci* var. *vaucheri* Pallary, 1899**

Original source. Pallary 1899: 140, pl. 9, fig. 12.

Type locality. “Fez” [Fes], Morocco.

† ***Melanopsis ventricosa* Neumayr, 1880**

Original source. Neumayr 1880b: 290, pl. 1, fig. 9.

Type horizon. Plio-Pleistocene.

Type locality. “Zwischen Pylle und Antimachia” [between Pýli and Antimácheia, Kos Island], Greece.

Remarks. Willmann (1981: 171) considered this taxon as a junior synonym of *Melanopsis sporadum* Tournouër, 1876.

† *Melanopsis bouei* var. *ventricosa* Handmann, 1882 [invalid]

Original source. Handmann 1882: 557.

Type horizon. Pannonian, zone D, late Miocene.

Type locality. “Kottingbrunn [...] Ziegelei a”, Austria.

Remarks. Junior homonym of *Melanopsis ventricosa* Neumayr, 1880. Wenz (1929: 2674) considered the taxon as a junior synonym of *M. bouei* Férussac, 1823.

“*Melanopsis dufouri* var. *cossoni* f. *ventricosior*” mentioned in Westerlund (1886: 126) [unavailable]

Locality. “Bei Valencia” [near Valencia], Spain.

Remarks. Introduced as infrasubspecific taxon (forma below a variety), which is not ruled by the provisions of the Code. Moreover, the name is a nomen nudum; Westerlund apparently considered the name a descriptive term (*ventricosior* = Latin “more bulbous”).

Melanopsis cariosa var. *ventrosa* Bourguignat, 1884 [unresolved]

Original source. Bourguignat 1884: 151.

Type locality. “Dans les aqueducs de Séville et dans le Guadalquivir” [in the aqueducts of Sevilla and in the Guadalquivir river], Spain.

Remarks. Homonym of the simultaneously published name *Melanopsis costata ventrosa* Bourguignat, 1884. This case requires the action of a First Reviser.

Melanopsis costata var. *ventrosa* Bourguignat, 1884 [unresolved]

Original source. Bourguignat 1884: 141.

Type locality. Not indicated, but apparently the specimens derive from somewhere along the Jordan river or the Sea of Galilee.

Remarks. Homonym of the simultaneously published name *Melanopsis cariosa ventrosa* Bourguignat, 1884. This case requires the action of a First Reviser.

† *Melanopsis venusta* Pallary, 1916

Original source. Pallary 1916: 80.

Type horizon. Pannonian, zone D, late Miocene.

Type locality. “Kottingbrunn [...] Ziegelei a” (Handmann 1882: 557), Austria.

Remarks. Replacement name for *M. nodosa* Handmann, 1882, non Férussac, 1822. Wenz (1929: 2681) considered this taxon as a junior synonym of *Melanopsis bouei* Férussac, 1823.

† “*Melanopsis impressa vera*” mentioned in Schréter (1912: 155) [unavailable]

Horizon. Miocene.

Locality. “Boesciwiese, östlich von Kohldorf [...] in dem von Norden, vom Balomberge hinablaufenden Seitengraben” [Poiana Boistii (?), east of Cărbunari, in the trench running down from the Mt. Balom (?) in the north], Hungary.

Remarks. Nomen nudum, appears without explanation in the text. Possibly, Schréter wanted to denote that it is the “real” *M. impressa* and not a variety. Wenz (1929: 2753) listed the record as a junior synonym of *M. impressa*.

Melanopsis vespertina Bourguignat, 1884

Original source. Bourguignat 1884: 124.

Type locality. “Lalla Maghnia, sur la frontière du Maroc” [Maghnia, at the border to Morocco], Morocco or Algeria.

† “*Melanopsis vetusta*” mentioned in Sandberger (1870: 88–89) [unavailable]

Horizon. Unknown.

Locality. “Plan d’Aups (Var)” [Plan-d’Aups-Sainte-Baume, Dép. Var].

Remarks. Nomen nudum, used by Matheron in correspondence with Sandberger (1870), who gave it as a junior synonym of *Paludomus lyra* var. *calva* (Paludomidae), which he described there as new.

† *Melanopsis vicentina* Oppenheim, 1890

Original source. Oppenheim 1890a: 135, pl. 4, figs 1–1b.

Type horizon. Ronca Beds, Bartonian, Eocene.

Type locality. “Lovara di Tressino, Mussolon, Monte Pulli bei Valdagno” [Lovara, Muzzolon, Monte Pulli near Valdagno], Italy.

† *Melanopsis vidovici* Brusina, 1903

Original source. Brusina 1903: 113.

Type horizon. Late Pleistocene–Holocene.

Type locality. “Bischofsbad” [Püspökfürdő, Băile 1 Mai, Lake Peșea], Romania.

Remarks. Neubauer et al. (2014d: 125) considered this taxon as a junior synonym of *Microcolpia parreyssii sikorai* (Brusina, 1903).

† *Melanopsis vilicici* Brusina, 1902

Original source. Brusina 1902: pl. 6, fig. 1.

Type horizon. Early–middle Miocene.

Type locality. “Kirin (Stipan)”, Croatia.

Types. Milan et al. (1974: 98) indicated a holotype, but it is uncertain whether the specimen was the only one Brusina had at hand (holotype by monotypy, Art. 73.1.2). The specimen is stored in the Croatian Natural History Museum, Zagreb, coll. no. 2502-148.

Microcolpia vileserriana Bourguignat, 1884

Original source. Bourguignat 1884: 58.

Type locality. “Dans les cours d’eau aux environs d’Ismidt, en Anatolie” [in rivers around İzmit], Turkey.

Melanopsis liocephala var. *villica* Pallary, 1936

Original source. Pallary 1936: 57.

Type locality. Not indicated, but probably in Morocco.

† *Melanopsis vincta* Blanckenhorn, 1897

Original source. Blanckenhorn 1897: 115, pl. 9, figs 6–7.

Type horizon. Plio-Pleistocene.

Type locality. “In der ersten Thonbank des linken Orontesufers bei Dschisr esch-Schurr” [in the first clay bank at the left riverside of the Orontes near Jisr Ash-Shughur], Syria.

Remarks. Introduced as “n. mut.” but clearly as a binomen and hence not infrasubspecific in the sense of ICZN Art. 45.6.

† *Melanopsis vindobonensis* Fuchs in Fuchs & Karrer, 1870

Original source. Fuchs and Karrer 1870: 140, textfig. 5.

Type horizon. Pannonian, zone E, late Miocene.

Type locality. “Zu Brunn, Inzersdorf, Rothneusiedel und Wien” [in Brunn am Gebirge, Inzersdorf and Rothneusiedl (today both localities lie within Vienna), and Vienna], Austria.

† *Melanopsis viquesneli* Pavlović, 1932

Original source. Pavlović 1932: 240, 247, pl. 1, figs 6–7.

Type horizon. Pontian (Dacian Basin), late Miocene–Pliocene.

Type locality. “Села Дрсника” [village Drsnik], Kosovo.

Types. Perhaps stored in the Natural History Museum, Belgrade, but not mentioned in the catalogue of Milošević (1962).

† *Melanopsis visianiana* Brusina, 1874

Original source. Brusina 1874: 37, pl. 1, figs 7–8.

Type horizon. Langhian, middle Miocene.

Type locality. “Miočić”, Croatia.

Types. Milan et al. (1974: 99) indicated a holotype, but it is uncertain whether the specimen was the only one Brusina had at hand (holotype by monotypy, Art. 73.1.2). The specimen is stored in the Croatian Natural History Museum, Zagreb, coll. no. 3205-851.

† *Melanopsis vitalisi* Strausz, 1942

Original source. Strausz 1942: 92, pl. 5, figs 14, 20, 29.

Type horizon. Transdanubian, Pannonian, late Miocene.

Type locality. The locality was indicated as number “85” by Strausz (1942: 63, 101), which refers to the *C. balatonica* horizon at Nyárád, Hungary (p. 23).

† *Melanopsis vitezovici* Brusina, 1902

Original source. Brusina 1902: pl. 5, figs 61–64.

Type horizon. Langhian, middle Miocene.

Type locality. “Džepe” [Džepi], Bosnia and Herzegovina.

Types. Neubauer et al. (2016c: 280) designated a lectotype, which is stored in the Croatian Natural History Museum, Zagreb, coll. no. 2501-147/1-4.

† *Melanopsis vittata* Pallary, 1916

Original source. Pallary 1916: 86.

Type horizon. Pannonian, zone D, late Miocene.

Type locality. “Kottingbrunn [...] Ziegelei a” (Handmann 1882: 560), Austria.

Remarks. Replacement name for *M. fasciata* Handmann, 1882, non Gassies, 1874. Wenz (1929: 2741) considered the taxon as a junior synonym of *Melanopsis haueri* Handmann, 1882.

† *Melanopsis vitzoui* Porumbaru, 1881

Original source. Porumbaru 1881: 27, pl. 8, fig. 4.

Type horizon. Early Romanian, Pliocene.

Type locality. “Bucovatzu” [Bucovăț], Romania.

Remarks. The name “*vitzui*” as mentioned in Pallary (1926a: 93) is an incorrect subsequent spelling.

***Melanopsis vondeli* Pallary, 1928**

Original source. Pallary 1928b: 16, pl. 2, figs 9–14.

Type locality. “Beni Mellal, dans l’oued Taguenout” [Beni Mellal, in the Oued Taguenout (?)], Morocco.

† *Melanopsis vrcinensis* Neubauer, Harzhauser, Georgopoulou, Mandić & Kroh, 2014

Original source. Neubauer et al. 2014a: 457.

Type horizon. Middle Pannonian, late Miocene.

Type locality. “У Капазачу” (Pavlović 1927: 60) [from Karagača near Vrcin], Serbia.

Types. Milošević (1962: 23) indicated that the syntype illustrated by Pavlović is stored in the Natural History Museum, Belgrade, coll. no. 209.

Remarks. Replacement name for *M. glabra* Pavlović, 1927, non Brusina, 1874 (see Note 1).

***Melanopsis wagneri* Roth, 1839**

Original source. Roth 1839: 24, pl. 2, fig. 11.

Type locality. “Smyrnae” [Izmir], Turkey.

Remarks. Martens (1874: 32) considered the taxon as a junior synonym of *Melanopsis praemorsa* (Linnaeus, 1758).

***Melanopsis waitaraensis* Marwick, 1926**

Original source. Marwick 1926: 317, pl. 73, figs 1–2.

Type locality. “Coast south side Wai-iti Stream”, New Zealand.

† *Melanopsis wilhelmi* Esu & Girotti, 2015

Original source. Esu and Girotti 2015b: 150, figs 1–2.

Type horizon. Late early Pleistocene.

Type locality. “Stirone river section, between Laurano and S. Nicomede (Emilia)”, Italy.

Types. Senckenberg Forschungsinstitut und Naturmuseum Frankfurt, coll. no. SMF 345834.

† *Melanopsis wolfgangfischeri* Neubauer, Harzhauser, Kroh, Georgopoulou & Mandic, 2014

Original source. Neubauer et al. 2014c: 17.

Type horizon. Pannonian, zone B–D, late Miocene.

Type locality. “Leobersdorf” (Handmann 1887), Austria.

Types. See statement for *M. martiniana rugosa* Handmann, 1887.

Remarks. Replacement name for *M. martiniana rugosa* Handmann, 1887, non Matheron, 1842 (see Note 1).

† *Melanopsis wuenschmanni* Mertin, 1939

Original source. Mertin 193: 206, pl. 6, fig. 3.

Type horizon. Heidelberg Formation, late Santonian, late Cretaceous.

Type locality. “Flugplatz Quedlinburg” [airfield at Quedlinburg], Germany.

† *Fagotia wuesti* Meijer, 1990

Original source. Meijer 1990: 152, pl. 2, figs 1–2.

Type horizon. Bavelian Complex, Pleistocene.

Type locality. “Clay-pit North of the village of Bavel (province of Noord Brabant, The Netherlands)”, Netherlands.

Types. Rijks Geologische Dienst, Haarlem, The Netherlands; no number indicated.

Remarks. Glöer (2002: 79) assigned the species to *Esperiana* (*Microcolpia*).

Melanopsis zarudnyi Izzatullaev & Starobogatov, 1984

Original source. Izzatullaev and Starobogatov 1984: 1479, fig. 1 (12), fig. 3 (6).

Type locality. “Золотоу кӀлоч околo АухабӀгa” [Zolotoy spring near Ashgabat], Turkmenistan.

Types. Zoological Institute of Russian Academy of Sciences, St.-Petersburg; no number indicated.

† ?*Melanopsis zea* Pezant, 1908

Original source. Pezant 1908: 200, pl. 7, figs 23–24.

Type horizon. Bartonian, Eocene.

Type locality. “Monneville”, France.

Remarks. Probably not a Melanopsidae.

Melanopsis zebra De Cristofori & Jan, 1832

Original source. De Cristofori and Jan 1832: 4.

Type locality. “Guinea”, indicated in the previous part of the same work (“Conchylia terrestria et fluviatilia [...]”, p. 7; there, the name is a nomen nudum).

Remarks. Probably not a Melanopsidae.

Melanopsis fasensis var. *zebrina* Pallary, 1920

Original source. Pallary 1920c: 148.

Type locality. “Sefrou”, Morocco.

Melanopsis zelandica Gould, 1847

Original source. Gould 1847: 225.

Type locality. “New Zealand” [no locality indicated].

Remarks. Suter (1893: 139) considered the taxon as a junior synonym of *Zemelanopsis trifasciata* (Gray in Dieffenbach, 1843). The name “*zelandica*” as mentioned in Hutton (1883: 67) is an incorrect subsequent spelling.

† *Melanopsis* (*Canthidomus*) *zitteli* Neumayr, 1869

Original source. Neumayr 1869: 357, pl. 11, figs 4–5.

Type horizon. Langhian, middle Miocene.

Type locality. “Miocic” [Miočić], Croatia.

Types. Illustrated syntypes are stored at the Geological Survey Austria, Vienna, coll. no. 1869/01/5/1-2.

Melanopsis maroccana var. *zonata* Gassies, 1856

Original source. Gassies 1856: 12, figs 5–6.

Type locality. “L’Aïn-Kadra, sur les hauts plateaux de l’Atlas, à deux mètres des Chots” [in the ‘Aïn Khadr, in the highlands of the Atlas, two meters in the Chott (probably he meant the lake Chott el Hodna)], Algeria.

***Melanopsis praemorsa* var. *zonata* Bourguignat, 1864 [invalid]**

Original source. Bourguignat 1864: 264.

Type locality. “Biskra”, Algeria.

Remarks. Junior homonym of *Melanopsis maroccana zonata* Gassies, 1856 (see Note 1), which Bourguignat listed as well (p. 260–261).

***Melanella afra* var. *zonata* Bourguignat, 1884**

Original source. Bourguignat 1884: 22.

Type locality. “Aus einem Mühlbache bei Nassenfuss in Unterkrain” [from a mill creek near Mokronog], Slovenia.

Remarks. Introduced for a specimen of *Melania holandri* var. *laevigata* figured in Rossmässler (1839: 38, pl. 50, fig. 666).

***Melanopsis maroccana* var. *zonatosubcostata* Paladilhe, 1875**

Original source. Paladilhe 1875: 94.

Type locality. “Meknès”, Morocco.

Remarks. Described originally as “*zonato-subcostata*”.

***Melanopsis zonites* Gassies, 1870**

Original source. Gassies 1870: 147.

Type locality. “Prope Saint-Vincent” [near Saint-Vincent], New Caledonia.

Remarks. Brot (1879: 457) considered the species as a junior synonym of *M. brevis* Morelet, 1857 (which is a junior homonym of *Melanopsis brevis* Sowerby, 1826; see *M. moreleti* Pallary, 1916).

***Melanopsis praerosa* [sic] var. *zonulata* Paetel, 1888**

Original source. Paetel 1888: 403.

Type locality. “Persia” (Nevill 1884: 208), Iran.

Remarks. Originally introduced as infrasubspecific taxon (“subvariety”) by Nevill (1884), but made available by Paetel (1888) who treated it as variety (Art. 45.5.1). Paetel clearly referred to the description of Nevill.

† *Melanopsis zujovici* Brusina, 1893

Original source. Brusina 1893: 200 (Serbian part), 33 (Italian part), pl. 2, fig. 5.

Type horizon. Pannonian, zone D–E, late Miocene.

Type locality. “Ripanj”, Serbia.

Types. Milan et al. (1974: 99) indicated a holotype, but it is uncertain whether the specimen was the only one Brusina had at hand (holotype by monotypy, Art. 73.1.2). The specimen is stored in the Croatian Natural History Museum, Zagreb, coll. no. 2531-177/1.

Acknowledgements

I am greatly indebted to the FreshGEN team at the Natural History Museum Vienna (NHM Vienna) for support in every respect. Specifically, I would like to thank Andreas Kroh and Mathias Harzhauser for lively and fruitful discussions on nomenclatural problems. The completeness and accuracy of the catalogue benefitted from the knowledge and help of many colleagues, particularly Philippe Bouchet and Jean-Michel Paccard (both at the Muséum National d’Histoire Naturelle, Paris). Furthermore, I would like to thank the following colleagues for help during literature research: Joaquin Albesa (University of València), Ion Andreescu (Bucharest), Vitaliy V. Anistratenko (Schmalhausen Institute of Zoology of NAS, Kiev), Ruud A. Bank (University of Groningen), Daniela Esu (University of Rome), Dilian Georgiev (Plovdiv University, Bulgaria), Elisavet Georgopoulou (NHM Vienna), Ursula Göhlich (NHM Vienna), Gerhard Haszprunar (Bavarian State Collection of Zoology and Ludwig Maximilian University, Munich), Sonja Herzog-Gutsch (NHM Vienna), Olaf Hölzke (State Museum of Natural History Stuttgart), Heinz Kollmann (NHM Vienna), Andrea Kourgli (NHM Vienna), Imre Magyar (MTA-MTM-ELTE Research Group for Paleontology, Budapest), Oleg Mandic (NHM Vienna), Michael W. Rasser (State Museum of Natural History Stuttgart), Maxim Vinarski (Saint-Petersburg State University) and Winfried Werner (Bayerische Staatssammlung für Paläontologie und Geologie, München). Many thanks go to Chen Yanlong (University of Graz) who helped with the translation of the Chinese literature. The constructive reviews by Philippe Bouchet and Abraham Breure are highly appreciated.

The catalogue was assembled during the project “Freshwater systems in the Neogene and Quaternary of Europe: Gastropod biodiversity, provinciality, and faunal gradients” supported by the Austrian Science Fund (FWF project no. P25365-B25).

References

Web-links to the first page of the publications are provided as far as known, mainly referring to the Biodiversity Heritage Library (<http://www.biodiversitylibrary.org/>), Internet Archive (<https://archive.org/>), and to the online catalogues of the Bibliothèque Nationale de France (<http://gallica.bnf.fr/>), the Stadtarchiv Baden (<http://jubilotheque.upmc.fr/>) and the OÖ. Landesmuseum (<http://www.zobodat.at/index.php>).

For some journals and books, the dates given on the cover or the first page of the article do not necessarily reflect the real publication dates. Examples are the *Journal de Conchyliologie* and the *Bulletin de la Société Géologique de France*, which occur many times in the reference list and where the dates deviated considerably (mostly regarding volumes published in the second half of the 19th century). Publication dates of the *Bulletin* were usually supplied in the appendix; for the *Journal de Conchyliologie*, the dates are based on Fischer-Piette (1937).

The special case of Matheron (1842, 1843) should be discussed here briefly because it led to erroneous conclusions on homonymy issues and invalid replacement names. The two references refer to two, slightly different versions of the same publication by Philippe Matheron. The first version was published in the *Répertoire des travaux de la Société de statistique de Marseille*, vol. 6, pp. 81–341, the second as book in May 1843. In a note at the end of the book (p. 269), Matheron stated: “Circumstances beyond my control have delayed the release of the discussion above, the work that I presented at the Statistical Society of Marseille at the beginning of the year 1842 and part of which was already printed in August of the same year. The time that has elapsed since the original wording of the catalogue until the moment I write this note (in May 1843) allowed me to mention many species recently published by M. d’Orbigny and make my job more complete than it would have been if it had been published a year ago.” Hence, we deal with two different publications that need to be cited separately. The melanopsids in question already appeared in the 1842-version and no changes were made for the 1843-version.

Adams H, Adams A (1853–1858) The genera of Recent Mollusca arranged according to their organizations. Van Voorst, London, 2 vol. of text (661 pp.), 1 vol. of plates pp. <http://www.biodiversitylibrary.org/bibliography/4772#/summary> (for publication dates see Bouchet and Rocroi 2005: 285).

Agassiz L (1847) *Nomenclatoris zoologici index universalis, continens nomina systematica classium, ordinum, familiarum et generum animalium omnium, tam viventium quam fossilium, secundum ordinem alphabeticum unicum disposita, adjectis homonymiis plantarum*. Soloduri, Jent et Gassmann, 393 pp. <http://www.biodiversitylibrary.org/item/16933#page/13/mode/1up> [re-issued version from 1848]

Ahuir Galindo J (2014a) New *Melanopsis* from Morocco. *Malacologia Mostra Mondiale* 84: 14–15.

Ahuir Galindo J (2014b) New *Melanopsis* from Morocco. *Malacologia Mostra Mondiale* 85: 7–8.

- Ahuir Galindo J (2015) A new subspecies of *Melanopsis tricarinata* (Bruguière, 1789) from Spain. *Malacologia Mostra Mondiale* 86: 22–23.
- Ahuir Galindo J (2016) New *Melanopsis* from Morocco. *Malacologia Mostra Mondiale* 90: 26–28.
- Aldrich TH (1886) Preliminary Report on the Tertiary Fossils of Alabama and Mississippi. Geological Survey of Alabama, Bulletin 1: 15–60. <http://www.biodiversitylibrary.org/item/67748#page/19/mode/1up>
- Aldrich TH (1895) New or little known Tertiary Mollusca from Alabama and Texas. *Bulletins of American Paleontology* 1 (2): 53–82. <http://www.biodiversitylibrary.org/item/96050#page/67/mode/1up>
- Allmon WD (1990) Review of the *Bullia* Group (Gastropoda: Nassariidae) with comments on its Evolution, Biogeography, and Phylogeny. *Bulletins of American Paleontology* 99 (335): 1–179. <http://www.biodiversitylibrary.org/item/40609#page/7/mode/1up>
- Almera J (1894) Descripción de los depósitos pliocénicos de la cuenca del Bajo Llobregat y llano de Barcelona. *Memorias de la Real Academia de Ciencias y Artes de Barcelona* 3: 1–355.
- Altaba CR (1998) Testing vicariance: melanopsid snails and Neogene tectonics in the western Mediterranean. *Journal of Biogeography* 25 (3): 541–551. doi: 10.1046/j.1365-2699.1998.2530541.x
- Anderson H-J (1964) Die miocene Reinbek-Stufe in Nord- und Westdeutschland und ihre Mollusken-Fauna. In: Hesemann J (Ed.) *Faunen aus dem Miocän Nordwestdeutschlands. Fortschritte in der Geologie von Rheinland und Westfalen* 17: 31–368.
- Andrae A (1893) [Book review of] P. Oppenheim: *Beiträge zur Kenntniss des Neogen in Griechenland. Mit einer geologischen Einleitung von A. Philippson. Mit 3 Taf. (Zeitschr. d. deutsch. geol. Ges. Bd. XLIII. 421. 1891.)*. *Neues Jahrbuch für Mineralogie, Geologie und Paläontologie* 2: 170–172. <http://www.biodiversitylibrary.org/item/151140#page/400/mode/1up>
- Andrusov N (1909) Materialy k poznaniyu Prikaspiyskogo neogena. *Ponticheskiye plasty She-makhinskogo uyezda. Trudy Geologicheskogo Komiteta, novaya seriya* 40: 1–177.
- Angelov AM (2000) *Catalogus Faunae Bulgaricae, Volume 4. Mollusca: Gastropoda et Bivalvia aquae dulcis*. Pensoft Publishers, Sofia, 57 pp. Parts of the work are available from https://books.google.at/books?id=m95hcMSai2cC&printsec=frontcover&hl=de&source=gbs_ge_summary_r&cad=0#v=onepage&q&f=false
- Anistratenko VV (1993) Mollyuski semeystv Melanopsidae i Bithyniidae (Gastropoda, Pectinibranchia) iz kimmeriyskikh otlozheniy Abkhazii. *Byulleten' Moskovskogo Obshchestva Ispytateley Prirody, Otdel biologicheskii* 98 (5): 68–79.
- Anistratenko VV, Anistratenko OYu (2001) *Fauna Ukraini* 29, Mollyuski 1(1): Class Polyplacophora or Chitons, class Gastropoda - Cyclobranchia, Scutibranchia and Pectinibranchia. *Natsional'na akademiia nauk Ukraïny, Instytut Zoologii, Kiev*, 240 pp. [not seen]
- Annandale N (1918) Freshwater shells from Mesopotamia. *Records of the Indian Museum* 15: 159–170. <http://www.biodiversitylibrary.org/item/41750#page/227/mode/1up>
- Annandale N, Prashad B (1919) The Mollusca of the inland waters of Baluchistan and of Seistan. With a note on the Liver-Fluke of Sheep in Seistan, by S. W. Kemp. *Records of*

- the Indian Museum 18: 17–63. <http://www.biodiversitylibrary.org/item/41762#page/45/mode/1up>
- Azpeitia Moros F (1929) Monografía de las *Melanopsis* vivientes y fósiles de España. Memorias del Instituto Geológico y Minero de España 36: 1–402.
- Bănărescu PM (1990) Zoogeography of Fresh Waters. Vol.1. General distribution and dispersal of freshwater animals. Aula-Verlag, Wiesbaden, 512 pp.
- Bandel K (2000) Speciation among the Melanopsidae (Caenogastropoda). Special emphasis to the Melanopsidae of the Pannonian Lake at Pontian time (Late Miocene) and the Pleistocene and Recent of Jordan. Mitteilungen aus dem Geologisch-Paläontologischen Institut der Universität Hamburg 84: 131–208.
- Bandel K, Riedel F (1994) The late Cretaceous gastropod fauna from Ajka (Bakony Mountains, Hungary): a revision. Annalen des Naturhistorischen Museums in Wien, Serie A 96: 1–65. http://www.zobodat.at/pdf/ANNA_96A_0001-0065.pdf
- Bartha F (1955) A várpalotai pliocén puhatestű fauna biosztratigrafiai vizsgálata. A Magyar Állami Földtani Intézet Évkönyve 43 (2): 275–359.
- Bellardi L, Michelotti G (1841) Saggio orittografico sulla classe dei Gasteropodi del Piemonte. Memorie della Reale Accademia delle Scienze di Torino, Serie II 3: 93–174.
- Beu AG, Maxwell PA (1990) Cenozoic Mollusca of New Zealand. New Zealand Geological Survey Paleontological Bulletin 58: 1–518. http://www.gns.cri.nz/static/catalog/p_bull_58.pdf
- Biggs HEJ (1934) Supposed new variety of *Melanopsis doriae* Issel. Journal of Conchology 20 (2): 60.
- Biggs HEJ (1937) Mollusca of the Iranian Plateau. Journal of Conchology 20 (12): 342–350.
- Biggs HEJ (1962) Mollusca of the Iranian Plateau-II. Journal of Conchology 25 (2): 67–72.
- Blanckenhorn M (1897) Zur Kenntnis der Süßwasserablagerungen und Mollusken Syriens. Palaeontographica 44: 71–144. <http://www.biodiversitylibrary.org/item/103439#page/81/mode/1up>
- Blanckenhorn M (1901) Neues zur Geologie und Paläontologie Aegyptens. IV. Das Pliocän und Quartärzeitalter in Aegypten ausschliesslich des Rothen Meergebietes. Zeitschrift der Deutschen Geologischen Gesellschaft 53 (3): 307–502. <http://www.biodiversitylibrary.org/item/150066#page/387/mode/1up>
- Blanckenhorn M, Oppenheim P (1927) Neue Beiträge zur Kenntnis des Neogens in Syrien und Palästina. Geologische und Palaeontologische Abhandlungen, Neue Folge 15 (4): 319–358.
- Bogachev V (1908) Verkhnemiotzenoviya unionidy Kavkaza. Izvestiya Geologicheskogo Komiteta 27 (4): 237–258.
- Bogachev VV (1938) Miotsen Zakavkaz'ya (V svete novykh dannykh 1932-1935 g.) [The Transcaucasian Miocene (in the light of new data from the years 1932-1935)]. Trudy Azerbaydzhanskogo filiala Akademii Nauk SSSR. Geologicheskaya seriya 10 (44): 1–67. [In Azerbaijan, Russian and German]
- Boistel A (1898) Nouvelles observations sur la bordure tertiaire du Jura a l'ouest du Bugey. Bulletin de la Société Géologique de France, troisième série 26: 11–35. http://jubilothèque.upmc.fr/img-viewer/fonds-bulsgf/GB_000072_001/Contenu/JPEG_HD/viewer.html?ns=GB_000072_001_J3_0013.jpg

- Bonelli A (1827) Denominaciones ineditae testaceorum Musei Zoologici Taurinensis. Unpublished catalogue, Torino.
- Botez G (1914) Ueber die levantinische Molluskenfauna von Moreni. Bulletin de la section scientifique de l'Academie Roumaine 2 (8): 235–241.
- Bouchet P, Poppe GT (1995) A review of the deep-water volute genus *Calliotectum* (Gastropoda: Volutidae). In: Bouchet P (Ed.) Résultats des Campagnes Musorstom, volume 14. Mémoires du Muséum national d'Histoire naturelle 167: 499–525. <http://www.vliz.be/nl/open-marien-archie?module=ref&refid=215190>
- Bouchet P, Rocroi J-P (2005) Classification and Nomenclator of Gastropod Families. Malacologia 47 (1–2): 1–397. <http://www.biodiversitylibrary.org/page/25127194#page/51mode/1up>
- Bourguignat JR (1853) Catalogue raisonné des mollusques terrestres et fluviatiles recueillis par M. F. de Saulcy pendant son voyage en Orient. Gide et J. Baudry, Paris, xxvi + 96 pp. <http://gallica.bnf.fr/ark:/12148/bpt6k99038p/f1.item>
- Bourguignat JR (1862) Paléontologie des mollusques terrestres et fluviatiles de l'Algérie. Baillière et fils, Paris, 126 pp. [https://books.google.at/books?id=k2hw4aoCjAMC&printsec=f](https://books.google.at/books?id=k2hw4aoCjAMC&printsec=frontcover&hl=de#v=onepage&q&f=false)
- Bourguignat JR (1864) Malacologie de l'Algérie, ou, Histoire naturelle des animaux mollusques terrestres et fluviatiles recueillis jusqu'à ce jour dans nos possessions du nord de l'Afrique. Tome second. Challamel Ainé, Paris, 380 pp. <http://www.biodiversitylibrary.org/item/106363#page/9/mode/1up>
- Bourguignat JR (1868) Mollusques nouveaux, litigieux ou peu connus. Revue et magasin de zoologie pure et appliquée, deuxième série 20: 422–433. <http://www.biodiversitylibrary.org/item/20627#page/428/mode/1up>
- Bourguignat JR (1870) Aperçu sur la faune malacologique du Bas Danube. Annales de Malacologie 1: 36–76. <http://www.biodiversitylibrary.org/item/54956#page/48/mode/1up>
- Bourguignat JR (1876) Species novissimae Molluscorum in Europaeo systemati detectae, notis diagnosticis succinctis breviter descriptae. Paul Klincksieck, Paris, 80 pp. doi: 10.5962/bhl.title.10357
- Bourguignat JR (1877) Descriptions de deux nouveaux genres algériens, suivies d'une classification des familles et des genres de mollusques terrestres et fluviatiles du système européen. Bulletin de la Société des Sciences Physiques et Naturelles de Toulouse 3 (1): 49–101. <http://www.animalbase.uni-goettingen.de/zooweb/servlet/AnimalBase/home/digireference?id=43>
- Bourguignat JR (1880) Étude sur les fossiles tertiaires et quaternaires de la vallée de la Cettina en Dalmatie. Imprimerie D. Bardin, Saint Germain, 55 pp.
- Bourguignat JR (1884) Histoire des mélaniens du système européen. Annales de Malacologie 2: 1–168. <http://www.biodiversitylibrary.org/item/55067#page/9/mode/1up>
- Bowdich TE (1822) Elements of Conchology, including the fossil genera and the animals. Part I. Univalves. Treuttel and Würtz, London, 79 pp. + 19 pls. <http://www.biodiversitylibrary.org/item/45861#page/5/mode/1up>
- Braun A (1851) Darstellung der geognostischen Verhältnisse des Mainzer Beckens und seiner fossilen Fauna und Flora. In: Walchner FA (Ed.) Handbuch der Geognosie zum Gebrauche

- bei seinen Vorlesungen und zum Selbststudium mit besonderer Berücksichtigung der geognostischen Verhältnisse des Grossherzogthums Baden. Christian Theodor Gross, Karlsruhe, 1121–1169. https://books.google.at/books?id=7h0-AAAAcAAJ&printsec=frontcover&hl=de&source=gbs_ge_summary_r&cad=0#v=onepage&q&f=false
- Briart A, Cornet FL (1873) Description des fossiles du calcaire grossier de Mons. Seconde partie. Gastéropodes. Ordre I. - Prosobranches. Section B. - Syphonostomes (première partie). Mémoires couronnés et Mémoires des savants étrangers, publiés par l'Académie Royale des Sciences, des Lettres et des Beaux-arts de Belgique 37: iv + 1–94. <http://www.biodiversitylibrary.org/item/28722#page/363/mode/1up>
- Briart A, Cornet FL (1882) Description des fossiles du calcaire grossier de Mons. Troisième partie. Supplément aux deux premières parties. Mémoires couronnés et Mémoires des savants étrangers, publiés par l'Académie Royale des Sciences, des Lettres et des Beaux-arts de Belgique 43: viii + 1–73. http://gdz.sub.uni-goettingen.de/dms/load/img/?PPN=PPN129323659_0043&DMDID=DMDLOG_0006
- Bronn HG (1848) Index palaeontologicus, oder Uebersicht der bis jetzt bekannten fossilen Organismen. Erste Abtheilung. A. Nomenclator palaeontologicus, in alphabetischer Ordnung. Erste Hälfte. A-M. Schweizerbart'sche Verlagshandlung und Druckerei, Stuttgart, 775 pp.
- Brot A (1862) Matériaux pour servir à l'étude de la famille des mélaniens. Catalogue systématique des espèces qui composent la famille des mélaniens, I. Jules-Guillaume Fick, Genève, 72 pp. <https://archive.org/stream/matriaupourse113brot#page/n9/mode/2up>
- Brot A (1868) Matériaux pour servir à l'étude de la famille des mélaniens. Catalogue systématique des espèces qui composent la famille des mélaniens, II. Georg, Genève, 55 pp. <https://archive.org/stream/matriaupourse113brot#page/n85/mode/2up>
- Brot A (1870) Catalogue of the recent species of the family Melanidae. American Journal of Conchology 6: 271–325. <http://www.biodiversitylibrary.org/item/31241#page/261/mode/1up>
- Brot A (1874–1879) Die Melaniaceen (Melanidae) in Abbildungen nach der Natur mit Beschreibungen. - Systematisches Conchylien-Cabinet von Martini und Chemnitz 1 (24). Bauer & Raspe, Nürnberg, pp. 1–32, pls 1–6 (1874), pp. 33–125, pls 7–18 (1875), pp. 129–192, pls 19–24 (1876), pp. 193–352, pls 25–36 (1877), pp. 353–400, pls 37–42 (1878), pp. 401–488, pls 43–49 (1879). <http://www.biodiversitylibrary.org/item/106973#page/9/mode/1up>
- Bruguière JG (1789) Encyclopedie methodique. Histoire naturelle des Vers. Tome sixième. Panckoucke & Plomteux, Paris & Liege, 344 pp. https://books.google.at/books?id=VFQ18ndarFYC&printsec=frontcover&hl=de&source=gbs_ge_summary_r&cad=0#v=onepage&q&f=false
- Bruguière JG (1791) Tableau encyclopédique et méthodique des trois règnes de la nature. Contenant l'helminthologie, ou les vers infusoires, les vers intestins, les vers mollusques, &c. Septième livraison. Panckoucke, Paris, 83 pp. + pl. 1–92. <http://biodiversitylibrary.org/item/130182#page/9/mode/1up>
- Brusina S (1866) Contribuzione pella fauna dei molluschi dalmati. Verhandlungen der kaiserlichen und königlichen zoologisch-botanischen Gesellschaft in Wien 16 (Beigabe zu Heft 1/2): iii + 1–134. <http://www.biodiversitylibrary.org/item/41695#page/7/mode/1up>

- Brusina S (1867) Prinesci malakologiji hrvatskoj. Rad Jugoslavenske akademije znanosti i umjetnosti 1: 78–105. [in Croatian] https://books.google.at/books?id=ciVJAAAAcAAJ&printsec=frontcover&hl=de&source=gbs_ge_summary_r&cad=0#v=onepage&q&f=false
- Brusina S (1870) Contribution à la malacologie de la Croatie. Ch. Albrecht, Zagreb (Agram), 40 pp. <http://www.biodiversitylibrary.org/item/106124#page/1/mode/1up>
- Brusina S (1874) Fossile Binnen-Mollusken aus Dalmatien, Kroatien und Slavonien nebst einem Anhang. Actienbuchdruckerei, Agram, 138 pp.
- Brusina S (1876) Description d'espèces nouvelles, provenant des terrains tertiaires de Dalmatie. Journal de Conchyliologie 24: 109–116. <http://www.biodiversitylibrary.org/item/54517#page/119/mode/1up>
- Brusina S (1878) Molluscorum fossilium species novae et emendatae in tellure tertiaria Dalmatiae, Croatiae et Slavoniae inventae. Journal de Conchyliologie 26: 347–356. <http://www.biodiversitylibrary.org/item/54320#page/363/mode/1up>
- Brusina S (1884a) Die Fauna der Congerienschichten von Agram in Kroatien. Beiträge zur Paläontologie und Geologie Österreich-Ungarns und des Orients 3: 125–187.
- Brusina S (1884b) Die *Neritodonta* Dalmatiens und Slavoniens nebst allerlei malakologischen Bemerkungen. Jahrbücher der Deutschen Malakozoologischen Gesellschaft 11: 17–120. <http://www.biodiversitylibrary.org/item/55039#page/31/mode/1up>
- Brusina S (1885) Bemerkungen über rumänische Paludinenschichten mit Bezug auf Prof. Cobalcescu's Werk "Studii geologice si paleontologice asupra unor Teramuri Tertiare din unile Parti ale Romaniei" (Mem. Geol. ale Scolei militare din Iasi, Mem. I, Bucuresti 1883). Verhandlungen der k.k. geologischen Reichsanstalt 6: 157–163. <http://www.biodiversitylibrary.org/item/120934#page/165/mode/1up>
- Brusina S (1892) Fauna fossile terziaria di Markuševac in Croazia. Con un elenco delle Dreissensidae della Dalmazia, Croazia e Slavonia. Glasnik Hrvatskoga naravoslovnoga društva 7: 113–210. <http://www.biodiversitylibrary.org/item/42638#page/131/mode/1up>
- Brusina S (1893) Odlmoci srpske tercijerne malakologije/Frammenti di Malacologia terziaria Serba. Annales Géologiques de la Péninsule Balkanique 4: 192–208 (Serbian part), 25–74 (Italian part, which contained much more text and was separately numbered but simultaneously published).
- Brusina S (1897) Gragja za neogensku malakološku faunu Dalmacije, Hrvatske i Slavonije uz neke vrste iz Bosne i Hercegovine i Srbije. Djela Jugoslavenske akademije znanosti i umjetnosti 18: 1–43. [in Croatian and French] <http://archive.org/stream/djelajugoslaven03umjegoog#page/n6/mode/2up>
- Brusina S (1902) Iconographia Molluscorum Fossilium in tellure tertiaria Hungariae, Croatiae, Slavoniae, Dalmatiae, Bosniae, Herzegovinae, Serbiae and Bulgariae inventorum. Officina Soc. Typographicae, Agram, 30 pls.
- Brusina S (1903) Eine subtropische Oasis in Ungarn. Mitteilungen des naturwissenschaftlichen Vereines für Steiermark 39: 101–121. <http://www.biodiversitylibrary.org/item/45396#page/203/mode/1up>
- Brusina S (1904) Šipovo i njegova tercijarna fauna. Glasnik Zemaljskog muzeja Bosne i Hercegovine 16 (4): 493–498. [in Bosnian] The German version published in Wissenschaftliche

- Mitteilungen aus Bosnien und der Herzegowina in 1909 is available from <http://www.biodiversitylibrary.org/item/192826#page/594/mode/1up>
- Brusina S (1905) Naravoslovne crtice sa sjevero-istočne obale Jadranskoga mora. Dio treći. Putopis. Rad Jugoslavenske akademije znanosti i umjetnosti 163: 1–40. [in Croatian]
- Brusina S (1907) Naravoslovne crtice sa sjevero-istočne obale Jadranskoga mora. Dio četvrti i posljednji. Specijalni. Rad Jugoslavenske akademije znanosti i umjetnosti 169: 195–251. [in Croatian]
- Bukowski G von (1892) Notiz über die Molluskenfauna der levantinischen Bildungen der Insel Rhodus. Anzeiger der Österreichischen Akademie der Wissenschaften, mathematisch-naturwissenschaftliche Classe 29: 247–250.
- Bukowski G von (1893) Die levantinische Molluskenfauna der Insel Rhodus, I. Theil. Denkschriften der mathematisch-naturwissenschaftlichen Classe der Kaiserlichen Akademie der Wissenschaften 60: 265–306. <http://www.biodiversitylibrary.org/item/31609#page/283/mode/1up>
- Burgerstein L (1877) Beitrag zur Kenntniss des Jungtertiären Süßwasser-Depôts bei Ueskueb. Jahrbuch der k. k. geologischen Reichsanstalt 27: 243–250. <http://www.biodiversitylibrary.org/item/110135#page/261/mode/1up>
- Calvert F, Neumayr M (1880) Die jungen Ablagerungen am Hellespont. Denkschriften der mathematisch-naturwissenschaftlichen Classe der Kaiserlichen Akademie der Wissenschaften 40: 357–378. <http://www.biodiversitylibrary.org/item/104344#page/395/mode/1up>
- Calzada S, Urquiola M de M (1994) Nuevos datos sobre *Melanopsis armatus* Mathéron, 1843. (Gasterópodo del Cretácico, facies Garumniense). Coloquios de Paleontología 46: 151–159.
- Capellini G (1873) La formazione gessosa di Castellina Marittima e i suoi fossili. Memorie della Reale Accademia delle Scienze dell'Istituto di Bologna, Ser. 3: 525–602. <http://www.biodiversitylibrary.org/item/120571#page/629/mode/1up>
- Capellini G (1880) Gli strati a Congerie e la formazione gessoso-solfifera nella provincia di Pisa e nei dintorni di Livorno. Atti della Reale Accademia dei Lincei, Ann. 277, Memorie, Ser. 3 5: 375–427.
- Carez L (1879) Description d'espèces nouvelle des terrains tertiaires du bassin de Paris. Bulletin de la Société Géologique de France, troisième série 7: 637–641. http://jubilotheque.upmc.fr/img-viewer/fonds-bulsgf/GB_000051_001/Contenu/JPEG_HD/viewer.html?ns=GB_000051_001_J3_0648.jpg
- Cerulli-Irelli S (1914) Fauna malacologica mariana. Parte settima. Melaniidae, Littorinidae, Fossaridae, Solariidae, Rissoidae, Capulidae, Calyptraeidae, Xenophoridae, Lamellariidae, Naticidae, Scalidae, Aclisidae, Eulimidae, Pyramidellidae. Palaeontographica Italica 20: 183–277. <http://www.biodiversitylibrary.org/item/104144#page/191/mode/1up>
- Chemnitz JH (1795) Neues systematisches Conchylien-Cabinet. Eilfter Band. Gabriel Nikolaus Raspe, Nürnberg, 310 pp. <http://www.biodiversitylibrary.org/item/94992#page/5/mode/1up>
- Chenu JC (1859) Manuel de conchyliologie et de paléontologie conchyliologique. Tome premier. Victor Masson, Paris, 508 pp. doi: 10.5962/bhl.title.10591
- Chevallier H (1969) Mollusques subfossiles récoltés par M. Henri Lhote dans le Sud Oranais et le Sahara. Bulletin du Muséum National d'Histoire Naturelle, deuxième série 41 (1): 266–294.

- Clessin S (1887–1890) Die Mollusken-Fauna Mitteleuropa's. II. Theil. Die Molluskenfauna Oesterreich-Ungarns und der Schweiz. Bauer & Raspe, Nürnberg, 1–160 pp. (1887), 161–320 pp. (1887), 321–480 pp. (1888), 481–624 pp. (1889), i–ii, 625–858 pp. (1890). <http://biodiversitylibrary.org/item/122836#page/8/mode/1up>
- Cobălcescu G (1883) Studii geologice și paleontologice asupra unor tărâmurii terțiare din unele părți ale României. Memoriile Geologice ale Școlii Militare din Iași 1: 1–161.
- Coelho AR, Dinis MT, Reis J (2013) Effect of Diet and Stocking Densities on Life History Traits of *Clea helenae* (Philippi 1847) Reared in Captivity. Journal of Aquaculture & Research Development 4 (5): 187: 4 pp. doi: 10.4172/2155-9546.1000187
- Conrad TA (1830) On the Geology and Organic Remains of a part of the Peninsula of Maryland. Journal of the Academy of Natural Sciences of Philadelphia 6: 205–230. <http://www.biodiversitylibrary.org/item/79409#page/217/mode/1up>
- Conrad TA (1860) Description of New species of Cretaceous and Eocene Fossils of Mississippi and Alabama. Journal of the Academy of Natural Sciences of Philadelphia, second series 4: 275–298. <http://www.biodiversitylibrary.org/item/113513#page/283/mode/1up>
- Conrad TA (1862) Descriptions of new genera, subgenera and species of Tertiary and Recent shells. Proceedings of the Academy of Natural Sciences of Philadelphia 14: 284–291. <http://www.biodiversitylibrary.org/page/1951756>
- Conrad TA (1866) Illustrations of Miocene fossils with descriptions of new species. American Journal of Conchology 2 (1): 65–74. <http://www.biodiversitylibrary.org/item/30582#page/83/mode/1up>
- Cossmann M (1886) Description d'espèces du Terrain tertiaire des environs de Paris (suite). Journal de Conchyliologie 34: 224–235. <http://biodiversitylibrary.org/page/16071119#page/232/mode/1up>
- Cossmann M (1888) Catalogue illustré des coquilles fossiles de l'Éocène des environs de Paris. Annales de la Société royale malacologique de Belgique 23: 3–324. <http://www.biodiversitylibrary.org/item/28336#page/33/mode/1up>
- Cossmann M (1889) Gastéropodes. In: Douvillé H (Ed.) Revue de paléontologie pour l'année 1888. Annuaire Géologique Universel 5: 1079–1109. <https://archive.org/stream/annuairegeologiq09unkngoog#page/n1095/mode/2up>
- Cossmann M (1899) Paléozoologie générale. Revue critique de paléozoologie 3 (4): 141–149. <http://www.biodiversitylibrary.org/item/50630#page/157/mode/1up>
- Cossmann M (1906) Estudio de algunos moluscos eocénicos del Pirineo catalán. Boletín de la Comisión del Mapa Geológico de España 28: 135–151.
- Cossmann M (1907) Appendice No. 4 au Catalogue illustré des coquilles fossiles de l'Éocène des environs de Paris. Annales de la Société royale zoologique et malacologique de Belgique 41: 186–286. <http://www.biodiversitylibrary.org/page/5599125#page/206/mode/1up> [the cover indicates the year 1906, but individual printing sheets are dated January to March 1907]
- Cossmann M (1909) Essais de Paléoconchologie Comparée. Huitième Livraison. Privately published, Paris, 248 pp. <http://www.biodiversitylibrary.org/item/113485#page/337/mode/1up>
- Cossmann M, Peyrot A (1918) Conchologie néogénique de l'Aquitaine. Actes de la Société linnéenne de Bordeaux 70 (3): 5–491. <http://www.biodiversitylibrary.org/item/103369#page/11/mode/1up>

- Cossmann M, Pissarro G (1906–1913) Iconographie complète des coquilles fossiles de l'éocène des environs de Paris. Tome 2e. Scaphopodes, Gastropodes, Brachiopodes, Céphalopodes & Supplément. Privately published, Paris, Fasc. 1: pls 1–9 (1906), Fasc. 2: pls 10–25 (1910), Fasc. 3: pls 26–45 (1911), Fasc. 4: pls 46–65 (1913). <https://archive.org/stream/iconographiecomp212coss#page/n199/mode/2up>
- Crosse H (1867) Description d'espèces inédites provenant de la Nouvelle-Calédonie. *Journal de Conchyliologie* 15 (4): 433–437. <http://biodiversitylibrary.org/item/52768#page/437/mode/1up>
- Crosse H (1869a) Diagnoses Molluscorum novorum. *Journal de Conchyliologie* 17 (1): 69–70. <http://biodiversitylibrary.org/item/53431#page/73/mode/1up>
- Crosse H (1869b) Description d'espèces inédites provenant de la Nouvelle-Calédonie. *Journal de Conchyliologie* 17 (3): 279–281. <http://biodiversitylibrary.org/item/53431#page/283/mode/1up>
- Crosse H (1877) Rapport sur un mémoire de M. A. Letourneux, intitulé: Mollusques terrestres et fluviatiles du Maroc (Bulletin de la Société des sciences physiques, naturelles et climatologiques d'Alger, 13e année, 1876, 4^e trimestre). *Revue des sociétés savantes*, deuxième série 11: 31–33. <http://gallica.bnf.fr/ark:/12148/bpt6k202298p/f33.item>
- Čtyrský P (1972) Die Molluskenfauna der *Rzehakia-* (*Oncophora*)-Schichten Mährens. *Annalen des Naturhistorischen Museums in Wien* 76: 41–141. http://www.zobodat.at/pdf/ANNA_76_0041-0141.pdf
- Dautzenberg P (1894) Liste des mollusques terrestres et fluviatiles recueillis par M. Th. Barrois en Palestine et en Syrie. *Revue biologique du nord de la France* 6: 329–353. <http://www.biodiversitylibrary.org/item/41024#page/339/mode/1up>
- De Cristofori G, Jan G (1832) *Catalogus in IV. sectiones divisus rerum naturalium in museo exstantium Josephi De Cristofori et Georgii Jan plurium Acad. Scient. et Societ. Nat. Cur. Sodalium complectens adumbrationem oryctognosiae et geognosiae atque prodromum faunae et floriae Italiae Superioris. Sectio II. Conchyliologia. Pars I. Conspectus methodicus molluscorum. Fasc. 1. Testacea terrestria et fluviatilia. Carmignani, Parma, 1–2 pp. [unnumbered] [Dispositio methodica generum], pp. 1–8 [Conchyilia terrestria et fluviatilia], pp. 1–4 [Mantissa in secundam partem catalogi testaceorum exstantium in collectione quam possident], pp. 1–4, [Excerptum e primo nostre programme], pp. 1–16 [Conchyilia fossilia].* <http://www.biodiversitylibrary.org/item/122163#page/5/mode/1up>
- Deffner C, Fraas O (1877) Begleitworte zur geognostischen Spezialkarte von Württemberg. *Atlasblätter Bopfingen und Ellenberg*. W. Kohlhammer, Stuttgart, 36 pp.
- De Gregorio A (1880) *Fauna di S. Giovanni Ilarione. Fasc. I. P. Montaina e C., Palermo, xxviii + 106 pp.*
- De Laubrière L, Carez L (1881) Sur les Sables de Brasles (Aisne). *Bulletin de la Société Géologique de France*, troisième série 8: 391–413. http://jubilothèque.upmc.fr/img-viewer/fonds-bulsgf/GB_000052_001/Contenu/JPEG_HD/viewer.html?ns=GB_000052_001_J3_471.jpg
- Delpéy G (1940) Les gastéropodes mésozoïques de la région libanaise. *Haut Commissariat de la République française en Syrie et au Liban, Service des travaux publics, Section d'études géologiques - Notes et mémoires* 3: 5–292.

- Deshayes G-P (1824–1837) Description des coquilles fossiles des environs de Paris, vol. II. Privately published, Paris, 1–80 pp. (1824), 81–146 pp. (1825), 147–290 pp. (1832), 291–426 pp. (1833), 427–498 pp. (1834), 499–780 pp. (1835), 781–814 pp. (1837). <http://www.biodiversitylibrary.org/item/112044#page/6/mode/1up> (text); <http://www.biodiversitylibrary.org/item/111960#page/5/mode/1up> (atlas)
- Deshayes GP (1830–1832) Encyclopédie méthodique ou par ordre de matières. Histoire naturelle des Vers et Mollusques. Privately published, Paris. Vol. 2, part 1: i–vi, 1–256 pp. (1830); part 2: 1–144 pp. (1830), 145–594 pp. (1832). Vol. 3: 595–1152 pp. (1832). <http://www.biodiversitylibrary.org/bibliography/49857>
- Deshayes GP (1861–1864) Description des animaux sans vertèbres découverts dans le bassin de Paris pour servir de supplément a la description des coquilles fossiles des environs de Paris comprenant une revue générale de toutes les espèces actuellement connues. Tome deuxième. - Texte. Mollusques Acéphalés Monomyaires et Brachiopodes. Mollusques céphalés. Première Partie. J.-B. Baillière et Fils, Paris, pp. 1–432, pls 1–26 (1861), pp. 433–640, pls 27–39 (1862), pp. 641–920, pls 40–62 (1863), pp. 921–968 (1864). <http://www.biodiversitylibrary.org/item/110230#page/9/mode/1up>
- Deshayes GP, Jullien J (1876) Mémoire sur les mollusques nouveaux du Cambodge envoyés au Muséum par M. le Docteur Jullien. Bulletin des Nouvelles Archives du Museum d'histoire naturelle 10 (1874): 115–162. <http://www.biodiversitylibrary.org/item/49424#page/381/mode/1up>
- De Serres M (1829) Géognosie des terrains tertiaires ou tableau des principaux animaux invertébrés des terrains marins tertiaires du midi de la France. Pomathio-Durville, Montpellier, Paris, 276 pp. https://books.google.at/books?id=e3EOAAAAQAAJ&pg=PA101&lpg=PA101&dq=%22melanopsis+deperdita%22+serres&source=bl&ots=sQALAl0_Dw&sig=gLGpFXBcKKjDj2c7BaoB0wrEB0M&hl=de&sa=X&ved=0ahUKEwjaw6uO1NPJAhUH_iwKHclpCGAQ6AEIjAA#v=onepage&q&f=false
- De Stefani C (1877) Molluschi continentali fino ad ora notati in Italia nei terreni pliocenici, ed ordinamento di questi ultimi. Atti della Società Toscana di Scienze Naturali, residente in Pisa 3 (2): 274–325. <http://www.biodiversitylibrary.org/item/108494#page/316/mode/1up>
- De Stefani C (1880) Molluschi continentali fino ad ora notati in Italia nei terreni pliocenici, ed ordinamento di questi ultimi (continuazione). Atti della Società Toscana di Scienze Naturali, residente in Pisa 5 (1): 9–108. <http://www.biodiversitylibrary.org/item/113405#page/13/mode/1up>
- Doderlein P (1863) Cenni geologici intorno la giacitura dei terreni miocenici superiori dell'Italia centrale. Atti del 10° Congresso degli scienziati Italiani 1862: 1–25.
- Dollfus GF (1906) Feuille de Bourges au 320 000e (Révision des Faunes continentales). Bulletin des services de la carte géologique de la France. Comptes rendus des collaborateurs pour la campagne de 1905 16 (110): 284–304.
- Dollfus GF (1912) Recherches critiques sur quelques genre et espèces d'*Hydrobia* vivants ou fossiles. Journal de Conchyliologie 59 (3): 179–270. <http://biodiversitylibrary.org/item/55064#page/207/mode/1up>
- Dollfus GF (1922) Faune malacologique du Miocène supérieur de Janina en Epire. Bulletin de la Société Géologique de France, quatrième série 22: 101–123. <http://jubilo->

- theque.upmc.fr/img-viewer/fonds-bulsgf/GB_000097_001/Contenu/JPEG_HD/viewer.html?ns=GB_000097_001_J3_128.jpg
- Dominici S, Kowalke T (2014) Early Eocene Cerithioidean gastropods of mangrove-fringed coasts (South-central Pyrenees, Spain). *Bollettino della Società Paleontologica Italiana* 53 (3): 137–162. doi: 10.4435/BSPI.2014.13
- Doncieux L (1908) Catalogue descriptif des fossiles nummulitiques de l'Aude et de l'Hérault. Deuxième partie (Fascicule I). Corbières septentrionale. *Annales de l'Université de Lyon, Nouvelle Série, I. Sciences, Médecine* 22: 1–288.
- D'Orbigny A (1850) *Prodrome de Paléontologie. Stratigraphique universelle des animaux mollusques et rayonnés faisant suite au cours élémentaire de paléontologie et de géologie stratigraphique. Deuxième volume.* Victor Masson, Paris, 427 pp. doi: 10.5962/bhl.title.62810
- D'Orbigny A (1852) *Prodrome de Paléontologie. Stratigraphique universelle des animaux mollusques et rayonnés faisant suite au cours élémentaire de paléontologie et de géologie stratigraphique. Troisième volume.* Victor Masson, Paris, 189 pp. <http://www.biodiversitylibrary.org/item/98989#page/9/mode/1up>
- Douvillé H (1904) *Mission scientifique en Perse. Tome troisième. Etudes géologiques. Partie IV. - Paléontologie. Mollusques fossiles.* Ernest Leroux, Paris, 191–368 pp. <http://gallica.bnf.fr/ark:/12148/bpt6k110553t/f300.item>
- Drensky P (1947) *Sustav i razprostranenie na sladkovodnite Mollusca v Bulgariya* [Synopsis and distribution of freshwater Mollusca in Bulgaria]. *Godishnik na Sofiiskiya Universitet, Fiziko-matematicheski fakultet, Kniga 3 - Estestrenna istoriya* 43 (3): 33–54. [In Bulgarian with English summary]
- Dunker W (1848) Über die in der Molasse bei Günzberg unfern Ulm vorkommenden Conchylien und Pflanzenreste. *Palaeontographica* 1: 155–168. <http://www.biodiversitylibrary.org/item/43708#page/171/mode/1up>
- Dunker W (1862) Beschreibung einiger von Herrn Dr. v. Hochstetter auf Neuseeland gesammelten Süßwasser-Mollusken. *Malakozoologische Blätter* 8: 150–154. <http://www.biodiversitylibrary.org/item/54069#page/418/mode/1up>
- Erber J (1868) Bericht über eine Reise nach Rhodus. *Verhandlungen der kaiserlichen und königlichen zoologisch-botanischen Gesellschaft in Wien* 18: 903–908. <http://www.biodiversitylibrary.org/item/81414#page/1081/mode/1up>
- Esu D, Girotti O (1975) La malacofauna continentale del Plio-Pleistocene dell'Italia centrale. I. Paleontologia. *Geologica Romana* 13: 203–294. http://www.dst.uniroma1.it/geologica-romana/Volumi/VOL%2013/GR_13_203_293_Esu%20et%20al.pdf
- Esu D, Girotti O (2015a) The late Early Pleistocene non-marine molluscan fauna from the Synania Formation (Achaia, Greece), with description of nine new species. *Archiv für Molluskenkunde* 144 (1): 65–81. doi: 10.1127/arch.moll/1869-0963/144/065-081
- Esu D, Girotti O (2015b) *Melanopsis wilhelmi* n. sp. and *Valvata ducati* n. sp., two new Pleistocene gastropods from a section of the Stirone River (Emilia, North Italy) (Gastropoda: Melanopsidae, Valvatidae). *Archiv für Molluskenkunde* 144 (2): 149–154. doi: 10.1127/arch.moll/1869-0963/144/149-154

- Etheridge R (1879) Notes on the Mollusca collected by C. Barrington Brown, Esq., A.R.S.M., from the Tertiary Deposits of Solimões and Javary Rivers, Brazil. *Quarterly Journal of the Geological Society* 35: 82–88. doi: 10.1144/GSL.JGS.1879.035.01-04.126
- Falsan A, Locard A (1879) Note sur les formations tertiaires et quaternaires des environs de Miribel (Ain). *Annales de la Société d'Agriculture. Histoire naturelle et arts utiles de Lyon*, cinquième série 1: 1–10.
- Férussac AEJPJFd'Audebard de (1812) Sur des Terres d'eau douce observés en divers lieux, et sur les Fossiles terrestres et fluviatiles. *Annales du Muséum National d'Histoire Naturelle* 19: 242–256. <http://www.biodiversitylibrary.org/item/23270#page/276/mode/1up>
- Férussac AEJPJFd'Audebard de (1814) Mémoires géologiques sur les terrains formés sous l'eau douce par les débris fossiles des mollusques vivant sur la terre ou dans l'eau non salée. Poulet, Paris, 76 pp. <http://books.google.at/books?pg=PR5&lpg=PA17&id=NqpAAAAAcAAJ&hl=de&ots=x7ErJq1S5a#v=onepage&q&f=false>
- Férussac AEJPJFd'Audebard de [continued by Deshayes GPP] (1819–1851) Histoire naturelle générale et particulière des mollusques terrestres et fluviatiles tant des espèces que l'on trouve aujourd'hui vivantes, que des dépouilles fossiles de celles qui n'existent plus ; classées d'après les caractères essentiels que présentent ces animaux et leurs coquilles. J.-B. Baillière, Paris, 2 vols of text + 210 pls.
- Férussac AEJPJFd'Audebard de (1823) Monographie des espèces vivantes et fossiles du genre mélanopsidé, *Melanopsis*, et observations géologiques à leur sujet. *Mémoires de la Société d'Histoire Naturelle de Paris* 1 (1): 132–164. <http://www.biodiversitylibrary.org/item/23445#page/148/mode/1up>
- Férussac JBLd'Audebard de, Férussac AEJPJFd'Audebard de (1807) Essai d'une méthode conchyliologique appliquée aux mollusques fluviatiles et terrestres d'après la considération de l'animal et de son test. Nouvelle édition augmentée d'une synonymie des espèces les plus remarquables, d'une table de concordance systématique de celles qui ont été décrites par Geoffroy, Poiret et Draparnaud, avec Müller et Linné, et terminée par un catalogue d'espèces observées en divers lieux de la France. Delance, Paris, xvi + 142 pp. http://www.wuser.gwdg.de/~subtypo3/gdz/pdf/PPN614791812/PPN614791812__LOG_0000.pdf
- Finlay HJ (1926) A Further Commentary on New Zealand Molluscan Systematics. *Transactions and Proceedings of the New Zealand Institute* 57: 320–485. http://rsnz.natlib.govt.nz/volume/rsnz_57/rsnz_57_00_003550.pdf
- Fischer-de-Waldheim G (1837) *Oryctographie du Gouvernement de Moscou*. Auguste Semen, Moscou, 202 pp. https://books.google.at/books?id=3F4qT7T4SE4C&printsec=frontcover&hl=de&source=gbg_summary_r&cad=0#v=onepage&q&f=false
- Fischer P (1883) Observations sur la Note précédente. *Journal de Conchyliologie* 31 (1): 60–62. <http://www.biodiversitylibrary.org/item/53857#page/70/mode/1up>
- Fischer P, Crosse H (1880–1902) Études sur les mollusques terrestres et fluviatiles du Mexique et du Guatemala. In: Milne Edwards M (Ed.) *Recherches Zoologiques pour servir à l'histoire de la faune de l'Amérique Centrale et du Mexique*. Septième partie. Tome second. Imprimerie nationale, Paris, 731 pp., pl. 32–72. <http://www.biodiversitylibrary.org/item/100311#page/9/mode/1up> (for publication dates see Bouchet and Rocroi 2005: 306).

- Fischer-Piette E (1937) Dates de publication du “Journal de Conchyliologie” de 1861 à 1900. *Journal de Conchyliologie* 81 (1): 88–92. <http://gallica.bnf.fr/ark:/12148/bpt6k5567298f/f95.image>
- Fischer W (1993) *Melanopsis hennersdorfensis* n. sp., eine neue fossile Melanopside (Gastropoda: Thiaridae) aus dem Wiener Becken (Miocän/Pannon). *Club Conchylia Informationen* 25 (2): 179–183.
- Fischer W (1996a) Beiträge zur Kenntnis der rezenten und fossilen Melanopsidae V. *Melanopsis handmanniana* nov. nomen und *Melanopsis fossilis* (Gmelin 1790) (Gastropoda, Prosobranchia). *Club Conchylia Informationen* 28 (1/2): 25–38.
- Fischer W (1996b) Beiträge zur Kenntnis der rezenten und fossilen Melanopsidae VI. Über *Melanopsis coaequata* Handmann 1887 und *Melanopsis rugosa* Handmann 1887 sowie ein Nachtrag zu Teil V. (Gastropoda, Prosobranchia). *Club Conchylia Informationen* 28 (3/4): 19–26.
- Fitton WH (1824) Inquiries respecting the Geological Relations of the Beds between the Chalk and the Purbeck Limestone in the South-East of England. *Annals of Philosophy, new series* 8: 365–383. <http://www.biodiversitylibrary.org/item/53939#page/385/mode/1up>
- Fitton WH (1836) Observations on some of the Strata between the Chalk and the Oxford Oolite, in the South-east of England. *Transactions of the Geological Society of London, Series 2* 4 (2): 103–388. doi: 10.1144/transgslb.4.2.103
- Fontannes F (1879–1882) Les invertébrés du bassin tertiaire du sud-est de la France. Les mollusques pliocènes de la vallée du Rhone et du Roussillon. Tome premier. Gastéropodes. Georg, F. Savy, Lyon, Paris, 276 pp., 12 pls. [1–76 pp. (March 1879); 77–168 pp. (February 1880); 169–268 pp. (June 1880); end date 1882 is given on front cover, possibly date of publication of plates, but not stated elsewhere]. <http://www.biodiversitylibrary.org/item/47327#page/7/mode/1up>
- Fontannes F (1881) Les terrains tertiaires du bassin de Crest. *Annales de la Société d’Agriculture. Histoire naturelle et arts utiles de Lyon* 3: 827–1046. <http://gallica.bnf.fr/ark:/12148/bpt-6k6154617d/f856.image>
- Fontannes F (1884) Description sommaire de la faune malacologique des formations saumâtres et d’eau douce du groupe d’Aix (Bartonien-Aquitainien) dans le Bas-Languedoc, la Provence et le Dauphiné. Georg, F. Savy, Lyon, Paris, 60 pp. <http://gallica.bnf.fr/ark:/12148/bpt6k55007367>
- Fontannes F (1887) Contribution à la faune malacologique des terrains néogènes de la Roumanie. *Archives du Muséum d’Histoire naturelle de Lyon* 4: 321–365.
- Forbes E (1856) On the Tertiary Fluvio-marine Formation of the Isle of Wight. Longman, Brown, Green, and Longmans, London, 162 pp. https://books.google.at/books?id=YibPAAAAMAAJ&printsec=frontcover&hl=de&source=gbs_ge_summary_r&cad=0#v=onepage&q&f=false
- Förster B (1892) Geologischer Führer für die Umgebung von Mülhausen i. E. Mittheilungen der Geologischen Landesanstalt von Elsass-Lothringen 3: 199–309. https://archive.org/stream/bub_gb_cY4MAAAAYAAJ#page/n267/mode/2up
- Fuchs T (1870a) III. Beiträge zur Kenntniss fossiler Binnenfaunen. III. Die Fauna der Congerenschichten von Radmanest im Banate. *Jahrbuch der k. k. geologischen Reichsanstalt* 20 (3): 343–364. http://opac.geologie.ac.at/wwwopacx/wwwopac.ashx?command=getcontent&server=images&value=JB0203_343_A.pdf

- Fuchs T (1870b) VII. Beiträge zur Kenntniss fossiler Binnenfaunen. IV. und V. Die Fauna der Congerenschichten von Tihany am Plattensee und Kúp bei Pápa in Ungarn. *Jahrbuch der k. k. geologischen Reichsanstalt* 20 (4): 531–548. https://opac.geologie.ac.at/wwwopacx/wwwopac.ashx?command=getcontent&server=images&value=JB0204_531_A.pdf
- Fuchs T (1873) Beiträge zur Kenntniss fossiler Binnenfaunen. VI. Neue Conchylienarten aus den Congerien-Schichten und aus Ablagerungen der sarmatischen Stufe. *Jahrbuch der k. k. geologischen Reichsanstalt* 23: 19–26. <http://www.biodiversitylibrary.org/item/110165#page/37/mode/1up>
- Fuchs T (1877) Studien über die jüngeren Tertiärbildungen Griechenlands. *Denkschriften der mathematisch-naturwissenschaftlichen Classe der Kaiserlichen Akademie der Wissenschaften* 37 (2): 1–42. <http://www.biodiversitylibrary.org/item/31616#page/381/mode/1up>
- Fuchs T, Karrer F (1870) Neue Brunnengrabungen in Wien und Umgebung. In: Fuchs T, Karrer F *Geologische Studien in den Tertiärbildungen des Wiener Beckens*. *Jahrbuch der k. k. geologischen Reichsanstalt* 20: 130–140. https://opac.geologie.ac.at/wwwopacx/wwwopac.ashx?command=getcontent&server=images&value=JB0201_113_A.pdf
- Gabb WM (1868) An attempt at a revision of the two families Strombidae and Aporrhaidae. *American Journal of Conchology* 4: 137–149. <http://www.biodiversitylibrary.org/item/54513#page/157/mode/1up>
- Gambetta L (1929) Molluschi. *Archivio zoologico italiano* 13: 45–117.
- Gassies J-B (1856) Description des coquilles univalves, terrestres et d'eau douce envoyées à la Société linnéenne de Bordeaux par M. le Capitaine Mayran. J.-B. Baillière, Paris, 13 pp. <http://gallica.bnf.fr/ark:/12148/bpt6k6529789x/f3.image>
- Gassies J-B (1857) Descriptions de coquilles terrestres et fluviatiles recueillies à la Nouvelle-Calédonie. *Journal de Conchyliologie* 6: 270–278. <http://biodiversitylibrary.org/item/53872#page/276/mode/1up>
- Gassies J-B (1858) Descriptions d'espèces nouvelles de l'Archipel Calédonien (3e article). *Journal de Conchyliologie* 7: 368–372. <http://biodiversitylibrary.org/item/53873#page/374/mode/1up>
- Gassies J-B (1861) Faune conchyliologique terrestre et fluvio-lacustre de la Nouvelle-Calédonie. *Actes de la Société Linnéenne de Bordeaux* 24: 211–330. <http://www.biodiversitylibrary.org/item/106381#page/225/mode/1up>
- Gassies J-B (1869) Description d'espèces inédites de la Nouvelle-Calédonie. - VIIIe article. *Journal de Conchyliologie* 17: 71–78. <http://biodiversitylibrary.org/item/53431#page/75/mode/1up>
- Gassies J-B (1870) Diagnoses d'espèces inédites provenant de la Nouvelle-Calédonie. Neuvième article. *Journal de Conchyliologie* 18: 140–150. <http://biodiversitylibrary.org/item/52282#page/144/mode/1up>
- Gassies J-B (1871) Faune conchyliologique terrestre et fluvio-lacustre de la Nouvelle-Calédonie. 2e partie. J.-B. Baillière et fils et F. Savy, Paris, 212 pp. <http://biodiversitylibrary.org/item/46322#page/9/mode/1up>
- Gassies J-B (1874) Descriptions d'espèces nouvelles, terrestres et fluvio-lacustres de la Nouvelle-Calédonie. *Journal de Conchyliologie* 22: 375–387. <http://biodiversitylibrary.org/item/53430#page/379/mode/1up>

- Gassies J-B (1880) Faune conchyliologique terrestre et fluvio-lacustre de la Nouvelle-Calédonie. Troisième partie. Baillière, Férét et Fils, Paris, Bordeaux, 108 pp. <http://biodiversitylibrary.org/item/42311#page/9/mode/1up>
- Gaudry A (1862–1867) Animaux fossiles et Géologie de l'Attique d'après les recherches faites en 1855–56 et en 1860 sous les auspices de l'Académie des Sciences. Librairie de la Société Géologique de France, Paris, Livr. 1–3: pp. 1–72, pls 1–12 (1862); Livr. 4–6: pp. 73–144, pls 13–24 (1863); Livr. 7–10: pp. 145–240, pls 25–40 (1864); Livr. 11–13: pp. 241–324, pls 41–52 (1865); Livr. 14–15: pp. 325–370, pls 53–60 (1866); Livr. 16–19: pp. 371–474, pls 61–75 (1867); Livr. 20: pp. 475–476 (1867). https://books.google.at/books?id=28VCdHzus08C&printsec=frontcover&hl=de&source=gb_s_ge_summary_r&cad=0#v=onepage&q&f=false (text); https://archive.org/details/bub_gb_bYbylmqfjV4C (atlas)
- Genth FA (1848) [Eocene Bildungen bei Wächtersbach im Hanauischen und ihre fossilen Reste; zur miocänen Geognosie des Mainzer-Beckens, seine Reste; Kalk-Tuff]. Neues Jahrbuch für Mineralogie, Geognosie, Geologie und Petrefaktenkunde 1848: 188–199. <http://www.biodiversitylibrary.org/page/36223720#page/216/mode/1up>
- Gentiluomo C (1868) Catalogo dei Molluschi terrestri e fluviali della Toscana. Bullettino Malacologico Italiano 1 (6): 81–100. <http://www.biodiversitylibrary.org/item/122154#page/131/mode/1up>
- Georgopoulou E, Neubauer TA, Harzhauser M, Kroh A, Mandic O (2015) An outline of the European Quaternary localities with freshwater gastropods: Data on geography and updated stratigraphy. Palaeontologia Electronica 18 (3): 48A (9 p.). <http://palaeo-electronica.org/content/2015/1328-quaternary-gastropods>
- Germain L (1921) Mollusques terrestres et fluviatiles de Syrie. Tome premier - Introduction et Gastéropodes. J.-B. Baillière et fils, Paris, 523 pp. <https://archive.org/details/mollusques-terres01germ>
- Germain L (1934) Études sur les Faunes malacologiques insulaires de l'Océan Pacifique. In: Berland L, Berlioz J, Bryan EH, Cheesman E, Joleaud L, Chopard L, Germain L, Guillaumin A, Holdhaus K, Mumford & Adamson, River P, Seurat L, Skottsberg C, Topsent E, Vallaux C (eds) Contribution à l'étude du peuplement zoologique et botanique des îles du Pacifique. Paul Lechevalier & Fils, Paris, 89–153. <http://gallica.bnf.fr/ark:/12148/bpt6k6395474n/f105.item.r=france+berberes.langFR>
- Gerstaecker A (1884) Vier Decaden von Neuropteren aus der Familie Megaloptera Burm. Mittheilungen aus dem naturwissenschaftlichen Vereine für Neu-Vorpommern und Rügen 16: 1–49.
- Gill T (1863) Systematic Arrangement of the Mollusks of the Family Viviparidae, and others, inhabiting the United States. Proceedings of the Academy of Natural Sciences of Philadelphia 15: 33–40. <http://www.biodiversitylibrary.org/item/18294#page/39/mode/1up>
- Gillet S, Gramann F, Steffens P (1978) Neue biostratigraphische Ergebnisse aus dem brackischen Neogen an Dardanellen und Marmara-Meer (Türkei). Newsletter on Stratigraphy 7 (1): 53–64.
- Gillet S, Marinescu F (1971) La faune malacologique pontienne de Radmanesti (Banat Roumain). Mémoires - L'Institut de Géologie et de Géophysique 15: 1–78.

- Girotti O (1972) Correlabilita', mediante molluschi d'acqua dolce, del Calabrian di Monte Mario (Roma) col Villafranchiano superiore. *Geologica Romana* 11: 229–235.
- Gistel J (1848) *Naturgeschichte des Thierreichs. Für höhere Schulen.* Hoffmann'sche Verlags-Buchhandlung, Stuttgart, xvi + 216 pp. Second edition from 1851 available from BHL: <http://www.biodiversitylibrary.org/item/175526#page/7/mode/1up>
- Glaubrecht M (1993) Mapping the diversity: geographical distribution of the freshwater snail *Melanopsis* (Gastropoda: ?Cerithioidea: Melanopsidae) with focus on its systematics in the Mediterranean Basin. *Mitteilungen aus dem Hamburger Zoologischen Museum und Institut* 90: 41–97.
- Glaubrecht M (1996) *Evolutionsökologie und Systematik am Beispiel von Süß- und Brackwasserschnecken (Mollusca: Caenogastropoda: Cerithioidea): Ontogenese-Strategien, Paläontologische Befunde und Zoogeographie.* Backhuys, Leiden, 499 pp.
- Glaubrecht M (2011) Toward solving Darwin's "mystery": speciation and radiation in freshwater gastropods. *American Malacological Bulletin* 29: 187–216. doi: 10.4003/006.029.0211
- Glibert M (1962) Les Mesogastropoda fossiles du Cénozoïque étranger des collections de l'Institut Royal des Sciences Naturelles de Belgique. Première partie. Cyclophoridae à Stiliferidae (inclus). Institut Royal des Sciences naturelles de Beligiques, Mémoires. Deuxième série 69: 1–305.
- Glöer P (2002) *Die Tierwelt Deutschlands, 73. Teil: Die Süßwassergastropoden Nord- und Mitteleuropas. Bestimmungsschlüssel, Lebensweise, Verbreitung.* ConchBooks, Hackenheim, 327 pp.
- Gmelin JF (1791) *Caroli a Linné, systema naturae per regna tria naturae, secundum classes, ordines, genera, species, cum characteribus, differentiis, synonymis, locis. Editio decima tertia, aucta, reformata. Tomus I. Pars VI: Vermes.* Georg Emanuel Beer, Lipsiae, 3021–3910 pp. <http://www.biodiversitylibrary.org/item/83098#page/5/mode/1up>
- Gojranović-Kramberger K (1892) Das Vorkommen der Paludinenschichten in den Maria-Goricaer Hügeln in Croatien. *Verhandlungen der k. k. geologischen Reichsanstalt* 1892 (4): 108–110. <http://biodiversitylibrary.org/page/39121934#page/534/mode/1up>
- Gould AA (1847) [Descriptions of the following species of *Melania*, from the Collection of the Exploring Expedition]. *Proceedings of the Boston Society of Natural History* 2: 222–225. doi: 10.5962/bhl.part.3322
- Gozhik PF (2002) *Pontychni prsnovodni molyusky pivdnya Ukrayiny i Moldovy.* Privately published, Kiev, 96 pp.
- Gozhik PF, Datsenko LN (2007) *Presnovodnyye Mollyuski pozdnego kaynozoya yuga Vostochnoy Yevropy. Chast' II. Semeystva Sphaeridae, Pisidiidae, Corbiculidae, Neritidae, Viviparidae, Valvatidae, Bithyniidae, Lithoglyphidae, Melanopsidae.* Natsional'naya Akademiya Nauk Ukrainy, Institut Geologicheskikh Nauk, Kiev, 253 pp.
- Graells M de la P (1846) *Catalogo de los moluscos terrestres y de agua dulce observado en España, y descripcion y notas de algunas especies nuevas o poco conocidas del mismo pais.* Viuda é Hijos de Don Antonia Calleja; Calleja, Ojea y Compañía, Madrid, Lima, 23 pp. <http://biodiversitylibrary.org/item/51553#page/7/mode/1up>
- Grateloup J-PS de (1828) *Tableau des coquilles Fossiles qu'on rencontre dans des terrains calcaires tertiaires (faluns) des environs de Dax, dans le département des Landes (Arti-*

- cles 1–3). Bulletin d'Histoire Naturelle de la Société linnéenne de Bordeaux 2: 72–109, 123–158, 192–204. <http://www.biodiversitylibrary.org/item/110491#page/82/mode/1up> [publication dates indicated on p. 261 of the volume]
- Grateloup J-PS de (1838) Conchyliologie fossile du bassin de l'Adour. (3.me Mémoire). V. Mémoire sur les Coquilles fossiles des Mollusques terrestres et fluviatiles (de la classe des Trachéliopodes) observées dans les terrains tertiaires du bassin de l'Adour. Actes de la Société linnéenne de Bordeaux 10: 92–152. <http://www.biodiversitylibrary.org/item/101658#page/30/mode/1up>
- Grateloup J-PS de (1840) Conchyliologie fossile des terrains tertiaires du bassin de l'Adour (environs de Dax). Tome 1: Univalves. Atlas. Th. Lafargue, Bordeaux, 48 pls. https://books.google.at/books?id=A-oQAAAAIAAJ&printsec=frontcover&hl=de&source=gbg_summary_r&cad=0#v=onepage&q&f=false
- Graves L (1847) Essai sur la topographie géognostique du département de l'Oise. Achile Desjardins, Beauvais, xv + 804 pp. <https://patrimoine.mines-paristech.fr/items/viewer/1400-#page/Image+2/mode/2up>
- Gray JE (1843) Notes on the Materials at present existing towards a Fauna of New Zealand. In: Dieffenbach E (Ed.) Travels in New Zealand; with contributions to the geography, geology, botany, and natural history of that country. Vol. II. John Murray, London, 177–295. <http://www.biodiversitylibrary.org/item/67828#page/191/mode/1up>
- Gray JE (1847) A list of the genera of recent Mollusca, their synonyma and types. Proceedings of the Zoological Society of London 15: 129–242.
- Griffith E, Pidgeon E (1833–1834) Vol. 12. The Mollusca and Radiata. Arranged by the Baron Cuvier, with supplementary additions to each order. In: Griffith E (Ed.) The Animal Kingdom arranged in conformity with its organization. Whittaker and Co., London, viii + 601 pp., 61 pls. [pp. 1–192, Mollusca pls. 1–27, 29–35, 38–39, 1833; pp. 193–601 + viii, Mollusca corrected pls. 28*, 36*, 37*; pls. 40, 41, 1834]. <http://www.biodiversitylibrary.org/item/40578#page/11/mode/1up>
- Guirao Á (1854) Zwei Novitäten. Malakozoologische Blätter 1: 32–33. <http://www.biodiversitylibrary.org/item/53876#page/48/mode/1up>
- Gümbel CW von (1861) Geognostische Beschreibung des bayerischen Alpengebirges und seines Vorlandes. Justus Perthes, Gotha, 950 pp. http://reader.digitale-sammlungen.de/fs1/object/display/bsb10806180_00005.html
- Guppy RJL (1866) On the Relations of the Tertiary Formations of the West Indies. The Quarterly Journal of the Geological Society of London 22: 570–590. doi: 10.1144/GSL.JGS.1866.022.01-02.51
- Guppy RJL (1867) Descriptions of a New Genus and Six New Species of Mollusca, from the Caribbean Miocene. The Geological Magazine 4 (41): 500–501. <http://www.biodiversitylibrary.org/item/101219#page/556/mode/1up>
- Halaváts G von (1903) Die Fauna der pontischen Schichten in der Umgebung des Balatonsees. Resultate der wissenschaftlichen Erforschung des Balatonsees 1 (1): 1–80.
- Halaváts G von (1914) Geologischer Aufbau der Gegend von Ujeháza, Holczmány und Oltszakadát. Jahresbericht der Königlich ungarischen Geologischen Anstalt 1913: 410–422.

- Handmann R (1882) Die fossile Molluskenfauna von Kottingbrunn. Jahrbuch der k. k. geologischen Reichsanstalt 32: 543–564. <http://www.biodiversitylibrary.org/item/110147-#page/579/mode/1up>
- Handmann R (1887) Die fossile Conchylienfauna von Leobersdorf im Tertiärbecken von Wien. Aschendorff, Münster, 47 pp.
- Hantken MRvP (1878) Die Kohlenflötze und der Kohlenbergbau in den Ländern der ungarischen Krone. Gebrüder Légrády, Budapest, 354 pp.
- Harris GD (1899) The Lignitic stage. Part II. Scaphopoda, Gastropoda, Pteropoda and Cephalopoda. Bulletins of American Paleontology 3 (11): 1–128. <http://www.biodiversitylibrary.org/item/95774#page/15/mode/1up>
- Harris TW (1841) A report on the insects of Massachusetts, injurious to vegetation. Folsom, Wells, and Thurston, Cambridge, 459 pp. <http://www.biodiversitylibrary.org/item/27609#page/5/mode/1up>
- Harzhauser M, Kowalke T, Mandic O (2002) Late Miocene (Pannonian) Gastropods of Lake Pannon with Special Emphasis on Early Ontogenetic Development. Annalen des Naturhistorischen Museums in Wien, Serie A 103: 75–141. http://www.zobodat.at/pdf/ANNA_103A_0075-0141.pdf
- Harzhauser M, Neubauer TA, Georgopoulou E, Esu D, D’Amico C, Pavia G, Giuntelli P, Carnevale G (2015) Late Messinian continental and Lago-Mare gastropod assemblages from the Tertiary Piedmont Basin, NW. Bollettino della Società Paleontologica Italiana 54 (1): 1–53. doi: 10.4435/BSPI.2015.1
- Haszprunar G (2014) A nomenclator of extant and fossil taxa of the Valvatidae (Gastropoda, Ectobranchia). ZooKeys 377: 1–172. doi: 10.3897/zookeys.377.6032
- Heller J, Mordan P, Ben-Ami F, Sivan N (2005) Conchometrics, systematics and distribution of *Melanopsis* (Mollusca: Gastropoda) in the Levant. Zoological Journal of the Linnean Society 144: 229–260. doi: 10.1111/j.1096-3642.2005.00172.x
- Heller J, Sivan N (2000) A new species of *Melanopsis* from the Golan heights, Southern Levant (Gastropoda: Melanopsidae). Journal of Conchology 37 (1): 1–5.
- Heller J, Sivan N (2001) *Melanopsis* from the Mid-Pleistocene site of Gesher Benot Ya’aqov (Gastropoda: Cerithioidea). Journal of Conchology 37 (2): 127–147.
- Heller J, Sivan N (2002a) *Melanopsis* from the Pleistocene site of ‘Ubeidiya, Jordan Valley: direct evidence of early hybridization (Gastropoda: Cerithioidea). Biological Journal of the Linnean Society 75: 39–57. doi: 10.1046/j.1095-8312.2002.00005.x.
- Heller J, Sivan N (2002b) *Melanopsis* from the Pliocene site of ‘Erq el-Ahmar, Jordan Valley (Gastropoda: Cerithioidea). Journal of Conchology 37 (6): 607–625.
- Heller J, Sivan N, Ben-Ami F (2002) Systematics of *Melanopsis* from the coastal plain of Israel (Gastropoda: Cerithioidea). Journal of Conchology 37 (6): 589–606.
- Heller J, Sivan N, Motro U (1999) Systematics, distribution and hybridization of *Melanopsis* from the Jordan Valley (Gastropoda: Prosobranchia). Journal of Conchology 36 (5): 49–81.
- Hermite H (1879) Études géologiques sur les îles Baléares. Première partie, Majorque et Minorque. F. Pichon, F. Savy, Paris, 362 pp. doi: 10.5962/bhl.title.13990

- Herrmannsen AN (1846–1847) *Indicis generum malacozoorum primordia. Nomina subgenerum, generum, familiarum, tribuum, ordinum, classium; adjectis auctoribus, temporibus, locis systematicis atque literariis, etymis, synonymis. Praetermittuntur Cirripedia, Tunicata et Rhizopoda. Vol. I.* Fischer, Cassellis, i–xxvii, 1–104 pp. (September 1846), 105–232 pp. (December 1846), 233–360 pp. (March 1847), 361–488 pp. (April 1847), 489–616 pp. (May 1847), 617–637 pp. (July 1847). <http://www.biodiversitylibrary.org/item/40627#page/9/mode/1up>
- Hoeninghaus FW (1831) Versuch einer geognostischen Eintheilung seiner Versteinerung-Sammlung (Dritter Theil. Tertiär-Gebirge.). *Jahrbuch für Mineralogie, Geognosie, Geologie und Petrefaktenkunde* 2: 132–170. <https://archive.org/stream/jahrbuchfrminer00unkngoog#page/n157/mode/2up>
- Hoernes R (1876) Ein Beitrag zur Kenntniss fossiler Binnenfaunen. (Süsswasserschichten unter den sarmatischen Ablagerungen am Marmorameere.). *Sitzungsberichte der mathematisch-naturwissenschaftlichen Classe der Kaiserlichen Akademie der Wissenschaften* 74 (1): 7–34. <http://www.biodiversitylibrary.org/item/35546#page/13/mode/1up>
- Hofmann K (1870) A Zsily-völgyi szénteknö. A Magyarhoni Földtani Társulat Munkálatai 5: 1–57. http://epa.oszk.hu/02000/02013/00005/pdf/EPA02013_magyarhoni_foldtani_tarsulat_1870.pdf
- Hörnes M (1851–1856) Die fossilen Mollusken des Tertiär-Beckens von Wien. I. Univalven. *Abhandlungen der Geologischen Reichsanstalt* 3: 185–296 (1853), 385–460 (1855), 461–736 (1856).
- Hörnes M (1855) Über einige neue Gastropoden aus den östlichen Alpen. *Denkschriften der Kaiserlichen Akademie der Wissenschaften. Mathematisch-naturwissenschaftliche Classe* 10 (2): 173–178. <http://www.biodiversitylibrary.org/item/30071#page/323/mode/1up>
- Huică IV (1977) Studiul geologic al depozitelor miocene și pliocene dintre Valea Sohodol și Valea Blahnița, județul Gorj (depresiunea getică). *Anuarul institutului de Geologie și Geofizică* 51: 5–68.
- Hutton FW (1873) *Catalogue of the Tertiary Mollusca and Echinodermata of New Zealand, in the collection of the Colonial Museum. Colonial Museum and Geological Survey Department, Wellington*, xvi + 48 pp. doi: 10.5962/bhl.part.22727
- Hutton FW (1883) List of the Freshwater Shells of New Zealand. *Proceedings of the Linnean Society of New South Wales* 7: 67–68. <https://archive.org/stream/proceedingsoflin0107linn#page/66/mode/2up>
- International Commission on Zoological Nomenclature (1944) Opinion 184. on the status of names first published in volumes 1 to 11 of Martini (F. H. W.) and Chemnitz (J. H.), *Neues Systematischer Conchylien Cabinet, Nürnberg 1769–1795. Opinions And Declarations Rendered By The International Commission On Zoological Nomenclature* 3: 25–35. <http://www.biodiversitylibrary.org/item/107759#page/47/mode/1up>
- International Commission on Zoological Nomenclature (1999) *International Code of Zoological Nomenclature. International Trust for Zoological Nomenclature, London*, 306 pp. Online available from <http://www.nhm.ac.uk/hosted-sites/iczn/code/index.jsp>

- Ippolito F (1947) Contributi alla conoscenza della malacofauna delle ligniti di Colle dell'Oro presso Terni. Rendiconti della Reale Accademia delle scienze fisiche e matematiche di Napoli, seria 4 15: 10 p. [page range unknown, only Separatum available]
- Iredale T (1915) A commentary on Suter's "Manual of the New Zealand Mollusca". Transactions and Proceedings of the New Zealand Institute 47: 417–497. <http://www.biodiversitylibrary.org/page/3343255#page/447/mode/1up>
- Issel A (1865) Catalogo dei molluschi raccolti dalla missione italiana in Persia. Stamperia Reale, Torino, 55 pp. (The work was reprinted in the Memorie della Reale Accademia delle Scienze di Torino, Serie seconda 23 [1866]: 387–439. <http://www.biodiversitylibrary.org/item/97653#page/445/mode/1up>)
- Issel A (1866) Dei molluschi raccolti nella provincia di Pisa. Memorie della Società Italiana di Scienze Naturali 2 (1): 1–38. <http://babel.hathitrust.org/cgi/pt?id=mdp.39015040091830;view=1up;seq=9>
- Izzatullaev ZI, Starobogatov YI (1984) Pod *Melanopsis* (Gastropoda, Pectinibranchia) i ego predstaviteli, obitayushchiye v vodoyemakh SSSR [The genus *Melanopsis* (Gastropoda, Pectinibranchia) and its representatives that live in the waters of the USSR]. Zoologicheskii Zhurnal 48 (10): 1471–1483. [In Russian]
- Jekelius E (1944) Sarmat und Pont von Soceni (Banat). Memoriile Institutului geologic al României 5: 1–167.
- Jodot P (1958) Les faunes des mollusques continentaux reparties dans le sud-est de l'Espagne entre le Miocène supérieur et le Quaternaire. Memorias y comunicaciones del Instituto Geológico 17: 3–133.
- Jodot P (1965) Malacologie continentale de quelques gisements quaternaires du littoral atlantique marocain [Avant-propos par Georges Lecointre]. Notes et Mémoires du Service Géologique Maroc 25 (185): 101–108.
- Kapan Yeşilyurt S, Taner G (2002) Datça yarımadasının geç pliyosen Pelecypoda ve Gastropoda faunası ve stratigrafisi (Muğla-Güneybatı Anadolu). MTA Dergisi 125: 89–120. [in Turkish] http://www.mta.gov.tr/v2.0/daire-baskanliklari/bdt/kutuphane/mtadergi/125_6.pdf
- Kennard AS (1942a) The Histoire and Prodrome of Férussac. Part I. Proceedings of the Malacological Society of London 25 (1): 12–17.
- Kennard AS (1942b) The Histoire and Prodrome of Férussac. Part II. Proceedings of the Malacological Society of London 25 (3): 105–110.
- Kennard AS (1942c) The Histoire and Prodrome of Férussac. Part III. Proceedings of the Malacological Society of London 25 (3): 111–118.
- Klein A (1852) Conchylien der Süßwasserkalkformation Württembergs. Jahreshefte des Vereins für vaterländische Naturkunde in Württemberg 8: 157–164.
- Klika G (1891) Die tertiären Land- und Süßwasser-Conchylien des nordwestlichen Böhmen. Archiv der naturwissenschaftlicher Landesdurchforschung von Böhmen 7: 1–122. <http://www.biodiversitylibrary.org/item/112127#page/331/mode/1up>
- Kobelt W (1879–1880) Iconographie der Land- & Süßwasser-Mollusken mit vorzüglicher Berücksichtigung der europäischen noch nicht abgebildeten Arten von E. A. Rossmässler. VII. Band. C. W. Kreidel, Wiesbaden, Hefte 1–3: 1–24 pp. (1879), Hefte 4–6: 24–94 pp. (1880) pp. <http://www.biodiversitylibrary.org/item/55038#page/377/mode/1up>

- Kobelt W (1881) Catalog der im europäischen Faunengebiet lebenden Binnen-Conchylien. 2. Auflage. Fischer, Cassel, 294 pp. <http://www.biodiversitylibrary.org/item/51396#page/111/mode/1up>
- Koch F (1917) Levantinska fauna Vukomerickih gorica. Glasnik Hrvatskoga prirodoslovnoga društva 29: 7–17. [in Croatian] <http://www.biodiversitylibrary.org/item/41618#page/217/mode/1up>
- Köhler F, Glaubrecht M (2010) Uncovering an overlooked radiation: molecular phylogeny and biogeography of Madagascar's endemic river snails (Caenogastropoda: Pachychilidae: *Madagasikara* gen. nov.). Biological Journal of the Linnean Society 99 (4): 867–894. doi: 10.1111/j.1095-8312.2009.01390.x
- Kollmann HA (1984) *Megalonoda* n. gen. (Melanopsidae, Gastropoda) aus der Oberkreide der Nördlichen Kalkalpen (Österreich). Annalen des Naturhistorischen Museums in Wien, Serie A 86: 55–62. http://www.zobodat.at/pdf/ANNA_86A_0055-0062.pdf
- Kormos T (1903) Palaeontologiai közlemények. Földtani Közlöny 33: 451–462 (hungarian), 496–503 (german). http://epa.oszk.hu/01600/01635/00054/pdf/Foldtani_kozlony_EPA01635_1903_10-12_451-462.pdf
- Kormos T (1904) Új adatok a Püspökfürdő élő csigáinak ismeretéhez. Állattani Közlemények 3: 102–111. [in Hungarian]
- Kormos T (1905) Über den Ursprung der Thermenfauna von Püspökfürdő. Földtani Közlöny 35 (8–9): 421–450.
- Koşal-Şahin S, Yıldırım MZ (2007) The Mollusc Fauna of Lake Sapanca (Turkey: Marmara) and Some Physico-Chemical Parameters of Their Abundance. Turkish Journal of Zoology 31: 47–52. <http://journals.tubitak.gov.tr/zoology/issues/zoo-07-31-1/zoo-31-1-7-0511-12.pdf>
- Krauss F (1852) Die Mollusken der Tertiärformation von Kirchberg an der Iller. Jahreshefte des Vereins für vaterländische Naturkunde in Württemberg 8: 136–157.
- Kühn O (1946) Das Alter der Prominaschichten und der innereozänen Gebirgsbildung. Jahrbuch der Geologischen Bundesanstalt 91: 49–94. http://opac.geologie.ac.at/wwwopac/wwwopac.ashx?command=getcontent&server=images&value=JB0911_049_A.pdf
- Kühn O (1951) Süßwassermiozän von bosnischem Typus in Griechenland. Annales Géologiques des Pays Helléniques 3: 185–192.
- Kühn O (1963) Das Süßwassermiozän von Attika. Praktika tis Akademias Athinon 38: 370–400.
- Kurr JG von (1856) Land- und Süßwasserconchylien der Tertiärformation Oberschwabens. Jahreshefte des Vereins für vaterländische Naturkunde in Württemberg 12: 38–43. <http://www.biodiversitylibrary.org/item/33518#page/46/mode/1up>
- Lamarck J-BPA de M de [1797] Tableau encyclopédique et méthodique des trois règnes de la nature. Vingt-unième partie. Mollusques testacés. Henri Agasse, Paris, pl. 287–390. <http://biodiversitylibrary.org/item/130185#page/9/mode/1up>
- Lamarck J-BPA de M de (1799) Prodrome d'une nouvelle classification des coquilles, comprenant une rédaction appropriée des caractères génériques, et l'établissement d'un grand nombre des genres nouveaux. Mémoires de la Société d'Histoire Naturelle de Paris 1: 63–91.
- Lamarck J-BPA de M de (1804) Suite des mémoires sur le fossiles des environs de Paris. Annales du Muséum national d'histoire naturelle 4: 429–436. <http://biodiversitylibrary.org/item/93709#page/432/mode/1up>

- Lamarck J-BPA de M de (1822) Histoire naturelle des animaux sans vertèbres, présentant les caractères généraux et particuliers de ces animaux, leur distribution, leurs classes, leurs familles, leurs genres, et la citation des principales espèces qui s'y rapportent; précédée d'une introduction offrant la détermination des caractères essentiels de l'animal, sa distinction du végétal et des autres corps naturels, enfin, l'exposition des principes fondamentaux de la zoologie. Tome sixième, deuxième partie. Privately published, Paris, 232 pp. <http://www.biodiversitylibrary.org/item/47441#page/359/mode/1up>
- Lang AF (1823) Nuncium pro conchyliologis. Isis von Oken 13: 430. <http://www.biodiversitylibrary.org/item/47603#page/977/mode/1up>
- Lea I (1837) Observations on the Naiades; and Descriptions of New Species of that, and other Families. Transactions of the American Philosophical Society, new series 5: 23–119. <http://www.biodiversitylibrary.org/item/110958#page/45/mode/1up>
- Lea I (1841) [untitled; about fifty-seven new species of the genus *Melania*]. Proceedings of the American Philosophical Society held at Philadelphia 2: 11–15. <http://www.biodiversitylibrary.org/item/107402#page/19/mode/1up>
- Lea I (1842) [untitled; replacement names for preoccupied names of *Melania*]. Proceedings of the American Philosophical Society held at Philadelphia 2 (24): 237. <http://www.biodiversitylibrary.org/item/107402#page/245/mode/1up>
- Lea I (1862) Description of a New Genus (*Goniobasis*) of the Family Melanidae and eighty-two new Species. Proceedings of the Academy of Natural Sciences of Philadelphia 14: 262–272. <http://www.biodiversitylibrary.org/item/18281#page/274/mode/1up>
- Lea I, Lea HC (1850) Description of a new genus of the family Melaniana and many new species of the genus *Melania* chiefly collected by Hugh Cuming Esq., during his zoological voyage in the east and now first described. Proceedings of the Zoological Society of London 18: 179–197. <http://www.biodiversitylibrary.org/item/46212#page/241/mode/1up>
- Letourneux A (1876a) Mollusques terrestres et fluviatiles du Maroc. Revue des Sciences Naturelles 5 (1): 45–49. <http://www.biodiversitylibrary.org/item/52495#page/55/mode/1up>
- Letourneux A (1876b) Mollusques terrestres et fluviatiles du Maroc. Bulletin de la Société des Sciences Physiques, Naturelles et Climatologiques d'Alger 13 (4): [page range unknown; not seen].
- Letourneux A, Bourguignat JR (1887) Prodrome de la malacologie terrestre et fluviatile de la Tunisie. Imprimerie nationale, Paris, 166 pp. http://doc.rero.ch/record/11944/files/mol_r5.pdf
- Leymerie A (1881) Description géologique et paléontologique des Pyrénées de la Haute Garonne. Édouard Privat, Toulouse, 1010 pp. [http://jubilotheque.upmc.fr/ead.html?id=GR_000322_001&c=GR_000322_001_page1#!{%22content%22:\[%22GR_000322_001_page1%22,false,%22%22\]}](http://jubilotheque.upmc.fr/ead.html?id=GR_000322_001&c=GR_000322_001_page1#!{%22content%22:[%22GR_000322_001_page1%22,false,%22%22]})
- Linnaeus C (1758) Systema naturae per regna tria naturae, secundum classes, ordines, genera, species, cum characteribus, differentiis, synonymis, locis. Tomus I. Editio decima, reformata. Laurentius Salvius, Holmiae, iv + 824 pp. <http://www.biodiversitylibrary.org/item/10277-#page/3/mode/1up>
- Linnaeus C (1767) Systema naturae, Tom. I. Pars II. Editio duodecima reformata. Laurentius Salvius, Holmiae, 533–1327, [1–37] pp. <http://biodiversitylibrary.org/item/137240-#page/5/mode/1up>

- Locard A (1878) Description de la faune de la mollasse marine et d'eau douce du Lyonnais et du Dauphiné. Archives du Muséum d'Histoire naturelle de Lyon 2: 1–284.
- Locard A (1883a) Malacologie des lacs de Tibériade, d'Antioche et d'Homs, Syrie. Archives du Muséum d'Histoire naturelle de Lyon 3: 195–293. <http://www.biodiversitylibrary.org/item/37326#page/207/mode/1up>
- Locard A (1883b) Recherches paléontologiques sur les dépôts tertiaires à *Milne-Edwardsia* et *Vivipara* du Pliocène inférieur du département de l'Ain. Annales de l'Académie de Mâcon, IIe Série 6: 2–166. <http://gallica.bnf.fr/ark:/12148/bpt6k2136624/f3.image>. Printed date at page 1 of the volume is 1888, but the paper was already offered with year “1883” in the book reviews at the end of vol. 32 (1) of the Journal de Conchyliologie (1884).
- Locard A (1893) Monographie des mollusques tertiaires terrestres et fluviatiles de la Suisse. Deuxième partie. Mémoires de la Société Paléontologique Suisse 19 (1892): 131–275.
- Łomnicki AM (1886) Słodkowodny utwór trzeciorzędny na Podolu galicyjskiem. Sprawozdania Komisji Fizjograficznej Polskiej Akademii Umiejętności 20: 48–119. [in Polish]
- Lörenthey E (1902) Die Pannonische Fauna von Budapest. Palaeontographica 48: 137–256. <http://www.biodiversitylibrary.org/item/103637#page/157/mode/1up>
- Lubenescu V (1981) Studiul biostratigrafic al Neogenului superior din sud-vestul Transilvaniei. Anuarul Institutului de Geologie si Geofizica 58: 123–202.
- Lubenescu V (1985) Nouvelles espèces de mollusques du Pannonien du SO de la dépression de Transylvanie. Revue Roumaine de Géologie, Géophysique et Géographie 29: 73–79.
- Ludwig R (1865) Fossile Conchylien aus den tertiären Süsswasser- und Meerwasser-Ablagerungen in Kurhessen, Grossherzogtum Hessen und der Bayer'schen Rhön. Palaeontographica 14: 40–97. <http://www.biodiversitylibrary.org/item/44419#page/48/mode/1up>
- Lueger JP (1980) Die Molluskenfauna aus dem Pannon (Obermiozän) des Fölligberges (Eisenstädter Bucht) im Burgenland (Österreich). Mitteilungen der österreichischen geologischen Gesellschaft 73: 95–134.
- Lycett J (1848) Notes on the distribution of the Fossil Conchology of the Oolitic Formations in the vicinity of Minchinhampton, Gloucestershire. The Annals and Magazine of Natural History, second series 2: 248–259. <http://www.biodiversitylibrary.org/page/18686254#page/262/mode/1up>
- Macarovici N (1940) Recherches géologiques et paléontologiques dans la Bessarabie méridionale (Roumanie). Annales scientifiques de l'Université de Jassy 36 (1): 177–422.
- Magrograssi A (1928) La fauna levantina di Coo e di Rodi. Atti della Società Italiana di Scienze Naturali e del Museo Civico di Storia Naturale in Milano 67: 249–264.
- Mandic O, Kurečić T, Neubauer TA, Harzhauser M (2015) Stratigraphic and palaeogeographic significance of lacustrine mollusks from the Pliocene *Viviparus* beds in central Croatia. Geologia Croatica 68 (3): 179–207. doi: 10.4154/gc.2015.15
- Manzoni A (1870) Della Fauna Marina di due lembi Miocenici dell'alta Italia. Sitzungsberichte der mathematisch-naturwissenschaftlichen Classe der Kaiserlichen Akademie der Wissenschaften 60 (1): 475–504. <http://www.biodiversitylibrary.org/item/108174#page/583/mode/1up>
- Marshall P (1918) The Tertiary Molluscan Fauna of Pakaurangi Point. Kaipara Harbour. Transactions and Proceedings of the New Zealand Institute 50: 263–278. http://rsnz.natlib.govt.nz/volume/rsnz_50/rsnz_50_00_003260.pdf

- Martens E von (1866) Über neue Landschnecken aus Ost-Indien und über zwei Seesterne von Costa-Rica. Monatsberichte der Königlich Preussische Akademie der Wissenschaften zu Berlin 1865: 51–59. <http://www.biodiversitylibrary.org/page/35168448#page/61/mode/1up>
- Martens E von (1874) Ueber vorderasiatische Conchylien nach den Sammlungen des Prof. Hausknecht. Novitates Conchologicae. Supplement 5: 1–127. <http://biodiversitylibrary.org/item/46803#page/7/mode/1up>
- Martini FHW (1777) Neues systematisches Conchylien-Cabinet. III. Band. Gabriel Nikolaus Raspe, Nürnberg, 434 pp. <http://www.biodiversitylibrary.org/item/94988#page/5/mode/1up>
- Marwick J (1926) New Tertiary Mollusca from North Taranaki. Transactions and Proceedings of the New Zealand Institute 56: 317–331. http://rsnz.natlib.govt.nz/volume/rsnz_56/rsnz_56_00_002910.pdf
- Matheron P (1842) Catalogue méthodique et descriptif des corps organisés fossiles du département des Bouches-du-Rhône et lieux circonvoisins; précédé D'un Mémoire sur les terrains supérieurs au Grès Bigarré du S. E. de la France. Répertoire des Travaux de la Société de statistique de Marseille 6: 81–341. <http://gallica.bnf.fr/ark:/12148/bpt6k5719359d/f81.image>
- Matheron P (1843) Catalogue méthodique et descriptif des corps organisés fossiles du département des Bouches-du-Rhône et lieux circonvoisins; précédé D'un Mémoire sur les terrains supérieurs au Grès Bigarré du S. E. de la France. Carnaud Fils, Marseille, 269 pp. + 41 pls. https://books.google.at/books?id=IC4-AAAAcAAJ&printsec=frontcover&hl=de&source=gbs_ge_summary_r&cad=0#v=onepage&q&f=false
- Matheron P (1878–1880) Recherches paléontologiques dans le Midi de la France, ou étude sur les animaux fossiles découverts dans cette région présentant la description et la figure des espèces nouvelles, douteuses ou peu connues avec l'énumération méthodique des corps organisés fossiles qui le accompagnent dans leurs gisements stratigraphiques. Privately published, Marseille, Livr. 1–2: 4 pp., pl. B15, B18, B20, C8, C9, C11, C13, C14, C17, C21, F22, G10 (March 1878), Livr. 3–4: pl. A2, B16, B17, C1, C2, C12, C20, D24, D25, E8, F15, G12 (June–July 1878), Livr. 5–6: 7 pp., pl. A1, B8, C3, C4, C8, C16, C18, C19, C24, J22, O1 (March 1879); Livr. 7: pl. B2, B27, F41, F42, F43, J13 (December 1880).
- Mayer C (1871) Découverte des couches à Congéries dans le bassin du Rhône. Vierteljahrsschrift der Naturforschenden Gesellschaft in Zürich 16: 185–203. <http://www.biodiversitylibrary.org/item/34569#page/201/mode/1up>
- Mayer-Eymar C (1903) Description de coquilles fossiles des terrains tertiaires inférieurs (Suite). Journal de Conchyliologie 51: 308–320. <http://biodiversitylibrary.org/page/16311599#page/328/mode/1up>
- Mayer-Eymar K (1887) Systematisches Verzeichniss der Kreide- und Tertiär-Versteinerungen der Umgegend von Thun, nebst Beschreibung der neuen Arten. Beiträge zur geologischen Karte der Schweiz, Beilage 24 (2): i–xxviii + 1–128.
- M'Coy F, Griffith RJ (1844) A synopsis of the characters of the Carboniferous limestone fossils of Ireland. University Press, Dublin, 207 pp. doi: 10.5962/bhl.title.11559
- Meek FB (1877) Palaeontology. United States Geological Exploration of the Fortieth Parallel, Report 4: 1–197. <http://www.biodiversitylibrary.org/item/104531#page/19/mode/1up>

- Meijer T (1990) Notes on Quaternary freshwater Mollusca of the Netherlands, with descriptions of some new species. *Mededelingen van de Werkgroep voor Tertiaire en Kwartaire Geologie* 26 (4): 145–181.
- Melleville M (1843) Mémoire sur les Sables tertiaires inférieurs du bassin de Paris, avec la description de 78 espèces de coquilles fossiles inédites de ce terrain. *Annales des Sciences Géologiques* 2 (1–2): 1–29, 77–120. reprint available from Google Books: https://books.google.at/books?id=rzF9m0OG85AC&printsec=frontcover&hl=de&source=gbs_ge_summary_r&cad=0#v=onepage&q&f=false
- Merian P (1849) Ueber eine marinische Tertiärformation am Randen bei Schaffhausen. *Bericht über die Verhandlungen der Naturforschenden Gesellschaft in Basel* 8: 30–31. <http://www.biodiversitylibrary.org/item/105004#page/36/mode/1up>
- Mertin H (1939) Über Brackwasserbildungen in der Oberen Kreide des nördlichen Harzvorlandes. *Nova Acta Leopoldina, Neue Folge* 7 (48): 142–263.
- Michelin H (1833) Section 2, Classe V. - Mollusques. Mélanopside. *Magasin de Zoologie* 3: pl. 29. <http://www.biodiversitylibrary.org/item/114093#page/183/mode/1up> (notice); <http://www.biodiversitylibrary.org/item/114093#page/453/mode/1up> (plate)
- Michelotti G (1840) Rivista di alcune specie fossili della famiglia dei gasteropodi. *Annali delle Scienze del Regno Lombardo-Veneto. Opera periodica di alcuni collaboratori* 10: 137–162.
- Miletić-Spajić O (1961) Beiträge zur Kenntnis der pannonischen (unteren Congerien-) Schichten in der Umgebung von Beograd (Zaklopaca, Begaljica, Vrcin). *Annales Géologiques de la Péninsule Balkanique* 28: 247–264.
- Montfort PD de (1810) *Conchyliologie systématique et classification méthodique de coquilles; offrant leurs figures, leur arrangement générique, leurs descriptions caractéristiques, leurs noms; ainsi que leur synonymie en plusieurs langues. Ouvrage destiné à faciliter l'étude des coquilles, ainsi que leur disposition dans les cabinets d'histoire naturelle. Coquilles univalves, non cloisonnées. Tome second.* Schoell, Paris, 676 pp. <http://www.biodiversitylibrary.org/item/41566#page/9/mode/1up>
- Morelet A (1853) Catalogue des Mollusques terrestres et fluviatiles de l'Algérie. *Journal de Conchyliologie* 4: 280–305. <http://www.biodiversitylibrary.org/item/53861#page/286/mode/1up>
- Morelet A (1857) Testacea nova Australiae. *Bulletin de la Société d'Histoire Naturelle du Département de la Moselle* 8: 26–33. <http://www.biodiversitylibrary.org/item/105437#page/38/mode/1up>
- Morelet A (1864) Descriptions de Coquilles inédites. *Journal de Conchyliologie* 12: 155–159. <http://biodiversitylibrary.org/item/54459#page/163/mode/1up>
- Morelet A (1880) La Faune malacologique du Maroc en 1880. *Journal de Conchyliologie* 28: 5–83. <http://biodiversitylibrary.org/item/54068#page/11/mode/1up>
- Moricand S (1838) Premier supplément au Mémoire sur les Coquilles terrestres et fluviatiles de la province de Bahia, envoyées par M. Blanchet. *Memoires de la Société de physique et d'histoire naturelle de Genève* 8: 139–148. <http://www.biodiversitylibrary.org/item/39750#page/175/mode/1up>
- Moricand S (1841) Second supplément au Mémoire sur les Coquilles terrestres et fluviatiles de la province de Bahia, envoyées par M. Blanchet. *Memoires de la Société de phy-*

- sique et d'histoire naturelle de Genève 9: 57–64. <http://www.biodiversitylibrary.org/item/39756#page/79/mode/1up>
- Morlet L (1881) Liste des coquilles recueillies par M. le Dr André pendant l'expédition des Chotts et description d'espèces nouvelles. Archives des missions scientifiques et littéraires, troisième série 7: 393–397. <http://www.biodiversitylibrary.org/item/111285#page/411/mode/1up>
- Mousson A (1854) Coquilles terrestres et fluviatiles recueillies par M. le prof Bellardi dans un voyage en Orient. Zurcher et Furrer, Zürich, 59 pp. doi: 10.5962/bhl.title.101714
- Mousson A (1861) Coquilles terrestres et fluviatiles recueillies par M. le Prof. J. R. Roth dans son dernier voyage en Orient. Vierteljahrsschrift der Naturforschenden Gesellschaft in Zürich 6: 1–34, 124–156. <http://www.biodiversitylibrary.org/item/35456#page/463/mode/1up>
- Mousson A (1863) Coquilles terrestres et fluviatiles, recueillies dans l'Orient par M. le Dr. Alex. Schläfli. Vierteljahrsschrift der Naturforschenden Gesellschaft in Zürich 8: 275–320, 368–426. <http://www.biodiversitylibrary.org/item/34519#page/733/mode/1up>
- Mousson A (1874) Coquilles terrestres et fluviatiles recueillies par M. le Dr Alex. Schläfli en Orient. Journal de Conchyliologie 22: 5–60. <http://biodiversitylibrary.org/item/53430#page/9/mode/1up>
- Munier-Chalmas C-P-E (1897) Note préliminaire sur les assises montiennes du bassin de Paris. Bulletin de la Société Géologique de France, troisième série 25: 82–91. http://jubilotheque.upmc.fr/img-viewer/fonds-bulsgf/GB_000071_001/Contenu/JPEG_HD/viewer.html?ns=GB_000071_001_J3_0087.jpg
- Münster G (1841) Beiträge zur Petrefacten-Kunde. Buchner'sche Buchhandlung, Bayreuth, 152 pp. https://books.google.at/books?id=k8MyAQAAMAAJ&printsec=frontcover&hl=de&source=gbs_ge_summary_r&cad=0#v=onepage&q&f=false
- Neubauer TA, Georgopoulou E, Kroh A, Harzhauser M, Mandic O, Esu D (2015a) Synopsis of European Neogene freshwater gastropod localities: updated stratigraphy and geography. *Palaeontologia Electronica* 18 (1): 3T (7 p.). <http://palaeo-electronica.org/content/2015/1153-neogene-freshwater-gastropods>
- Neubauer TA, Harzhauser M, Georgopoulou E, Mandic O, Kroh A (2014a) Replacement names and nomenclatural comments for problematic species-group names in Europe's Neogene freshwater Gastropoda. *Zootaxa* 3785 (3): 453–468. doi: 10.11646/zootaxa.3785.3.7
- Neubauer TA, Harzhauser M, Georgopoulou E, Wrozyńska C (2014d) Population bottleneck triggering millennial-scale morphospace shifts in endemic thermal-spring melanopsids. *Palaeogeography, Palaeoclimatology, Palaeoecology* 414: 116–128. doi: 10.1016/j.palaeo.2014.08.015
- Neubauer TA, Harzhauser M, Kroh A, Georgopoulou E, Mandic O (2014c) Replacement names and nomenclatural comments for problematic species-group names in Europe's Neogene freshwater Gastropoda. Part 2. *ZooKeys* 429: 13–46. doi: 10.3897/zookeys.429.7420
- Neubauer TA, Harzhauser M, Mandic O, Georgopoulou E, Kroh A (2016a) Paleobiogeography and historical biogeography of the non-marine caenogastropod family Melanopsidae. *Palaeogeography, Palaeoclimatology, Palaeoecology* 444: 124–143. doi: 10.1016/j.palaeo.2015.12.017

- Neubauer TA, Harzhauser M, Pipík R (2015b) Upper Miocene endemic lacustrine gastropod fauna of the Turiec Basin: addressing taxonomic, paleobiogeographic and stratigraphic issues. *Geologica Carpathica* 66 (2): 139–156. doi: 10.1515/geoca-2015-0016
- Neubauer TA, Mandić O, Harzhauser M (2011) Middle Miocene Freshwater Mollusks from Lake Sinj (Dinaride Lake System, SE Croatia; Langhian). *Archiv für Molluskenkunde* 140 (2): 201–237. doi: 10.1127/arch.moll/1869-0963/140/201-237
- Neubauer TA, Mandić O, Harzhauser M (2014b) A new melanopsid species from the Middle Miocene Kupres Basin (Bosnia and Herzegovina). *The Nautilus* 128 (2): 51–54.
- Neubauer TA, Mandić O, Harzhauser M (2016b) The freshwater mollusk fauna of the Middle Miocene Lake Drniš (Dinaride Lake System, Croatia): a taxonomic and systematic revision. *Austrian Journal of Earth Sciences* 108 (2): 15–67. doi: 10.17738/ajes.2015.0013
- Neubauer TA, Mandić O, Harzhauser M (2016c) The early Middle Miocene lacustrine gastropod fauna of Džepi, Bosnia and Herzegovina (Dinaride Lake System): high endemism in a small space. *Bulletin of Geosciences* 91 (2): 271–296. doi: 10.3140/bull.geosci.1584
- Neubauer TA, Mandić O, Harzhauser M, Hrvatić H (2013) A new Miocene lacustrine mollusc fauna of the Dinaride Lake System and its palaeobiogeographic, palaeoecologic, and taxonomic implications. *Palaeontology* 56 (1): 129–156. doi: 10.1111/j.1475-4983.2012.01171.x
- Neumayr M (1869) II. Beiträge zur Kenntniss fossiler Binnenfaunen. *Jahrbuch der k. k. geologischen Reichsanstalt* 19: 355–382. <http://www.biodiversitylibrary.org/item/111699-#page/407/mode/1up>
- Neumayr M (1876) Ueber einige neue Vorkommnisse von jungtertiären Binnenmollusken. *Verhandlungen der k. k. geologischen Reichsanstalt* 1876 (16): 366–368. <http://www.biodiversitylibrary.org/item/120885#page/756/mode/1up>
- Neumayr M (1880a) Der geologische Bau des westlichen Mittel-Griechenland. *Denkschriften der mathematisch-naturwissenschaftlichen Classe der Kaiserlichen Akademie der Wissenschaften* 40 (1): 91–128. <http://www.biodiversitylibrary.org/item/104344#page/114/mode/1up>
- Neumayr M (1880b) Über den geologischen Bau der Insel Kos und über die Gliederung der jungtertiären Binnenablagerungen des Archipels. *Denkschriften der mathematisch-naturwissenschaftlichen Classe der Kaiserlichen Akademie der Wissenschaften* 40 (1): 213–314. <http://www.biodiversitylibrary.org/item/104344#page/243/mode/1up>
- Neumayr M (1880c) V. Tertiäre Binnenmollusken aus Bosnien und der Hercegovina. *Jahrbuch der k. k. geologischen Reichsanstalt* 30 (2): 463–486. <http://www.biodiversitylibrary.org/item/110179#page/491/mode/1up>
- Neumayr M, Paul CM (1875) Die Congerien- und Paludien-schichten Slavoniens und deren Faunen. *Abhandlungen der k. k. Geologischen Reichsanstalt* 7 (3): 1–106.
- Nevill G (1884) Hand list of Mollusca in the Indian Museum, Calcutta. Part II. Gastropoda. Prosobranchia-Neurobranchia (contd.). Printed by order of the trustees, Calcutta, 306 pp. <http://www.biodiversitylibrary.org/item/65259#page/7/mode/1up> (the date “1885” is printed in the preface, but at the end of the volume on p. 306 the date is given as “14-8-84”; perhaps the preface was added later)

- Newton RB (1891) Systematic list of the Frederick E. Edwards collection of British Oligocene and Eocene Mollusca in the British Museum (Natural History), with references to the type-specimens from similar horizons contained in other collections belonging to the Geological Department of the Museum. Printed by order of the Trustees, London, xxviii + 365 pp. <http://www.biodiversitylibrary.org/item/65563#page/5/mode/1up>
- Nicolas H (1898) Mollusques nouveaux. *Comptes Rendus de l'Association Francaise pour l'Avancement des Sciences* 26 (2): 528–537. <http://gallica.bnf.fr/ark:/12148/bpt-6k201187c/f529.image>
- Noetling F (1886) Ueber die Lagerungsverhältnisse einer quartären Fauna im Gebiete des Jordansthal. *Zeitschrift der Deutschen Geologischen Gesellschaft* 38: 807–823. <http://www.biodiversitylibrary.org/item/141361#page/817/mode/1up>
- Noulet J-B (1857) Coquilles fossiles nouvelles des terrains d'eau douce du Sud-Ouest de la France. Editions Masson, Paris, 24 pp.
- Nuttall CP (1990) A review of the Tertiary non-marine molluscan faunas of the Pebasian and other inland basins of north-western South America. *Bulletin of The British Museum (Natural History) Geology* 45 (2): 165–371.
- Olivier GA (1801) Voyage dans l'Empire Othoman, l'Égypte et la Perse, fait par ordre du gouvernement, pendant les six premières années de la République. Tome premier. Agasse, Paris, 432 pp. http://gdz.sub.uni-goettingen.de/dms/load/img/?PPN=PPN625322428&DMDID=&LOGID=LOG_0007&PHYSID=PHYS_0008
- Olivier GA (1804) Voyage dans l'Empire Othoman, l'Égypte et la Perse, fait par ordre du gouvernement, pendant les six premières années de la République. Tome second. Agasse, Paris, 466 pp. http://gdz.sub.uni-goettingen.de/dms/load/img/?PPN=PPN625322681&DMDID=&LOGID=LOG_0005&PHYSID=PHYS_0008
- Olsson AA (1929) Contributions to the Tertiary Paleontology of northern Peru: Part 2, Upper Eocene Mollusca and Brachiopoda. *Bulletins of American Paleontology* 15 (57): 69–117. <http://www.biodiversitylibrary.org/item/95850#page/83/mode/1up>
- Olujčić J (1999) O razvojnim nizovima nekoliko melanopsida i prozostenida iz sarmatskih naslaga okolice Sinja (Dalmacija, Hrvatska). *Hrvatski prirodoslovni muzej / Provincijalat Franjevačke provincije presvetog Otkupitelja, Zagreb / Sinj*, 7–32; 33–60 pp.
- Oppenheim P (1890a) Die Land- und Süßwasserschnecken der Vicentiner Eocänbildungen. Eine paläontologisch-zoographische Studie. *Denkschriften der mathematisch-naturwissenschaftlichen Classe der Kaiserlichen Akademie der Wissenschaften* 57: 113–150. <http://www.biodiversitylibrary.org/item/31471#page/151/mode/1up>
- Oppenheim P (1890b) Neue oder wenig gekannte Binnenschnecken des Neogen im Peloponnes und im südlichen Mittel-Griechenland. *Zeitschrift der Deutschen Geologischen Gesellschaft* 42 (3): 588–592. <http://www.biodiversitylibrary.org/item/37733#page/598/mode/1up>
- Oppenheim P (1891) Beiträge zur Kenntniss des Neogen in Griechenland. *Zeitschrift der Deutschen Geologischen Gesellschaft* 43: 421–487. <http://www.biodiversitylibrary.org/item/141359#page/429/mode/1up>
- Oppenheim P (1892) Ueber einige Brackwasser- und Binnenmollusken aus der Kreide und dem Eocän Ungarns. *Zeitschrift der Deutschen Geologischen Gesellschaft* 44: 697–818. <http://www.biodiversitylibrary.org/item/141293#page/713/mode/1up>

- Pacaud J-M, Harzhauser M (2012) *Jponsia*, *Moniquia* et *Eginea*, trois nouveaux genres de Pachychilidae (Gastropoda, Caenogastropoda) du Paléogène européen. *Annales du Muséum d'histoire naturelle de Nice* 27: 105–153.
- Pacaud J-M, Le Renard J (1995) Révision des Mollusques paléogènes du bassin de Paris. IV - Liste systématique actualisée. *Cossmanniana* 3 (4): 151–187.
- Paetel F (1888) *Catalog der Conchylien-Sammlung, mit Hinzufügung der bis jetzt publicirten recenten Arten, sowie der ermittelten Synonyma. Erste Abtheilung: Die Cephalopoden, Pteropoden und Meeres-Gastropoden.* Gebrüder Paetel, Berlin, 639 pp. <http://biodiversitylibrary.org/item/120992#page/7/mode/1up>
- Paladilhe A (1874) Description de quelques nouvelles espèces de coquilles fossiles provenant des marnes pleistocènes d'estuaire des environs d'Oran. *Revue des Sciences Naturelles* 3: 399–405. <http://www.biodiversitylibrary.org/item/52514#page/419/mode/1up>
- Paladilhe A (1875) Étude sur les coquilles terrestres et fluviatiles rapportées du Maroc par le Dr. Bleicher. *Revue et Magasin de Zoologie pure et appliquée, troisième série* 3: 75–101. <http://www.biodiversitylibrary.org/item/103201#page/83/mode/1up>
- Pallary P (1899) Deuxième contribution à l'étude de la Faune malacologique du Nord-Ouest de l'Afrique. *Journal de Conchyliologie* 46: 49–170. <http://biodiversitylibrary.org/item/54071#page/57/mode/1up>
- Pallary P (1901a) Sur les mollusques fossiles terrestres, fluviatiles et saumâtres de l'Algérie. *Mémoires de la Société Géologique de France, Paléontologie, Mémoire* 22 9 (1): 1–213. <http://www.biodiversitylibrary.org/item/133868#page/11/mode/1up>
- Pallary P (1901b) Diagnoses de quelques coquilles nouvelles provenant du Maroc. *Journal de Conchyliologie* 49 (3): 226–228. <http://biodiversitylibrary.org/item/53864#page/238/mode/1up>
- Pallary P (1904) Quatrième contribution à l'étude de la Faune malacologique du Nord-Ouest de l'Afrique. *Journal de Conchyliologie* 52: 5–58. <http://biodiversitylibrary.org/item/55042#page/13/mode/1up>
- Pallary P (1911) Description de quelques *Melanopsis* nouveaux ou peu connus du Maroc. *Bulletin de la Société d'Histoire Naturelle de l'Afrique du Nord* 2 (8): 127–134. <http://bibdigital.rjb.csic.es/spa/Libro.php?Libro=1833> (first page of volume)
- Pallary P (1912) Étude sur quelques *Melanopsis* du Sahara et de la Tunisie. *Bulletin de la Société d'Histoire Naturelle de l'Afrique du Nord* 3 (1): 14–22. <http://bibdigital.rjb.csic.es/spa/Libro.php?Libro=1834> (first page of volume)
- Pallary P (1913) Description de quelques Mollusques terrestres nouveaux du Sud du Maroc. *Bulletin du Muséum National d'Histoire Naturelle* 19 (6): 360–365. <http://www.biodiversitylibrary.org/item/27226#page/418/mode/1up>
- Pallary P (1915) Description de quelques Mollusques nouveaux du Grand Atlas. *Bulletin du Muséum National d'Histoire Naturelle* 21 (1): 21–28. <http://www.biodiversitylibrary.org/item/27185#page/37/mode/1up>
- Pallary P (1916) Observations relatives à la nomenclature des *Melanopsis* fossiles. *Bulletin de la Société d'Histoire Naturelle de l'Afrique du Nord* 7 (3): 70–87. <http://bibdigital.rjb.csic.es/spa/Libro.php?Libro=1840> (first page of volume)

- Pallary P (1918) Diagnoses d'une cinquantaine de mollusques terrestres nouveaux du Nord de l'Afrique. Bulletin de la Société d'Histoire Naturelle de l'Afrique du Nord 9 (7): 137–152. <http://bibdigital.rjb.csic.es/spa/Libro.php?Libro=1842> (first page of volume)
- Pallary P (1920a) Descriptions d'une nouvelle cinquantaine de mollusques terrestres nouveaux du Nord-Ouest de l'Afrique. Bulletin de la Société d'Histoire Naturelle de l'Afrique du Nord 11 (2): 18–34. <http://bibdigital.rjb.csic.es/spa/Libro.php?Libro=1844> [first page of volume]
- Pallary P (1920b) Deuxième note sur la nomenclature des *Melanopsis* fossiles. Bulletin de la Société d'Histoire Naturelle de l'Afrique du Nord 11 (7): 104–119. <http://bibdigital.rjb.csic.es/spa/Libro.php?Libro=1844> (first page of volume)
- Pallary P (1920c) Récoltes malacologiques du Capitaine Paul Martel dans la partie septentrionale du Maroc. Journal de Conchyliologie 65 (1–2): 1–39, 131–160. <http://biodiversitylibrary.org/item/54318#page/15/mode/1up>
- Pallary P (1922) Faune malacologique du Grand Atlas (Suite). Journal de Conchyliologie 66 (3): 185–217. <http://gallica.bnf.fr/ark:/12148/bpt6k5571699n/f3.item>
- Pallary P (1923) Faune malacologique des eaux douces de la Tunisie. Archives de l'Institut Pasteur de l'Afrique du Nord 3 (1): 22–47.
- Pallary P (1924) Révision des *Melanopsis* de l'Espagne. Bulletin de la Société d'Histoire Naturelle de l'Afrique du Nord 15 (6): 240–255. <http://bibdigital.rjb.csic.es/spa/Libro.php?Libro=1849> (first page of volume)
- Pallary P (1925) Troisième note relative à la nomenclature des *Melanopsis* fossiles. Bulletin de la Société d'Histoire Naturelle de l'Afrique du Nord 16 (8): 256–258. <http://bibdigital.rjb.csic.es/spa/Libro.php?Libro=1853> (first page of volume)
- Pallary P (1926a) Répertoire des *Melanopsis* fossiles et vivantes connus en 1925. Bulletin de la Société d'Histoire Naturelle de l'Afrique du Nord 17 (2–3): 73–93, 126–136. <http://bibdigital.rjb.csic.es/spa/Libro.php?Libro=1854> (first page of volume)
- Pallary P (1926b) Supplément rectificatif au répertoire des *Melanopsis*. Bulletin de la Société d'Histoire Naturelle de l'Afrique du Nord 17 (8): 236–237. <http://bibdigital.rjb.csic.es/spa/Libro.php?Libro=1854> (first page of volume)
- Pallary P (1926c) Sur une faunule aquatique pliocénique d'Oran. Bulletin de la Société d'Histoire Naturelle de l'Afrique du Nord 17 (9): 284–289. <http://bibdigital.rjb.csic.es/spa/Libro.php?Libro=1854> (first page of volume)
- Pallary P (1928a) Complément à la faune malacologique de la Berbérie (suite). Journal de Conchyliologie 71 (3): 197–277.
- Pallary P (1928b) Notice sur seize mollusques nouveaux du Maroc découverts en 1926 et 1927. Journal de Conchyliologie 72 (1): 1–24. <http://gallica.bnf.fr/ark:/12148/bpt6k5566998j/f8.item>
- Pallary P (1930) Mollusques aquatiques nouveaux du Levant. Bulletin du Muséum National d'Histoire Naturelle, deuxième série 2 (3): 286–290. <http://gallica.bnf.fr/ark:/12148/bpt-6k5532113m/f44.image>
- Pallary P (1933) Diagnoses de quinze mollusques continentaux du Maroc. Bulletin de la Société d'Histoire Naturelle de l'Afrique du Nord 24 (7): 243–248. <http://bibdigital.rjb.csic.es/spa/Libro.php?Libro=1891> (first page of volume)

- Pallary P (1936) Deuxième complément à la faune malacologique de la Berbérie. *Journal de Conchyliologie* 80 (1): 5–65. <http://gallica.bnf.fr/ark:/12148/bpt6k5568411h/f12.image>
- Pallary P (1939) Deuxième addition à la faune malacologique de la Syrie. *Mémoires de l'Institut d'Égypte* 39: 1–141.
- Palmer KvW, Brann DC (1966) Catalogue of the Paleocene and Eocene Mollusca of the southern and eastern United States. Part II. Gastropoda (excluding Pteropoda, Part I). *Bulletins of American Paleontology* 48 (218): 471–1057. <http://www.biodiversitylibrary.org/item/40262#page/9/mode/1up>
- Pană I (2003) Les nannogastropodes. In: Papaianopol I, Marinescu F, Krstić N, Macalet R (eds) *Chronostratigraphie und Neostatotypen. Neogen der Zentrale Paratethys, Bd. X, Pl2. Romanien*. Editura Academiei Române, Bucuresti, 296–349.
- Pană I, Enache C, Andreescu I (1981) Fauna de moluște a depozitelor cu ligniți din Oltenia. Institutul de cercetări, inginerie tehnologică și proiectări miniere pentru lignit, Craiova, 276 pp.
- Pantaneli D (1879) Sugli strati miocenici del Casino (Siena) e considerazioni sul miocene superiore. *Atti della Reale Accademia dei Lincei, Ann. 276, Memorie, Ser. 3* 3: 309–327.
- Pantaneli D (1886a) Monografia degli strati pontici del Miocene superiore nella Italia settentrionale e centrale. *Memorie della Reale Accademia di Scienze, Lettere ed Arti di Modena, Ser. 2a* 4: 127–231.
- Pantaneli D (1886b) *Melanopsis* fossili e viventi d'Italia. *Bollettino della Società Malacologica Italiana* 12: 63–82. <http://www.biodiversitylibrary.org/item/125693#page/71/mode/1up>
- Papp A (1947) Brack- und Süßwasserfaunen Griechenlands. 3. Über Lebensspuren an Süßwassergastropoden. *Annales Géologiques des Pays Helléniques* 1: 126–128.
- Papp A (1951) Das Pannon des Wiener Beckens. *Mitteilungen der Geologischen Gesellschaft in Wien* 39–41: 99–193. http://www.zobodat.at/pdf/MittGeolGes_39_41_0099-0193.pdf
- Papp A (1953a) Brack- und Süßwasserfaunen Griechenlands. IV. Süßwassermollusken aus dem Pliozän von Elis (Peloponnes). *Annales Géologiques des Pays Helléniques* 5: 107–113.
- Papp A (1953b) Die Molluskenfauna des Pannon des Wiener Beckens. *Mitteilungen der geologischen Gesellschaft in Wien* 44: 85–222. http://www.zobodat.at/pdf/MittGeolGes_44_0085-0222.pdf
- Papp A (1955) Brack- u. Süßwasserfaunen Griechenlands. V. Bemerkungen über Melanopsiden der Untergattung *Melanosteira* Oppenheim, 1891. *Annales Géologiques des Pays Helléniques* 6: 122–132.
- Papp A (1979) Zur Kenntnis neogener Süßwasserfaunen in Attika (Griechenland). *Annales Géologiques des Pays Helléniques* 29: 664–678.
- Papp A, Psarianos P (1955) Über einige Süßwassergastropoden aus pliozänen Ablagerungen Griechenlands. *Annales Géologiques des Pays Helléniques* 6: 145–150.
- Papp A, Thenius E (1952) Paläontologische Ergebnisse einer mit Unterstützung der Akademie der Wissenschaften durchgeführten Griechenlandfahrt 1951. *Anzeiger der Österreichischen Akademie der Wissenschaften, mathematisch-naturwissenschaftliche Klasse* 10: 100–110.
- Pashko P (1968) Molusquet e suitës së Burrelit. *Përmbledhje studimesh* 9–10: 133–152.

- Pauč M (1937) Les Mollusques pléistocènes de Băile Episcopoești. Bulletin de la Société Roumaine de Géologie 3: 130–142.
- Pavlović PS (1903) Građa za poznavanje tercijara u Staroj Srbiji. Annales Géologiques de la Péninsule Balkanique 6 (1): 155–189. [in Serbian; a German translation was published in 1911 in the same journal, volume 6 (2)] <https://archive.org/stream/geoloshkianalib-00zavogoog#page/n176/mode/2up>
- Pavlović PS (1922) Prilozi za poznavanje Tercijara u Srbiji. Annales Géologiques de la Péninsule Balkanique 7 (1): 42–50. [In Serbian]
- Pavlović PS (1927) Donjopontiski mekušci iz okoline Beograda (s naročitim obzirom na fosilnu faunu okoline sela Vrčina). Sprska Akademija nauka, posebna izdanja 66 (Prirodnjački i matematički spisi 17): 1–121. [in Serbian; a French translation with a reduced text was published in 1928 in the Annales Géologiques de la Péninsule Balkanique 9 (2): 1–74]
- Pavlović PS (1931) O fosilnoj fauni mekusača iz Skopske Kotline. Glasnik Skopskog naucnog društva, Odeljenje prirodnih Nauka 9 (3): 1–28. [In Serbian]
- Pavlović PS (1932) Novi prilozi za poznavanje fosilne faune iz Kosovske i Metohiskopodrimske Kotline. Bulletin du Service Géologique du Royaume de Yougoslavie 1 (2): 231–253. [in Serbian, with French summary]
- Pecchioli V (1864) Descrizione di alcuni nuovi fossili delle argille subappennine Toscane. Atti della Società Italiana di Scienze Naturali 6: 498–529. <http://www.biodiversitylibrary.org/item/123039#page/506/mode/1up>
- Penecke KA (1884) Beiträge zur Kenntniss der Fauna der slavonischen Paludinschichten. II. *Congerina*, *Pisidium*, *Cardium* und die Gasteropoden. Beiträge zur Paläontologie Österreich-Ungarns und des Orients 4: 15–44. <http://www.biodiversitylibrary.org/item/50742#page/27/mode/1up>
- Penecke KA (1885) Das Eocän des Krappfeldes in Kärnten. Sitzungsberichte der Akademie der Wissenschaften, mathematisch-naturwissenschaftliche Classe 90: 327–371. <http://www.biodiversitylibrary.org/item/31594#page/373/mode/1up>
- Pérès J-M (1938a) Aperçu sur les variations de quelques *Melanopsis* marocains. Journal de Conchyliologie 82: 312–321.
- Pérès J-M (1938b) Sur quelques *Melanopsis* du Maroc. Bulletin de la Société Zoologique de France 63 (2): 127–133.
- Pérès J-M (1939) Contribution à l'étude des *Melanopsis* du Maroc. Journal de Conchyliologie 83: 129–162.
- Pérès J-M (1946) Contribution à l'étude du genre *Melanopsis*. Journal de Conchyliologie 86 (3–4): 109–136, 137–174.
- Pezant A (1908) Mollusques fossiles de Monneville (Oise). La Feuille des Jeunes Naturalistes, quatrième série 8 (451–455): 132–138, 158–161, 173–178, 198–202, 224–229. <http://www.biodiversitylibrary.org/item/137703#page/152/mode/1up>
- Pfeiffer C (1828) Naturgeschichte deutscher Land- und Süsswasser-Mollusken. Dritte Abtheilung. Landes-Industrie-Comptoir, Weimar, vi + 84 pp. https://books.google.at/books?id=UtrcAAAAcAAJ&printsec=frontcover&hl=de&source=gbg_ge_summary_r&cad=0#v=onepage&q&f=false

- Philippi RA (1845) Abbildungen und Beschreibungen neuer oder wenig gekannter Conchylien. Erster Band. Theodor Fischer, Cassel, ii + 212 pp. <http://www.biodiversitylibrary.org/item/41582#page/9/mode/1up>
- Philippi RA (1847) Abbildungen und Beschreibungen neuer oder wenig gekannter Conchylien. Zweiter Band. Theodor Fischer, Cassel, ix + 231 pp. <http://www.biodiversitylibrary.org/item/41581#page/7/mode/1up>
- Philippi RA (1887) Die tertiären und quartären Versteinerungen Chiles. F. A. Brockhaus, Leipzig, 266 pp. doi: 10.5962/bhl.title.15048
- Picard L (1934) Mollusken der levantinischen Stufe Nordpalästinas (Jordantal). Archiv für Molluskenkunde 66 (3): 105–139.
- Pichler A (1856) Zur Geognosie der nordöstlichen Kalkalpen Tirols. Jahrbuch der k. k. geologischen Reichsanstalt 7: 717–738. http://opac.geologie.ac.at/wwwopacx/wwwopac.ashx?command=getcontent&server=images&value=JB0074_717_A.pdf
- Poiret JLM (1801) Coquilles fluviatiles et terrestres observées dans le Département de l'Aisne et aux environs de Paris. Théophile Barrois; Soissons, chez l'auteur, Paris, 119 pp. <http://www.biodiversitylibrary.org/item/81286#page/9/mode/1up>
- Ponder WF (1984) A review of the genera of the Irvadiidae (Gastropoda: Rissoacea) with an assessment of the relationships of the family. Malacologia 25: 21–71. <http://biodiversitylibrary.org/page/13145132#page/31/mode/1up>
- Popescu-Voitești I (1910) Contribution à l'étude stratigraphique du Nummulitique de la Dépression Gétique (Roumanie occidentale). Anuarul Institutului Geologic al României 3 (2): 275–394.
- Porumbaru RC (1881) Étude géologique des environs de Craiova, parcours Bucovatzu-Cretzes-ci. Première partie. Gauthier-Villars, Imprimeur-Libraire, Paris, 42 pp.
- Potiez V-L-V, Michaud A-L-G (1838) Galerie des mollusques, ou catalogue méthodique, descriptif et raisonné des mollusques et coquilles du Muséum de Douai. Tome premier. Baillière, Paris, Londres, xxxvi + 560 pp. <http://www.biodiversitylibrary.org/item/41188#page/9/mode/1up>
- Preston HB (1913) New minute Terrestrial and Aquatic Mollusca from the Dutch East Indian Island of Beilan-Beilan, with Descriptions of Four new Genera and Subgenera. The Annals and Magazine of Natural History, eighth series 12 (71): 432–439. <http://www.biodiversitylibrary.org/item/71910#page/444/mode/1up>
- Preston HB (1914) A Molluscan Faunal List of the Lake of Tiberias, with Descriptions of new Species. Journal & Proceedings of the Asiatic Society of Bengal, new series 9: 465–475. <http://biodiversitylibrary.org/page/31620706#page/497/mode/1up>
- [Prevost C] (1821) Sur une nouvelle espèce de mollusque testacé du genre Mélanopside. Bulletin des Sciences, par la Société philomatique de Paris 1821: 136–137. <http://www.biodiversitylibrary.org/item/99191#page/346/mode/1up>
- Quaggiotto E, Mellini A (2008) Catalogo aggiornato dei molluschi fossili eocenici di San Giovanni Ilarione (Verona - Italia settentrionale). Prima parte: Mollusca, Gastropoda. Studi e Ricerche - Associazione Amici del Museo - Museo Civico "G. Zannato" 15: 41–58.
- Ramírez R, Paredes C, Arenas J (2003) Moluscos del Perú. Revista de Biología Tropical Universidad de Costa Rica 51 (Suppl. 3): 225–284. <http://www.ots.ac.cr/rbt/attachments/suppls/sup51-3%20malacol/12-Ramirez-Mol-225-284.pdf>

- Reeve LA (1860) Monograph of the genus *Melanopsis*. In: Reeve LA (Ed.) Conchologia iconica: or, Illustrations of the shells of molluscous animals. Vol. XII. Privately published by author, London, pls. 1–3 and unpaginated text. <https://archive.org/stream/conchologiaiconi12reev#page/n537/mode/2up>
- Reid DG (2014) The genus *Cerithidea* Swainson, 1840 (Gastropoda: Potamididae) in the Indo-West Pacific region. *Zootaxa* 3775 (1): 3–65. doi: 10.11646/zootaxa.3775.1.1
- Robles F (1975) El género “*Melanopsis*” Férussac en el Neógeno continental de la cuenca del río Júcar (1.a nota). *Real Sociedad Española de Historia Natural, Vol. Extraordinario Primer Centenario* (1971) 1: 357–369.
- Röding PF (1798) Museum Boltenianum, sive, Catalogus cimeliorum e tribus regnis naturae quae olim collegerat Joa. Fried. Bolten. Pars Secunda continens Conchylia sive Testacea univalvia, bivalvia & multivalvia. Johann Christian Trappe, Hamburg, 199 pp. [In German] <http://www.biodiversitylibrary.org/item/41571#page/13/mode/1up> (reprint from 1906)
- Rolle F (1858) Über die geologische Stellung der Sotzka-Schichten in Steiermark. *Sitzungsberichte der mathematisch-naturwissenschaftlichen Classe der Kaiserlichen Akademie der Wissenschaften* 30: 3–33. <http://www.biodiversitylibrary.org/item/30198#page/13/mode/1up>
- Rolle F (1860) Die Lignit-Ablagerung des Beckens von Schönstein in Unter-Steiermark und ihre Fossilien. *Sitzungsberichte der mathematisch-naturwissenschaftlichen Classe der Kaiserlichen Akademie der Wissenschaften* 41 (1): 7–46. <http://www.biodiversitylibrary.org/item/30221#page/19/mode/1up>
- Rosen O (1914) Katalog der schalentragenden Mollusken des Kaukasus. *Izvestija Kavkazskago Muzeja* 6: 141–253.
- Rossmässler EA (1838–1844) Iconographie der Land- und Süsswasser-Mollusken, mit vorzüglicher Berücksichtigung der europäischen noch nicht abgebildeten Arten. 2. Band. Arnold, Dresden, Leipzig. Hefte 1–2 (7/8): 43 pp. (1838); Hefte 3–4 (9/10): 46 pp. (1839); Heft 4 (11): 15 pp. (1842); Heft 6 (12): 37 pp. (1844). <http://www.biodiversitylibrary.org/item/80434#page/325/mode/1up>
- Rossmässler EA (1854–1859) Iconographie der Land- und Süsswassermollusken Europa's, mit vorzüglicher Berücksichtigung kritischer und noch nicht abgebildeter Arten. 3. Band. Hermann Costenoble, Leipzig. Hefte 1–2 (13/14): pp. 1–39 (1854), Hefte 3–4 (15/16): pp. 41–77 (1856); Hefte 5–6 (17/18): pp. 81–140 (1859). <http://www.biodiversitylibrary.org/item/55211#page/9/mode/1up>
- Roth JR (1839) Molluscorum species, quas in itinere per orientem facto comites clariss. Schuberti Doctores M. Erdl et J. R. Roth collegerunt. C. Wolf, Monachii, i–viii + 9–26 pp. <http://www.biodiversitylibrary.org/item/41720#page/3/mode/1up>
- Roule L (1884) Description de quelques nouvelles fossiles du calcaire lacustre de Rognac (Bouches-du-Rhône). *Bulletins de la Société malacologique de France* 1: 311–328. <http://www.biodiversitylibrary.org/item/51550#page/329/mode/1up>
- Rovereto G (1899) Prime ricerche sinonimiche sui generi dei Gasteropodi. *Atti della Società Ligustica di Scienze Naturali e Geografiche* 10 (2): 101–110.
- Royo Gómez J (1922) El Mioceno continental ibérico y su fauna malacológica. *Comisión de investigaciones paleontológicas y prehistóricas* 30: 1–230.

- Royo Gómez J (1929) Moluscos del Terciario continental de la provincia de Burgos. Boletín de la Real Sociedad Española de Historia Natural 29: 239–244. <http://bibdigital.rjb.csic.es/ing/Libro.php?Libro=1245> (first page of volume)
- Rzehak A (1883) Beiträge zur Kenntniss der Tertiärformation im ausseralpinen Wiener Becken (Der Grunder Horizont in Mähren). Verhandlungen des naturforschenden Vereins in Brünn 21: 31–49. <http://www.biodiversitylibrary.org/item/148627#page/467/mode/1up>
- Sacco F (1886) Nuove species terziarie di molluschi terrestri e d'acqua dolce e salmastra del Piemonte. Atti della Società Italiana di Scienze Naturali 29: 427–476. <http://www.biodiversitylibrary.org/item/54923#page/443/mode/1up>
- Sacco F (1889) Aggiunte alla fauna malacologica estramarina fossile del Piemonte e della Liguria. Memorie della Reale Accademia delle Scienze di Torino, Serie II 39 (1): 61–98. <http://www.biodiversitylibrary.org/item/138459#page/97/mode/1up>
- Sacco F (1895) I molluschi dei terreni Terziarii del Piemonte e della Liguria. Parte XVIII (Mela-niidae, Littorinidae, Fossaridae, Rissoidae, Hydrobiidae, Paludinidae e Valvatidae). Carlo Clausen, Torino, 52 pp. <http://www.biodiversitylibrary.org/item/46676#page/9/mode/1up>
- Sandberger CLF (1870–1875) Die Land- und Süßwasser-Conchylien der Vorwelt. C. W. Kreidel, Wiesbaden, Livr. 1–3: 1–96 pp., pls 1–12 (1870); Livr. 4–5: 97–160 pp., pls 13–20 (1871); Livr. 6–8: 161–256 pp., pls 21–32 (1872); Livr. 9–10: 257–352 pp., pls 33–36 (1873); Livr. 11: i–viii, 353–1000 pp. (1875).
- Sandberger F (1850) Ueber die geognostische Zusammensetzung der Umgegend von Wiesbaden. Jahrbücher des Vereins für Naturkunde im Herzogthum Nassau 6: 1–27. https://books.google.at/books?id=h_8IAQAIAAJ&printsec=frontcover&hl=de&source=gbs_ge_summary_r&cad=0#v=onepage&q&f=false
- Sandberger F (1851) [Analogie der Land- und Süßwasser-Fauna des Maynzer Becknes und des Mittelmeeres]. Neues Jahrbuch für Mineralogie, Geognosie, Geologie und Petrefaktenkunde 1851: 676. <http://www.biodiversitylibrary.org/item/44254#page/708/mode/1up>
- Sandberger F (1852) Über die Analogieen der fossilen Land- und Süßwasserfauna des Mainzer Becknes mit der lebenden der Mittelmeerländer. Zeitschrift der Deutschen Geologischen Gesellschaft 6: 680–685. <http://www.biodiversitylibrary.org/item/108484#page/692/mode/1up>
- Sandberger F (1886) Bemerkungen über fossile Conchylien aus dem Süßwasserkalke von Leobersdorf bei Wien (Inzersdorfer Schichten). Verhandlungen der k. k. geologischen Reichsanstalt 1886 (14): 331–332. <http://www.biodiversitylibrary.org/item/120934#page/793/mode/1up>
- Sauerzopf F (1952) Beitrag zur Entwicklungsgeschichte des südburgenländischen Pannons. Burgenländische Heimatblätter 14 (1): 1–16.
- Schenk V (1969) Zur Gastropodenfauna und Biofazies der mittleren Gosau (O.Kreide) von Brandenburg in Tirol. Dissertation, München: Ludwig-Maximilians-Universität München (The dissertation constitutes a publication in the sense of the Code because at least 150 identical copies were issued; W. Werner, München, pers. commun.).
- Schlotheim EF von (1820) Die Petrefactenkunde auf ihrem jetzigen Standpunkte durch die Beschreibung seiner Sammlung versteinerter und fossiler Überreste des Thier- und Pflanzenreichs der Vorwelt erläutert. Becker, Gotha, 457 pp.
- Schmidt FJ (1847) Systematisches Verzeichniss der in der Provinz Krain vorkommenden Land- und Süßwasser-Conchylien, mit Angabe der Fund-Orte. Joseph Blasnik, Laibach, 27 pp.

http://digital.staatsbibliothek-berlin.de/werkansicht/?PPN=PPN690062699&PHYSID=PHYS_0005

Schréter Z (1912) Beiträge zur Tektonik des südlichen Teiles des westlichen Krassószörényer Kalkgebirges. Jahresbericht der Königlich ungarischen Geologischen Anstalt 1910: 134–173.

Schréter Z (1975) Tanulmány az alsópleisztocén korú Melanopsidák köréből. Földtani Közlöny 105 (1): 1–22. [In Hungarian]

Schumacher CF (1817) Essai d'un nouveau système des habitations des vers testacés. Schultz, Copenhagen, iv + 288 pp. <http://www.biodiversitylibrary.org/item/81329#page/7/mode/1up>

Schütt H (1986) Mollusken aus jungpliozänen Seesedimenten von Antirrhion in Akarnanien. In: Symeonidis N, Theodorou G, Schütt H, Velitzelos E (eds) Palaeontological and stratigraphical observations in the area of Achaia and Etoloakamania (W-Greece). Annales Géologiques des Pays Helléniques 33: 329–365.

Schütt H (1988a) Über eine relikte *Melanopsis* aus Jordanien (Ein Beitrag zur Kenntnis der Rassenbildung durch geographische Isolation). Annalen des Naturhistorischen Museums in Wien, Serie B 90: 215–219.

Schütt H (1988b) Ergänzungen zur Kenntnis der Molluskenfauna oberpliozäner Süßwasserkonglomerate Syriens. Archiv für Molluskenkunde 118 (4/6): 129–143.

Schütt H, Bilgin FH (1974) Recent Melanopsines of the Aegean. Archiv für Molluskenkunde 104 (1/3): 59–64.

Schütt H, Ortal R (1993) A preliminary correlation between the plio-pleistocene malacofaunas of the Jordan Valley (Israel) and the Orontes Valley (Syria). Zoology in the Middle East 8 (1): 69–112. doi: 10.1080/09397140.1993.10637638

Sedgwick A, Murchison RI (1832) A Sketch of the Structure of the Eastern Alps; with Sections through the Newer Formations on the Northern Flanks of the Chain, and through the Tertiary Deposits of Styria. Transactions of the Geological Society of London, second series 3 (2): 301–420. doi: 10.1144/transgslb.3.2.301

Seninski K (1905) Novyya Dannyya o neogenovykh 'plastach' Yugozapadnogo Zakavkaz'ya. Trudy Obshchestva Estestvoispytateley pri Imperatorskom Yuryevskom Universitete 16: 1–80.

Servain G (1882) Histoire des mollusques acéphales des environs de Francfort. S. Lejay & Cie, Poissy, 68 pp. <http://www.biodiversitylibrary.org/item/41714#page/5/mode/1up>

Servain G (1884) Excursions malacologiques en Bosnie aux environs de Serajewo et aux sources de la Bosna. Annales de Malacologie 1 (4): 341–380. <http://biodiversitylibrary.org/page/16253419#page/379/mode/1up>

Silva e Castro J de (1883) Contributions à la faune malacologique du Portugal. Jornal de ciencias mathematicas, physicas, e naturaes 9 (35): 121–152. <http://www.biodiversitylibrary.org/item/54851#page/135/mode/1up>

Sinzov I (1884) Opisanie novykh i maloissledovannykh form rakovin iz tretichnykh obrazovaniy Novorossii [Description of new and less well known shells from Tertiary deposits of Novorossia]. Zapiski novorossiiskoe obshchestvo estestvoispytateley 9: 1–13. [In Russian]

Sismonda E (1847) Synopsis methodica animalium invertebratorum pedemontii fossilium (exceptis speciebus ineditis), vol. 2. Augustae Taurinorum, Torino, 62 pp.

Șoverth E (1953) Contribuții la studiul Pliocenului superior din Oltenia. Levantinul de la Hurezani. Revista Universității "C.I.Parhon", Seria Științele Naturii 3: 202–211.

- Sowerby J (1815–1818) The mineral conchology of Great Britain; or, Coloured figures and descriptions of those remains of testaceous animals or shells, which have been preserved at various times and depths in the earth. Vol. II. Privately published by author, London, 251 pp., pls. 103–203. <http://www.biodiversitylibrary.org/item/50341#page/5/mode/1up> (for publication dates see Wright and Cleevely 1985)
- Sowerby J (1821–1823) The mineral conchology of Great Britain; or, Coloured figures and descriptions of those remains of testaceous animals or shells, which have been preserved at various times and depths in the earth. Vol. IV. Privately published by author, London, 151 pp., pls. 307–407. <http://www.biodiversitylibrary.org/item/50366#page/7/mode/1up> (for publication dates see Wright and Cleevely 1985)
- Sowerby J (1826–1829) The mineral conchology of Great Britain; or, Coloured figures and descriptions of those remains of testaceous animals or shells, which have been preserved at various times and depths in the earth. Vol. VI. Privately published by author, London, 230 pp., pls. 504–609. <http://www.biodiversitylibrary.org/item/50339#page/5/mode/1up> (for publication dates see Wright and Cleevely 1985)
- Sowerby J (1845) Conchyliologie minéralogique de la Grande Bretagne. Jent & Gassmann, Soleure, 395 pls. https://books.google.at/books?id=icosAAAAQAAJ&printsec=frontcover&hl=de&source=gbs_ge_summary_r&cad=0#v=onepage&q&f=false
- Speyer O (1870) Die Conchylien der Casseler Tertiär-Bildungen. V. *Pyramidella*, *Odontostoma*, *Turbonilla*, *Eulima*, *Niso*, *Nerita*, *Sigaretus*, *Paludina*, *Bithynia*, *Nematura*, *Melanopsis*, *Melania*. Palaeontographica 19 (2): 47–101. <http://www.biodiversitylibrary.org/item/103320#page/67/mode/1up>
- Squires RL (1998) New Information on Morphology, Stratigraphy, and Paleoclimate Implications of the Eocene Brackish-Marine Gastropod *Loxotrema turritum* Gabb, 1868, from the West Coast of the United States. The Veliger 41 (4): 297–313.
- Squires RL (1999) Middle Eocene Brackish-Marine Mollusks from the Matilija Sandstone at Matilija Hot Springs, Ventura County, Southern California. Contributions in Science 480: 1–30.
- Squires RL, Saul LR (1997) Late Cretaceous Occurrences on the Pacific Slope of North America of the Melanopsid Gastropod Genus *Boggsia* Olsson, 1929. The Veliger 40 (3): 193–202.
- Stache G (1880) Die Liburnische Stufe. Verhandlungen der k. k. Geologischen Reichsanstalt 1880 (12): 195–209. <http://www.biodiversitylibrary.org/item/120884#page/603/mode/1up>
- Stache G (1889) Die Liburnische Stufe und deren Grenz-horizonte. Eine Studie über die Schichtenfolgen der cretacisch-eocänen oder protocänen Landbildungsperiode im Bereiche der Küstenländer von Österreich-Ungarn. Abhandlungen der k. k. Geologischen Reichsanstalt 13: 1–170.
- Starmühlner F (1970) Die Mollusken der neukaledonischen Binnengewässer. Cahiers ORSTOM, série Hydrobiologie 4 (3–4): 3–127. <http://www.biodiversitylibrary.org/item/47219#page/258/mode/1up>
- Starmühlner F (1973) Die Gattung *Melanopsis* Ferussac 1807 auf Neukaledonien. Malacologia 14: 242–243. <http://www.biodiversitylibrary.org/item/47219#page/258/mode/1up>
- Starmühlner F (1974) The freshwater gastropods of Ceylon. Bulletin of the Fisheries Research Station, Sri Lanka 25 (1–2): 97–181. http://aquaticcommons.org/17196/1/NARA25_097%20part%201.pdf

- Starobogatov YI, Aleksenko TL, Levina OV (1992) Rody *Fagotia* i *Microcolpia* (Gastropoda Pectinibranchia Melanopsidae) i ikh predstaviteli v sovremennoj faune [The genera *Fagotia* and *Microcolpia* (Gastropoda Pectinibranchia Melanopsidae) and their representatives in the modern fauna]. Byulleten' Moskovskogo Obshchestva Ispytateley Prirody, Otdel biologicheskiiy 97 (3): 57–72. [In Russian]
- Starobogatov YI, Izzatullaev ZI (1985) Vidy *Melanopsis* pustynnykh rayonov vosto-ka Irana [Species of *Melanopsis* from the desert regions of east Iran]. Trudy Zoologicheskogo Instituta Akademya Nauk, SSSR 135: 33–37. [in Russian] <http://malacolog.com/files/,%20%D0%98%D0%B7%D0%B7%D0%B0%D1%82%D1%83%D0%BB%D0%B0%D0%B5%D0%B2,%201985%20Melanopsis%20%D0%98%D1%80%D0%B0%D0%BD%D0%B0.pdf>
- Stefanescu S (1896) Études sur les Terrains tertiaires de Roumanie. Contribution à l'étude des faunes sarmatique, pontique et levantine. Mémoires de la Société Géologique de France, Mémoire 15. Paléontologique 6 (2–3): 1–147.
- Stefanescu S (1897) Contribution à l'étude des faunes Éogène et Néogène de Roumanie. Bulletin de la Société Géologique de France, troisième série 25: 310–314. http://jubilotheque.upmc.fr/img-viewer/fonds-bulsgf/GB_000071_001/Contenu/JPEG_HD/viewer.html?ns=GB_000071_001_J3_0310.jpg
- Stepanović B (1938) Gornji Miocen u Kamenom Potoku kod sela Kumodraža. Annales Géologiques de la Péninsule Balkanique 15: 51–102.
- Stoliczka F (1860) Über eine der Kreideformation angehörige Süßwasserbildung in den nordöstlichen Alpen. Sitzungsberichte der mathematisch-naturwissenschaftlichen Classe der Kaiserlichen Akademie der Wissenschaften 38: 482–496. <http://www.biodiversitylibrary.org/item/30206#page/542/mode/1up>
- Stoliczka F (1862) Beitrag zur Kenntnis der Molluskenfauna der Cerithien- und Inzersdorfer Schichten des ungarischen Tertiärbeckens. Verhandlungen der kaiserlichen und königlichen zoologisch-botanischen Gesellschaft in Wien 12: 529–538. http://www.zobodat.at/pdf/VZBG_12_0529-0538.pdf
- Strausz L (1941) Melanopsisok változékonyága. Földtani Közlöny 71 (4–6): 135–146, 163–170. [In Hungarian]
- Strausz L (1942) Das Pannon des mittleren Westungarns. Annales historico-naturales Musei Nationalis Hungarici. Pars mineralogica, geologica et palaeontologica 35: 1–102.
- Strong EE, Colgan DJ, Healy JM, Lydeard C, Ponder WF, Glaubrecht M (2011) Phylogeny of the gastropod superfamily Cerithioidea using morphology and molecules. Zoological Journal of the Linnean Society 162 (1): 43–89. doi: 10.1111/j.1096-3642.2010.00670.x
- Strong EE, Gargominy O, Ponder WF, Bouchet P (2008) Global diversity of gastropods (Gastropoda; Mollusca) in freshwater. Hydrobiologia 595: 149–166. doi: 10.1007/s10750-007-9012-6
- Suter H (1893) Check-list of the New Zealand Land and Freshwater Mollusca. Transactions and Proceedings of the Royal Society of New Zealand 26: 139–154. http://rsnz.natlib.govt.nz/volume/rsnz_26/rsnz_26_00_001490.html
- Swainson W (1840) A Treatise on Malacology; or, the natural classification of shells and shell fish. Longman, Orme, Brown, Green & Longmans; John Taylor, London, 419 pp. doi: 10.5962/bhl.title.8027

- Syrides GE (1998) Paratethyan mollusc faunas from the Neogene of Macedonia and Thrace, Northern Greece. *Romanian Journal of Stratigraphy* 78: 171–180.
- Tambareau Y, Villatte J (1966) A propos d'un gastropode nouveau, *Melanopsis (Canthidomus) fontanensis* nov. sp. *Bulletin de la Société d'histoire naturelle de Toulouse* 102 (2): 511–516. <http://gallica.bnf.fr/ark:/12148/bpt6k6556256q/f183.item.r=>
- Taner G (1997) Das Pliozän des östlichen Dardanellen-Beckens, Türkei. *Molluskenfauna und Stratigraphie. Annalen des Naturhistorischen Museums in Wien* 98 A: 35–67. http://www.zobodat.at/pdf/ANNA_98A_0035-0067.pdf
- Tausch L von (1886) Über die Fauna der nicht-marinen Ablagerungen der oberen Kreide des Csongerthales bei Ajka im Bakony (Veszprimer Comitatus, Ungarn) und über einige Conchylien der Gosaumergel von Aigen bei Salzburg. *Abhandlungen der k. k. Geologischen Reichsanstalt* 12: 1–32. http://opac.geologie.ac.at/wwwopacx/wwwopac.ashx?command=getcontent&server=images&cvalue=Abh_Bd_12_1.pdf
- Thomä C (1845) Fossile Conchylien aus den Tertiärschichten bei Hochheim und Wiesbaden gesammelt und im naturhistorischen Museum zu Wiesbaden ausgestellt. *Jahrbücher des Vereins für Naturkunde in Nassau* 2: 125–162.
- Tournouër R (1874) Sur les terrains tertiaires supérieurs du bassin de Théziers (Gard) et sur le niveau des *Potamides Basteroti* dans le bassin du Rhône. *Bulletin de la Société Géologique de France, troisième série* 2: 287–307. http://jubilothèque.upmc.fr/img-viewer/fonds-bulsgf/GB_000046_001/Contenu/JPEG_HD/viewer.html?ns=GB_000046_001_J3_293.jpg
- Tournouër R (1875) Diagnose d'espèces nouvelles de Coquilles d'eau douce, recueillies par M. Gorceix dans les terrains tertiaires supérieurs de l'île de Cos. *Journal de Conchyliologie* 23: 76–78. <http://www.biodiversitylibrary.org/item/53453#page/82/mode/1up>
- Tournouër R (1876) Étude sur les fossiles tertiaires recueillis par M. Gorceix dans l'île de Cos, en 1873. *Annales scientifiques de l'École normale supérieure, deuxième série* 5: 445–475. http://archive.numdam.org/article/ASENS_1876_2_5_445_0.pdf [without plates]
- Tournouër R (1877) Description d'une nouvelle espèce fossile de Melanopside, provenant des terrains tertiaires supérieurs de la province de Constantine. *Journal de Conchyliologie* 25: 275–276. <http://www.biodiversitylibrary.org/item/54878#page/289/mode/1up>
- Tournouër R (1879) Conchyliorum fluviatilium fossilium, quae in stratis tertiariis superioribus Rumaniae Dr. Gregorio Stefanescu collegit, novae species. *Journal de Conchyliologie* 27: 261–264. <http://www.biodiversitylibrary.org/item/54087#page/275/mode/1up>
- Tournouër R (1880) Conchyliorum fluviatilium fossilium, in stratis tertiariis superioribus Rumaniae collectorum, novae species. *Journal de Conchyliologie* 28: 96–100. <http://biodiversitylibrary.org/item/54068#page/102/mode/1up>
- Tournouër R (1882) Description d'un nouveau genre de Melanopsinae fossiles des terrains tertiaires supérieurs de l'Algérie. *Journal de Conchyliologie* 30: 59. <http://www.biodiversitylibrary.org/item/54877#page/67/mode/1up>
- Traub F (1938) Geologische und paläontologische Bearbeitung der Kreide und des Tertiärs im östlichen Rupertiwinkel, nördlich von Salzburg. *Paläontographica Abt. A* 88 (1–3): 1–114.
- Tristram HB (1865) Report on the Terrestrial and Fluvial Mollusca of Palestine. *Proceedings of the Zoological Society of London* 1865: 530–545. doi: 10.1111/j.1469-7998.1865.tb02386.x

- Tschapeck H (1881) Einige Süßwasser-Mollusken des Sanngebietes in Untersteiermark. Jahrbücher der Deutschen Malakozoologischen Gesellschaft 8: 101–109. <http://www.biodiversitylibrary.org/item/55035#page/115/mode/1up>
- Van Damme, D. 2014. *Melanopsis nodosa*. The IUCN Red List of Threatened Species 2014: e.T20593629A42422265. doi: 10.2305/IUCN.UK.2014-1.RLTS.T20593629A42422265.en [Accessed on 15 February 2016]
- Vergnau-Saubade AM (1966) Monographie de la faune malacologique des gisements types de l'Oligocène aquitain. Thèse de Doctorat d'État, Bordeaux: Université de Bordeaux (unpublished).
- Vergnau-Saubade AM (1968) Quelques espèces nouvelles de mollusques de l'Oligocène aquitain. Bulletin de l'Institut de Géologie du Bassin d'Aquitaine 4: 196–211.
- Vidal LM (1874) Datos para el conocimiento del terreno garumnense de Cataluña. Boletín de la Comisión del Mapa Geológico de España 1: 209–247. <http://hemerotecadigital.bne.es/pdf.raw?query=parent%3A0005421014+type%3Apress%2Fpage&name=Bolet%C3%ADn+de+la+Comisi%C3%B3n+del+Mapa+Geol%C3%B3gico+de+Espa%C3%B1a.+1-1-1874>
- Vidal LM (1917) Edad geológica de los lignitos de Selva y Binisalm (Mallorca) y descripción de algunas especies fósiles. Memorias de la Real Sociedad Española de Historia Natural 10 (7): 343–358. <http://bibdigital.rjb.csic.es/spa/Libro.php?Libro=1381> [first page of volume]
- Villa A, Villa GB (1841) Dispositio systematica conchyliarum terrestrium et Fluviatilium; conspectu abnormitatum novarumque specierum descriptionibus adjectis. Borroni et Scotti, Mediolani, 62 pp. https://books.google.at/books?id=_1ZOAAAAcAAJ&printsec=frontcover&hl=de&source=gbs_ge_summary_r&cad=0#v=onepage&q&f=false
- Villa A, Villa GB (1854) Due nuove melanie. Giornale di Malacologia 2 (8): 113.
- Villatte J (1952) Étude de la Faune campanienne saumâtre de Bélesta (Ariège). Bulletin de la Société d'Histoire Naturelle de Toulouse 8 (3–4): 315–339. <http://gallica.bnf.fr/ark:/12148/bpt6k6555410/f93.image.r=>
- Viquesnel A (1846) Journal d'un Voyage dans la Turquie d'Europe II. Mémoires de la Société Géologique de France, deuxième série 1 (2): 207–303. <http://www.biodiversitylibrary.org/item/133679#page/231/mode/1up>
- Voltz F (1852) Geologische Bilder aus dem Mainzer Becken. J. G. Wirth Sohn, Mainz, 88 pp. <https://download.digitale-sammlungen.de/pdf/1449821542bsb10284821.pdf>
- Walch JEI (1768) Die Naturgeschichte der Versteinerungen zur Erläuterung der Knorr'schen Sammlung von Merkwürdigkeiten der Natur. Des zweiten Bandes erster Abschnitt. Paul Jonathan Felßcker, Nürnberg, 184 pp. http://reader.digitale-sammlungen.de/de/fs1/object/display/bsb10213743_00005.html
- Walderdorff R (1864) Systematisches Verzeichniss der im Kreise Cattaro (Süd-Dalmatien) mit Ausnahme der Biela-Gora und in einigen angrenzenden Theilen von Montenegro und türkisch Albanien vorkommenden Land- und Süßwasser-Mollusken. Verhandlungen der kaiserlichen und königlichen zoologisch-botanischen Gesellschaft in Wien 14: 503–514. <http://www.biodiversitylibrary.org/item/55310#page/649/mode/1up>
- Watelet A (1851) [Des Recherches curieuses faites dans les sables tertiaires inférieurs des environs de Soissons, et donne la description de dix-sept espèces de coquilles inédites]. Bulletin

- de la Société archéologique, historique et scientifique de Soissons 5: 113–126. <http://gallica.bnf.fr/ark:/12148/bpt6k4539913/f110.item>
- Watelet A (1853) [Second mémoire sur les fossiles inédites du Soissonnais]. Bulletin de la Société archéologique, historique et scientifique de Soissons 7: 12–23. <http://gallica.bnf.fr/ark:/12148/bpt6k453993v/f11.image>
- Welter-Schultes FW (2012) European non-marine molluscs, a guide for species identification. Planet Poster Editions, Göttingen, 679 pp.
- Wenz W (1919a) Zur Nomenklatur tertiärer Land- und Süßwassergastropoden. I. Nachrichtenblatt der Deutschen Malakozoologischen Gesellschaft 51 (2): 68–76. <http://www.biodiversitylibrary.org/item/110094#page/82/mode/1up>
- Wenz W (1919b) Zur Nomenklatur tertiärer Land- und Süßwassergastropoden. II. Senckenbergiana 1: 63–67. <http://www.biodiversitylibrary.org/item/95236#page/75/mode/1up>
- Wenz W (1929) Fossilium Catalogus I: Animalia. Gastropoda extramarina tertiaria. W. Junk, Berlin, vol. IX: pp. 2503–2886.
- Wenz W (1928a) Zur Nomenklatur tertiärer Land- und Süßwassergastropoden. VIII. Senckenbergiana 10: 119–120.
- Wenz W (1928b) Zur Nomenklatur tertiärer Land- und Süßwassergastropoden. IX. Senckenbergiana 10: 219–220.
- Wenz W (1930) Zur Nomenklatur tertiärer Land- und Süßwassergastropoden. XI. Senckenbergiana 12: 64–66.
- Wenz W (1938–1944) Gastropoda. Teil 1: Allgemeiner Teil und Prosobranchia. In: Schindewolf OH (Ed.) Handbuch der Paläozoologie, Band 6. Verlag Gebrüder Bornträger, Berlin, 1–240 pp. (March 1938), 241–480 pp. (October 1938), 481–720 pp. (July 1939), 721–960 pp. (August 1940), 961–1200 pp. (October 1941), 1201–1506 pp. (October 1943), i–xii, 1507–1639 pp. (November 1944).
- Westerlund CA (1886) Fauna der in der paläarktischen Region (Europa, Kaukasien, Sibirien, Turan, Persien, Kurdistan, Armenien, Mesopotamien, Kleinasien, Syrien, Arabien, Egypten, Tripolis, Tunesien, Algerien und Marocco) lebenden Binnenconchylien. VI. Fam. Ampullariidae, Paludinidae, Hydrobiidae, Melaniidae, Valvatidae & Neritidae. Håkan Ohlsson, Lund, 156 + 13 pp. <http://www.biodiversitylibrary.org/item/41051#page/161/mode/1up>
- Westerlund CA (1892) Spicilegium Malacologicum. Neue Binnen-Conchylien in der paläarktischen Region. III. Nachrichtenblatt der Deutschen Malakozoologischen Gesellschaft 24 (11/12): 185–201. <http://www.biodiversitylibrary.org/item/53278#page/1047/mode/1up>
- Westerlund CA (1898) Novum specilegium malacologicum. Neue Binnenconchylien aus der paläarktischen Region. Ezhegodnik, Zoologicheskago Muzeya Imperatorskoi Akademii Nauk 3 (2): 155–183. <http://www.biodiversitylibrary.org/item/35243#page/261/mode/1up>
- White CA (1883) New molluscan forms from the Laramie and Green River groups, with discussion of some associated forms heretofore known. Proceedings of the United States National Museum 5: 94–99. doi: 10.5479/si.00963801.5-266.94
- Willmann R (1980) Die neogenen Süßwassergastropoden von Chersonisos (Kreta). Neues Jahrbuch für Geologie und Paläontologie, Abhandlungen 159 (2): 273–295.

- Willmann R (1981) Evolution, Systematik und stratigraphische Bedeutung der neogenen Süßwassergastropoden von Rhodos und Kos/Ägäis. *Palaeontographica Abt. A* 174: 10–235.
- Wood W (1828) Supplement to the Index testaceologicus; or A Catalogue of Shells, British and foreign. Privately published, London, 59 pp. + 8 pls. <http://www.biodiversitylibrary.org/item/74400#page/5/mode/1up>
- Woodward BB (1906) On the Dates of Publication of C. L. F. von Sandberger's "Die Land- und Süßwasser-Conchylien der Vorwelt", 4to, Wiesbaden (C. W. Kreidel), 1870–75. *Proceedings of the Malacological Society of London* 7 (1): 3–4. <http://www.biodiversitylibrary.org/item/53746#page/29/mode/1up>
- Yen JT-C (1958) Systematics and Distributions of *Pyrgulifera* Meek. *Annalen des Naturhistorischen Museums in Wien* 62: 193–209. http://www.zobodat.at/pdf/ANNA_62_0193-0209.pdf
- Yen T-C (1939) Die chinesischen Land- und Süßwasser-Gastropoden des Natur-Museums Senckenberg. *Abhandlungen der Senckenbergischen Naturforschenden Gesellschaft* 444: 1–234.
- Youluo (i.e. "the editorial group") (1978) Early Tertiary gastropod fossils from the coastal region of Bohai. Geological Publishing House, Nanjing, China, 6 + 157 pp. [In Chinese with English summary]
- Živković M (1893) Über das Tertiär des mittleren Timok-Beckens. *Annales Géologiques de la Péninsule Balkanique* 4 (1–2): 37–118, 147–168.